



# Prevalence of Non-Dermatophytic Molds Associated with Cutaneous Mycoses in Cattle in Abia and Imo States, Nigeria

C. N. Nwofor<sup>1</sup>, C. A. Oyeka<sup>2</sup>, N. E. Onyenwe<sup>3\*</sup>, M. O. Echeta<sup>4</sup>  
and Y. M. Tatteng<sup>5</sup>

<sup>1</sup>Department of Microbiology, Imo State University, Owerri, Nigeria.

<sup>2</sup>Department of Applied Microbiology, Nnamdi Azikiwe University, Awka, Anambra State, Nigeria.

<sup>3</sup>Department of Pharmaceutical Microbiology, College of Pharmacy, Igbinedion University Okada, Edo State, Nigeria.

<sup>4</sup>Department of Science Laboratory Technoogy, Umuagwo, Imo State, Nigeria.

<sup>5</sup>Department of Medical Laboratory Science, Niger Delta University, Willberforce Island, Bayelsa State, Nigeria.

## Authors' contributions

This work was carried out in collaboration among all authors. Author CNN designed the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Author CAO supervised the study. Author NEO proof read and managed the analyses of the study. Authors MOE and YMT managed the literature searches. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/ACRI/2021/v21i230232

Editor(s):

(1) Dr. Sung-Kun Kim, Northeastern State University, USA.

Reviewers:

(1) Agata Piecuch, University of Wroclaw, Poland.

(2) R. Meena Siddha, Central Research Institute, India.

Complete Peer review History: <http://www.sdiarticle4.com/review-history/68108>

Original Research Article

Received 10 March 2021

Accepted 14 May 2021

Published 04 June 2021

## ABSTRACT

**Aims:** To study the prevalence and identification of non-dermatophytic molds associated with cattle in Abia and Imo States, Nigeria.

**Study Design:** The systematic random sampling method was adopted.

**Place and Duration of Study:** A total of 2255 cattle was encountered, out of which 451 skin samples from both infected and asymptomatic animals were sampled from six cattle markets in Abia and Imo States, Nigeria between January and August, 2018.

**Methodology:** The 451 samples were analysed for their colonial and microscopic morphology and

\*Corresponding author: Email: o\_nathejik@yahoo.com, onyenwe.ejikeme@iuokada.edu.ng;

molecular analysis. Polymerase chain reaction of the 16SrRNA internal transcriber spacer (ITS) sequence techniques were carried out on the isolates. Pathogenicity of the isolates were determined.

**Results:** A total of 16 non-dermatophytic molds were obtained from 451 samples analysed at different frequency of occurrences from the cattle skins in both states and these include *Aspergillus fumigatus* (3.6%), *Aspergillus terreus* (2.7%), *Aspergillus wewitschiae* (13.5%), *Aspergillus flavus* (10.0%), *Aspergillus aculeatus* (9.0%), *Aspergillus sydowii* (5.0%), *Fusarium solani* (3.2%), *Fusarium lichenicola* (17.9%), *Fusarium succisae* (12.0%), *Fusarium oxysporum* (2.0%), *Penicillium citrinum* (3.0%), *Curvularia kusanol* (0.6%), *Cladosporium tenuissimum* (4.9%), *Pestalotiopsis microspora* (0.1%), *Talaromyces kendrickii* (0.1%) and *Absidia* specie (12.9%).

**Conclusion:** The study revealed that non-dermatophytic molds were highly prevalent and are possible causative agent of cutaneous mycoses in cattle. Pathogenicity evaluation carried out in this study revealed that *Aspergillus wewitschiae*, *Cladosporium tenuissimum* and *Absidia* sp. were highly virulent.

**Keywords:** Cattle; non-dermatophytic molds; cutaneous mycoses; Abia; Imo; fungi.

## 1. INTRODUCTION

Cattle is a major protein source in Nigeria and cattle rearing is one of the major occupations practiced by the fulanis in the northern part of Nigeria. They breed and nurture these animals till when they get matured and ready to be sold to the business men in other parts of the country. These business men make their supply to the cattle markets within the States, who later distributes them to smaller retailers who in turn sell to the masses, after butchering the animals. This process gives rise to a large business chain that is used to earn a source of livelihood. It can also be processed and used as a source of meat to prepare different delicacies within the country [1]. In the cause of breeding these animals, the fulani herdsmen move them from one geographical location to another within the bushes in search of grazing lands. This exposure therefore predisposes them to a lot of challenges.

Cutaneous mycosis in cattle is a worldwide zoonotic infection which are usually responsible for economic losses in the farm [2]. It is a highly contagious skin infection all over the world especially in the tropics (Africa, India) [3] and it is known to be caused by a dermatophyte *Trichophyton verrucosum* [4,5,6].

However, recent studies have shown that non-dermatophytic molds (NDMs); *Aspergillus* species, *Penicillium* species, *Fusarium* species, *Cladosporium* species, *Alternaria* species are fast replacing the dermatophytes in causing cutaneous mycoses [7,8,9]. These non-dermatophytic molds were known to be mere environmental contaminants [10]. They are also

known to be good enzymes secretors such as amylase, protease, cellulase, keratinase, lipases which can also act as a virulent factor [11].

The animal handlers are at high risk of this infections due to poor personal hygiene, since they are at close contact with these animals [12-13]. It can also be transmitted from animal to animal [12].

Therefore, considering the indispensable nature of cattles in Nigeria and the world at large, the highly contagious nature of cutaneous mycoses amongst animal communities have being reported as a public health challenge in Nigeria and the world at large, including its prolonged duration of treatment and mismanagement [14]. Therefore, it becomes evident that this study identifies these non-dermatophytic molds commonly associated with cutaneous mycoses in cattle in Abia and Imo States, Nigeria.

## 2. MATERIALS AND METHODS

### 2.1 Study Area

This study was carried out in the three major cattle markets in Abia and Imo States Southeastern, Nigeria respectively. Applying the systematic random sampling method [15], since it is a migrate population. a total of 2255 cattle were encountered from the six cattle markets in both States, dividing by the desired sample size which is 5, the total number of cattle skin sampled was 451. Samples were collected from both infected and asymptomatic animals. Only matured cows were found in the markets and mainly male cows were seen, since the females are usually left in the farms for procreation.

## 2.2 Method of Sample Collection

Samples of skin scrapings were collected from different parts of the cattle body especially areas suspected to be infected. These areas were cleaned with 70% disposable alcohol swab pad and allowed to evaporate. A sterile tooth brush was used to scrape the surface of the required areas of the cattle skin. These skin samples were collected in a sterile paper packet and placed in sterile bottles. Each collected sample was labeled based on location /zone where it was collected and date of collection. The sample containers were wrapped with aluminum foil and taken to the microbiology laboratory, Imo State University, Owerri, Nigeria and were processed within two hours.

## 2.3 Microscopic Examination of Samples

Some of the skin samples was examined by direct microscopy as described by Chesbrough [16], which were prepared with 10% KOH while the remaining were then cultured on the selected media.

## 2.4 Culture Technique

Each sample was cultured directly on the plates of Sabourauds dextrose agar (SDA) by spread plate method. The sterile brushes which was used for sample collection, was spread on the plates containing 20 mg chloramphenicol each to inhibit bacteria growth. This was later kept in the incubators at 37°C for 1-2 weeks. This was done on duplicate plate.

Discrete pure colonies of each isolates were inoculated on Sabourauds dextrose agar slants and incubated at 37°C for 1-2 weeks. The slants containing the pure cultures were stored in the refrigerator until required for further studies.

## 2.5 Identification of Isolates

The identification was based on growth rate, colonial and microscopic morphology. The isolates were identified on plates using standard methods as in Chesbrough (2010). The microscopic observations from slide culture of the isolates were compared with standard mycology atlas for identification [17]. Molecular analysis of the isolates was also carried on sixteen (16) isolates using sequencing the BigDye Terminator kit on a 3510 ABI sequencer by Inaqaba Biotechnological, Pretoria South Africa.

## 2.6 Pathogenicity Test

Twenty mice were bought from the department of Biochemistry, Imo State Polytechnic Umuagwo and checked for skin infection. The mice skin was cleaned with sterile water, fed for two weeks and allowed to adapt before introducing ten isolates that were recovered from cattle with lesion on them. The rats were paired and the cages were labeled with the name of the isolate for this investigation. A fragment of the isolate were diluted in 5ml of distilled water containing beads to break the fungal strands after which it was then rubbed on the skin of the mice with a sterile stick [18].

## 2.7 Method of Analysis of Data

The data from this study were analyzed statistically using multiple comparison in percentages and analysis of variance (ANOVA) as in Martins [19].

## 3. RESULTS AND DISCUSSION

Out of 2255 cattle encountered, 451 cattle skin were sampled of which 53(11.7%) had lesions which is an indication of fungal infections. The samples analyzed showed positive results on KOH and culture. Out of 451 cattle skin, 223 were sampled from Abia state of which 24(10.8%) had lesions while 228 were sampled from Imo state of which 29(12.7%) had lesion (Fig. 3). Among the sampled population all 451 (100%) were matured male cows ready to sell. Out of 31 skin sample sampled from the head non had lesion (0%), 58 were sampled from the leg and 10 (18.9%) had lesion, 15 from ear and 3(6%) had lesion, 16 from neck and 0(0%) had lesion, 8 were sampled from the groin and 6 (11.3%) had lesion, 285 were sampled from the abdomen and 21(40%) had lesion while 38 were sampled from the tail region and 7(13.2%) had lesion (Table 1). The amplified ITS fragments of the fungal isolates visualized under the blue transilluminator was shown in Fig. 1 (Lane 1- 11). Isolates from lanes 1,2, 4, 6-11 represents ITS bands at 600bp while lane 3 represents 500bp. L represents the 100bp molecular ladder. Fig. 2 shows results obtained from ITS sequence from the isolate which produced an exact match during the megablast search for highly similar sequences from the NCBI non reductant nucleotide (nr/nt) database. The ITS of the isolates showed 99-100% similarity to other species. The evolutionary distances which were computed using the Jukes-cantor method were in

agreement with the phylogenetic placement of ITS of C18 within the *Penicillium* sp. and revealed a closely relatedness to *Penicillium citrinum* (MH990629) other than any other *Penicillium* sp., C9 and C10 revealed a closely relatedness to *Aspergillus fumigatus* (MK461083), C14 and C16 to *Aspergillus terreus* (MK418744), C6 and C13 revealed a closely relatedness to *Aspergillus welwitschiae* (MG576117), C7, C11, C12 is related to *Aspergillus aculeatus* (MK461093), C3 is closely related to *Aspergillus flavus* (Mk 299130), C2 and C19 is related to *Fusarium succisae* (Mk 418691), C17 is closely related to *Aspergillus sydowii* (Mk 396475), C23 is related to *Talaromyceskendrickii* (kko 98055), C 25 closely related to *Curvulariakusanol* (MG975624), C15 is related to *Cladosporium tenuissimum* (MK 357638), C22 is related to *Pestalotiopsis microspora* (MK 224482), C24 was also found to be related to *Fusarium solani* (MH517359), C1 to *Fusarium lichenicola* (KH921661) and C20 was found closely related to *Fusarium oxysporum* (KM203578).

The DNA sequencing identified a total of 16 non dermatophytic molds species belonging to 8 genera; *Aspergillus* genera, *Talaromyces* genera, *Curvularia* genera, *Cladosporium* genera, *Pestalotiopsis* genera, *Fusarium* genera, *Penicillium* genera and *Absidia* genera (Table 2) were mostly isolated from the samples and the all occurred at different frequencies. In Abia State, out of 223 cattle skin sampled a total of 12 non-dermatophytic molds species were recovered which belongs to *Penicillium citrinum* (4%), *Aspergillus fumigatus* (4.7%), *Aspergillus terreus* (2%), *Aspergillus welwitschiae* (15%), *Aspergillus flavus* (11.1%), *Aspergillus aculeatus* (7.3%), *Aspergillus sydowii* (3.6%), *Fusarium solani* (2.2%), *Cladosporium tenuissimum* (4.5%), *Fusarium lichenicola* (19.4%), *Fusarium succisae* (12.5%) and *Absidia* specie (14%). Based on the result, *Fusarium lichenicola* was the most frequently isolated followed by *Aspergillus welwitschiae*, *Absidia* specie while the least isolated were *Aspergillus terreus* and *Fusarium solani*. Among these isolates, ten were recovered from the 24 samples that had lesion within the State and they include *Penicillium citrinum* (5%), *Aspergillus fumigatus* (2%), *Aspergillus welwitschiae* (21.3%), *Aspergillus aculeatus* (5%), *Aspergillus flavus* (5%), *Aspergillus sydowii* (3.3%), *Cladosporium tenuissimum* (7%), *Fusarium lichenicola* (13.1%), *Fusarium succisae* (20%) and *Absidia* specie (20%) (Tables 3, 4). Based on the results, *Aspergillus welwitschiae*, *Absidia* specie and

*Fusarium succisae* were the most prevalent non-dermatophytic mold recovered. In Imo State, out of 228 cattle skin sampled a total of 16 non-dermatophytic molds were recovered and the include *Penicillium citrinum* (1.5%), *Aspergillus fumigatus* (2.1%), *Aspergillus terreus* (4%), *Aspergillus welwitschiae* (12%), *Aspergillus flavus* (8%), *Aspergillus aculeatus* (11%), *Aspergillus sydowii* (7.1%), *Talaromyceskendrickii* (0.3%), *Curvulariakusanol* (1.5%), *Cladosporium tenuissimum* (5.5%), *Pestalotiopsis microspora* (0.3%), *Fusarium solani* (4.6%), *Fusarium lichenicola* (16%), *Fusarium succisae* (10.4%), *Fusarium oxysporum* (4.6%) and *Absidia* specie (11.3%). The result showed that *Fusarium lichenicola* was the most frequently isolated, followed by *Aspergillus welwitschiae* and *Absidia* specie while the least isolated were *Talaromyceskendrickii* and *Pestalotiopsis microspora*. Of these non-dermatophytic molds recovered, ten were among those recovered from the 28 that had lesion within the State and the include *Aspergillus welwitschiae* (28.2%), *Aspergillus aculeatus* (15.5%), *Aspergillus flavus* (1.4%), *Aspergillus sydowii* (8.5%), *Talaromyceskendrickii* (1.4%), *Cladosporium tenuissimum* (5.6%), *Pestalotiopsis microspora* (1.4%), *Fusarium lichenicola* (14.1%), *Fusarium succisae* (9.9%) and *Absidia* specie (14.1%) (Tables 3, 4). Out of the twelve isolates recovered from the cattle skins that had lesions from both States, ten were subjected to pathogenicity test, only three *Aspergillus welwitschiae*, *Cladosporium tenuissimum* and *Absidia* specie were able to elicit clinical lesions suggestive of cutaneous mycoses.

Based on the analysis shown in this study, it has revealed a high prevalence of non-dermatophytic molds associated with cattle skin (including those with lesion and those without lesion) in Abia and Imo States, Nigeria. Infection was mostly recorded from the abdomen than the other anatomical sites. This could be attributed to the fact that these animals lay and roll on the soil with their abdomen than other parts of their body, thereby predisposing their glabrous skin to the contaminated soil. These animals, often lay side by side to each other, thereby coming in contact with other infected animals in the market. Some of the cattle due have trauma on their glabrous skin due to the stacking effect they suffer on transit in the trucks while coming down to the East from the North. Also, trauma incurred in the bushes during grazing, paves way for these non-dermatophytic molds to gain access through the

skin. This agrees with work of [20]. In their work they stated that lesions were mainly observed on the animals (goats and sheep) glabrous skin. Also, [21] in their work also reported that lesions were seen more on the limbs (18.7%) of horses than other parts of the body where samples were collected (tail, head and abdomen), when they analyzed the prevalence and distribution of dermatophytes among domestic horses in Kwara State, Nigeria.

It is also important to note that only matured male cows were encountered in this study and as such age and sex variables were eliminated. This was attributed to the fact that only ready to sell cows (matured cow) were found in the markets. For the sex, according to the veterinary doctors, the female cattle were hardly sold out by the farmers as they are usually left in the farms for procreation. The sellers are also business men who buy commodities that they can sell and make profit. According to them buyers do prefer male cows than the female cows. They attributed these to the fact that the male cows are easy to prepare as meat for food than the female cows, so they rarely purchase them. Among the non-dermatophytic molds isolated from this study, *Aspergillus* species 338(43.6%) were the most frequently isolated followed by *Fusarium* species 269(34.7%) then *Absidia* specie 100(12.9%), *Cladosporium* specie 38(4.9%), *Penicillium*

specie 23(3.0%), *Curvularia* specie 5(0.6%). *Pestalotiopsis* specie 1(0.1%) and *Talaromyces* specie 1(0.1%) from both States. This can be attributed to the fact that most of these isolates have been reported to be major soil and vegetation inhabitants [22,23,24]. Since these cows lay mostly on these soils, there is a high tendency of these molds colonizing the animal skin and also considering that they were usually taken into the bushes for grazing, there is a high tendency of picking some from the grasses. Similar findings have been revealed, though not in cattle by [25] that they isolated *Aspergillus* species as the most prevalent followed by *Penicillium* sp., *Alternaria* sp., and *Fusarium* specie. when he examined the mycological skin swabs from dog skin. Also, [26] isolated saprophytes in the frequency order of *Aspergillus*, *Penicillium*, *Alternaria* species when he sampled hair from sheep and goat in Libya. [27], further more in his study, He reported that they isolated *Aspergillus* specie (37.9%) as the highest saprobe followed by *Penicillium* specie (21.4%). *Cladosporium* specie (8.7%), *Fusarium* species (7.8%), *Curvularia* specie (3.9%) when he surveyed saprophytic fungi from dogs and cats in the city of Fortaleza, Brazil. Another researcher [28] in his work recovered *Curvularia* specie from skin infection in goat in Cauvery delta region of Tamil Nadu, India.

**Table 1. Distribution of samples collected based on anatomical sites including sites infected with lesions**

Anatomical sites	Total samples collected from anatomical sites	Total samples collected from anatomical sites with lesion	% occurrence of anatomical sites with lesions
Head	31	0	0.0
Leg	58	10	18.9
Ear	15	3	6.0
Neck	16	0	0.0
Groin	8	6	11.3
Abdomen	285	27	51.0
Tail	38	7	13.2
Total	451	53	

**Table 2. Frequency of occurrence of Non- dermatophytic mold genera isolated from Abia and Imo states Nigeria**

Genera	Total occurrence	Percentage occurrence (%)
<i>Aspergillus</i> species	338	43.6
<i>Fusarium</i> species	269	34.7
<i>Penicillium</i> species	23	3.0
<i>Cladosporium</i> species	38	4.9
<i>Curvularia</i> species	5	0.6
<i>Pestalotiopsis</i> species	1	0.1
<i>Talaromyces</i> species	1	0.1
<i>Absidia</i> species	100	12.9
Total	775	

**Table 3. Frequency of occurrence of non-dermatophytic molds isolated from cattle skin (symptomatic and asymptomatic) in Abia and Imo states Nigeria**

Species of non-dermatophytic Molds	Number of occurrence		Total occurrence (%)
	Abialmo		
<i>Penicilliumcitrinum</i>	18	5	23 (3.0)
<i>Aspergillus fumigatus</i>	21	7	28 (3.6)
<i>Aspergillus terreus</i>	8	13	21(2.7)
<i>Aspergillus welwitschiae</i>	67	38	105 (13.5)
<i>Aspergillus flavus</i>	50	26	76 (10.0)
<i>Aspergillus aculeatus</i>	33	36	69 (9.0)
<i>Aspergillus sydowii</i>	16	23	39 (5.0)
<i>Talaromyceskendrickii</i>	0	1	1(0.1)
<i>Curvulariakusanol</i>	0	5	5(0.6)
<i>Cladosporium tenuissinum</i>	20	18	38 (4.9)
<i>Pestalotiopsismicrospora</i>	0	1	1 (0.1)
<i>Fusarium solani</i>	10	15	25 (3.2)
<i>Fusarium linchenicola</i>	87	52	139 (17.9)
<i>Fusarium succisae</i>	56	34	90 (12.0)
<i>Fusarium oxysporum</i>	0	15	15 (2.0)
<i>Absidia specie</i>	63	37	100 (12.9)
Total	449	326	775

**Table 4. Frequency of occurrence of non-dermatophytic molds isolated from cattle skin with lesion (symtomatic) in Abia and Imo states Nigeria**

Species of non-dermatophytic molds (%)	Frequency of occurrence	
	Abia State	Imo State
<i>Penicilliumcitrinum</i>	3 (5.0)	0
<i>Aspergillus fumigatus</i>	1 (2.0)	0
<i>Aspergillus welwitschiae</i>	13 (21.3)	20 (28.2)
<i>Aspergillus aculeatus</i>	3 (5.0)	11 (15.5)
<i>Aspergillus flavus</i>	3 (5.0)	1 (1.4)
<i>Aspergillus sydowii</i>	2 (3.3)	6 (8.5)
<i>Cladosporium tenuissinum</i>	4 (7.0)	4 (5.6)
<i>Talaromyceskendrickii</i>	0	1 (1.4)
<i>Pestalotiopsismicrospora</i>	0	1 (1.4)

Species of non-dermatophytic molds (%)	Frequency of occurrence	
	Abia State	Imo State
<i>Fusarium linchenicola</i>	8 (13.1)	10 (14.1)
<i>Fusarium succisae</i>	12 (20.0)	7 (9.9)
<i>Absidia</i> specie	12 (20.0)	10 (14.1)
Total	61	71




Fig. 1. Agarose gel electrophoresis showing the amplified ITS fragment of the fungal isolates. Lanes 1,2,4 and 6-11 represent the ITS bands at 600bp, lane 3 represents 500bp bands, while lane L, represents the 100bp molecular ladder





Fig. 2. The Phylogenetic tree from the internal transcriber spacer (ITS) obtained from the isolate in this study





**Fig. 3. Shows population of cattle sampled (with or without lesion) in Abia and Imo**

In this present study *Aspergillus* spp., *Fusarium* spp., *Absidia* spp., *Cladosporium* spp. and *Penicillium* spp. were the most frequently isolated non-dermatophytic mold in this study; this is not surprising as these molds have been reported to be highly ubiquitous in nature [29, 30,31,32,33]. A number of skin infections in man and animal has also been linked to them especially in immunocompromised host and onychomycosis [34,35,36,37].

In this study, there were no significant difference between the non-dermatophytic molds recovered from both States, although they were recovered at different frequencies. This is understandable since these animals were being produced in the same area in the Northern part of the country. Four non-dermatophytic molds recovered from Imo State were not recovered from Abia State which includes *Curvularia kusanol* 5(1.5%), *Pestalotiopsis microsporum* 1(0.1%), *Talaromyces kendrickii*1(0.3%) and *Fusarium oxysporum*15(4.6%). Judging by their low frequency of occurrence there is a high possibility that these molds were picked up by these cows on transit or within the environment of the cattle market.

In this study, it was also observed that similar non-dermatophytic molds recovered from asymptomatic cows were also recovered from those cows with lesions. This strengthens the

notion that these non-dermatophytic molds naturally colonize the cattle skin but if by coincidence the immunity of the host is lowered, they can take advantage by causing cutaneous infection. Also, the possibility of some of these molds being categorized as true pathogens cannot be underestimated. This is also in accordance with other published works by [38], their work analyzed the current status of *Fusarium* infection in animal and human and concluded that this filamentous fungus is widely distributed in the soil and on plants and can cause superficial infections especially on compromise host. [39] in their study, also agrees that *Cladosporium*spp. are opportunistic pathogens that are capable of causing cutaneous infections. In another study by [40] and [2], they also observed that most mold floras are known to cause superficial or cutaneous infections due to its ability to utilize keratin.

Among the twelve non-dermatophytic molds isolated from the cattles with lesion in both States, 10 of them were subjected to pathogenicity test and only *Aspergillus welwitschiae*, *Cladosporium tenuissimum* and *Absidia* specie were able to elicit alopecia, nodule and discoloration on healthy albino mice skin respectively. These observations were clinical symptoms suggestive of cutaneous mycoses. Thus, this observation therefore establishes that the non-dermatophytic molds

observed as mere opportunistic pathogens may actually be the -true pathogens. Previous works have also implicated the non-dermatophytic molds in causing cutaneous skin infections. Reports by [41] in their work revealed that loss of fur is an indication of lesion production. Also, [24] also demonstrated that *Cladosporium* species were capable of causing localized superficial infections. Also considering that these molds are known to be good producers of industrial enzymes such as keratinase, protease, lipase, amylase [42,43,44]. This can obviously explain their capability to degrade skin tissues as these enzymes can also act as virulent factors to cutaneous mycoses [45].

#### 4. CONCLUSION

This study also indicates that irrespective of the well-established advantages of the cattle fur, it acts as a potential receptacle to these molds and as such makes it easier for the molds to gain access into the skin once it is traumatized. It cannot also be overemphasized that this association between these non-dermatophytic molds and the animal might encourage a parasitic relationship between them which might invariably affect the size of the cow. This present study also revealed that movement of cattle from one geographical area to another might encourage the dispersal and picking up of new non-dermatophytic molds by these animals and nomads. These study therefore suggests that government should encourage farmers to keep these animals in well constructed barns, whereby their forages can be brought to them instead of moving them about. It is also important that they should be kept in columns in trucks while transporting them to reduce the tendency of incurring trauma on their skins and moreso regular checks by veterinary doctors should be ensured as this will go a long way in controlling the colonization of the animal skin by these non-dermatophytic molds. There is also need for those taking care of these animals to observe precautionary measures such as improving on their personal hygiene, wearing of protective kits while taking care of these animals as some of these fungal spores can easily be picked up ignorantly from them since majority of the cows are asymptomatic carriers. Pathogenicity tests carried out also showed a high possibility of some of these non-dermatophytic molds infecting human skin. Further research will be carried out to screen some of these non-dermatophytic molds for enzyme production.

#### ETHICAL APPROVAL

Based on the compliance of the ethical standard of Imo and Abia State in Nigeria , the Ethical permit No; NAR/Vet/XX of the Ministry of Agriculture and Natural Resources Owerri, Imo state dated 13th March ,2017 and that of the Ministry of Agriculture Umuahia, Abia state Ref. No; DVS/01/RCH/01/18, dated 4th June,2018, was also issued before the commencement of the screening of the cattles in both States.

#### ACKNOWLEDGEMENTS

We sincerely acknowledge the help of the Veterinary doctors in charge of the cattle market in both Abia and Imo state. The entire staff of the College of Pharmacy, Igbiniedion University Okada for their Assistance offered to this research in their Laboratory. Also my gratitude goes to Prof. Tetteng of Niger Delta University for his assistance in the Biotechnology analysis of this research.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Philip T. Live stockproduction: recent trends, future prospects. Philos Trans R Soc Lond B Biol Sci. 2010; 365(1554):2853-2867.
2. Didier P. Non-dermatophytes dermatoses mimicking dermatophytoses in animals. Mycopathologia. 2017;18:113-126.
3. Modupeola OS, Kabiru A. Cutaneous manifestation of Deep mycosis: An experience in a Tropical pathology laboratory. Indian J Dermatol. 2011;56(3): 282-286.
4. Akbarmehr J. The prevalence of cattle ringworm in native dairy farms of sarab city (East Azarbayjan Province), Iran. African Journal of Microbiology Research. 2011; 5(11):1268-1271.
5. El-diasty EM, Ahmed MA, Okasha N, Mansour SF, El-dek SL, Abd El-Khalek HM, Youssif MH. Antifungal activity of zinc oxide nanoparticles against dermatophytic lesions of cattle. Romanian Journal Biophys. 2013;23(3):191-202.
6. Agnetti F, Righi C, Soccia E, Felici A, Crotti S., Moretta L, Morretti A, Maresca C, Troiani L and Pappini M. *Trichophyton vernicosum* Infection in cattle farms of

- Umbria (Central Italy) and transmission to humans. *Mycosis*. 2014;57(7):400-405.
7. Vyas A, Pathan N, Sharma R, Vyas LA. Clinicomycological study of cutaneous mycoses in SawaiManSingh Hospital of Jaipur, North India. *Annals of Medical Health Science Research*. 2013;3:593-597.
  8. Pietro N, Constanze K, Gabriela G and Hans-jurgen, T. Mycology – an update. Part 1: Dermatophytes. Causative agents, epidemiology and pathogenesis. *Journal of the German Society of dermatology*. 2014;12(3):188-210.
  9. Ravinder KP, Swagatika P, Sahanawaj K. Non dermatophyte molds causing onychomycosis: a rising trend in North India. *International Journal of Community Medicine and Public Health*. 2017;4(12): 4532-4537.
  10. Tampieri, MP. Update on the diagnosis of dermatomycosis. *Parassitologia*. 2004; 46(1-2):183-186.
  11. Luis HG, Simone CP, Michele M, Ana CSR, Valeria CS, Fubiana FZ, Ana Carla MM.A, Altino BJ, Maria de Lourdes TMP. Screening of filamentous fungi for production of enzyme of biotechnological interest. *Brazilian Journal of Microbiology*. 2006;37:474-480.
  12. Songer JG and Post KW. *Veterinary Microbiology. Bacteria and Fungal Agents of Animal Disease*. Published by Elsevier Saunders. 2005;360-369.
  13. Zarei M. and Baharea I. Prevalence of cutaneous mycoses among workers. *Turk. J. Med. Sci*. 2011;41(2):291-294.
  14. Nweze EI, Eke I. Dermatophytosis in northern Africa. *Mycoses*. 2016;59(3):137-144.
  15. Cochran WG. *Sampling Techniques*, Wiley, New York. 3<sup>rd</sup> Edition. 1977;205-212 published in Canada.
  16. Cheesbrough M. *District Laboratory Practice in Tropical Countries*. Part 2, second edition, update. Tropical Health Technology/ Butterworths and Co. Ltd. Cambridge/ sevanaks; 2010.
  17. Lambert WC. ed. *Problem in dermatologic and diagnosis and management*. American Academic of Dermatology Inc. Evaton, Illinois. 1988;311.
  18. McEvoy JM, Nde CW, Sherwood JS, Logue CM. An evaluation of sampling methods for the detection of *Echerichia coli* and *Salmonella* on turkey carcasses. *Journal of Food Protection*. 2005;68:34-39.
  19. Martins OO and Igwemma AA. *Applied Statistical Techniques for Business and Basic Sciences*. 2<sup>nd</sup>. Skillmark media Ltd; Owerri, Nigeria. 2000;216-304.
  20. Emenuga VN, Oyeka CA. Epidemiology, Health Effects and Treatment of Cutaneous Mycoses of Goat and Sheep from some Eastern States of Nigeria. *American Journal of Infectious Diseases and Microbiology*. 2013;1(16):106 -110.
  21. Balogun RB, Jegede HO, Jibril A, Kwanashie CN, Kazeem HM. Prevalence and distribution of dermatophytes among domestic horses in Kwara State, Nigeria. *Sokoto Journal of Veterinary Sciences*. 2017;15(2):1-6.
  22. Deshmukh SK. Incidence of keratinophilic fungi from selected soils Kerala State (India). *Mycopathologia*. 2002;156(3): 177-181.
  23. Plancastell C, Ghidini F, Donotrio G, Jottini S, Taddal S, Cavirani S, Cabassi CS. Isolation and characterization of a strain *Lichtheimiacyrmbifera* (ex *Absidiacyrmbifera*) from a case of bovine abortion. *Reproduction Biology and Endocrinology*. 2009;30(7):138.
  24. Marcelo SD, Deanna AS, Adela MV, Jose FCL, Nathan WJ and Josepa G. Cladosporium species recovered from clinical samples in the united states. *Journal of clinical Microbiology*. 2015;53(9):2990-3000.
  25. Igor S, Sandra J, ProdanovJasna Z. Presence and importance of saprophyte fungal organisms on dog Skin. *Zbornik Matrice Srpske Za PrirodneNauke*. 2007;113(261):261-265.
  26. El-said, AHM., Sohair, THand El-Hadi., AG. Fungi associated with hairs of Goat and Sheep in Libya. *Mycobiology*. 2009; 37(2):82-88.
  27. Paixao GC, Sidrim JJC, Campos GMM, Brilhante RSN and Rocha MEG. Dermatophyte, and Saprobe Fungi collected from dogs and cats in the city of Fortaleza, Brazil. *Arqyivo Brasileiro de Medicina Veterinariaootecnia*. 2001;53(5): 568-573.
  28. Ponnusamy P, Saravanan M, Lurthu T. Rectha, Ronald BSM, Marickam R, Puvarajan R. *Curvularia* dermatomycosis in a goat: TamilNadu, India. *Journal of Entomology and Zoology Studies*. 2018; 6(3):1446-1448.
  29. Moallaei H, Zaini F, Pihet M, Mahmoudi M, Hsahemi J. Isolation of keratinophilic fungi

- from soils samples of forests and farm yards. Iran Journal of Public Health. 2006;35:62-69.
30. Farid MT, Nareen QF. Isolation, Identification and Seasonal Distribution of soil borne fungi in different areas of Erbil Governorate. Journal of Advanced Laboratory Research in Biology. 2012; 3(4): 246-255.
  31. Malek E, Moosazadeh M, Hanafi P, Abbasi NZ, Amini A. Isolation of keratinophilic fungi and aerobic Actinomycetes from park soils in Gorgan, North of Iran. Jundishapur Journal of Microbiology. 2013;6(10):11250.
  32. Nostratabadi M, Kordbacheh P, Kachueri R, Safara M, Rezala S and Afshan MA, Isolation of keratinophilic fungi from the soil of Agareater Tunb, Abu-Musa and Sirri Islands in the Persian Gulf, Iran. Current Medical Mycology. 2017;3(2):13-19.
  33. Tahereh R, Rasoul M. Identification of dermatophyte and nondermatophyte molds isolated from animal lesions suspected to dermatomycoses. Advanced Biomedical Research. 2020;9:4.
  34. Marjan M, Zeinab G, Mohammed RS, Leila H, Hossein K, Kamiar Z, Hossein M. Growing incidence of non-dermatophyte onychomycosis in Tehran Iran. Jundishapur Journal of Microbiology. 2016;9(8):e40543.
  35. Shayla EH, Paul ML, John HB and Adrian MZ. *Penicillium citrinum*: opportunistic pathogen or idle bystander? A case analysis with demonstration of galactomannan cross reactivity. Medical mycology case reports. 2017;17:8-10.
  36. Didier P. Non-dermatophytes dermatoses mimicking dermatophytoses in animals. Mycopathologia. 2017;182:113-126.
  37. Charu J, Deepti R, Ravinder K. Non-dermatophyte molds; Not so uncommon agents of onychomycosis. International Journal of Scientific Research. 2017; 6(11):204.
  38. Jain PK, Gupta VK, Misra AK, Gaur R, Bajpai V, Issar S. Current status of *Fusarium* infection in Human and Animal. Asian Journal of animal and veterinary Advances. 2011;6(3):201-227.
  39. Suzana T, Natasha MT. Cladosporium spp: cause of opportunistic mycoses. Acta Fac Med Naiss. 2007;24(1):15-19.
  40. Bakheshwain S, Khizzi NEI, Al-Rasheed AM., Al-Ajlan HH. and Parvez S. Isolation of opportunistic fungi from Dermatophytic samples. Asian Journal of Dermatology, 2011;(3):13-19.
  41. Subha G, ParveezAhnad P and Parveen KP. Case study: Investigation of dermatophytosis from the skin scrapings collected from cow. International Journal of current Microbiology and Applied Sciences. 2017;6(3):640-643. [43]
  42. Gropp K, Schild L, Schinder S, Hube B, Ziptel PE. and Skerka C. The yeast *Candida albicans* evades human complement attack by secretion of aspartic proteases. Molecular Immunology. 2009; 47(2):405-475.
  43. He J, Thomas K and Leif JJ. Enzyme production by filamentous fungi: analysis of the secretome of *Trichoderma reesei* grown on unconventional carbon source. Microbial cell factories. 2011; 10:68.
  44. Sunitha VH, Ramesha A, Savitha J, Srinivas C. Amylase production by endophytic fungi *Cylindrocephalum* sp isolated from medicinal plant *Alpinia calcarata* (Haw) Roscoe. Brazilian Journal of Microbiology. 2012;43(3):1213-1221.
  45. NaluTeixera de AP, Fernando Cristina AM, Antonio R and Nila Maria MR. Dermatophytes: host-pathogen interaction and antifungal resistance. Anais Brasileiros de Dermatologia. 2010;85(5): 657-667.

## APPENDIX

Examples of some lesions on different parts of cattle skin encountered during this research work





© 2021 Nwofor et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
*The peer review history for this paper can be accessed here:*  
<http://www.sdiarticle4.com/review-history/68108>