



# Impact of Foliar Boron and Fertilizer Management Systems on Growth, Yield and Quality of Brinjal

Bhagchand Yadav<sup>a\*</sup>, Anil Kumar Soni<sup>a</sup>,  
Atma Ram Meena<sup>a</sup>, Mahendra Jakhar<sup>b</sup> and Suman Meena<sup>a</sup>

<sup>a</sup> Department of Horticulture, S.K.N. College of Agriculture, Jobner S.K.N. Agriculture University, Jobner, Jaipur, Rajasthan, India.

<sup>b</sup> Department of Horticulture, College of Agriculture, SKRAU, Bikaner, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/IJECC/2023/v13i92366

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/97905>

Original Research Article

Received: 22/01/2023

Accepted: 24/03/2023

Published: 18/07/2023

## ABSTRACT

A field experiment was conducted during the rainy season of 2017–18 at Horticulture Farm, S.K.N. Agriculture University, Jobner, Jaipur, Rajasthan to study the role of foliar spray of boron on the growth, yield, and quality of Brinjal (*Solanum melongena* L.) The study was laid with 18 treatments (Three replicates of six treatments) viz., INM levels (control, 100% inorganic NPK, 75% inorganic NPK +25%VC (vermicompost), 50% inorganic NPK +50%VC, 25% inorganic NPK +75% VC, and 100%VC and boron levels (control, 100 ppm, and 200 ppm). It was observed that there was a significant ( $P < 0.05$ ) increase in the plant height and plant spread at 30, 60, and 90 DAT on the application of 75% inorganic NPK +25%VC on the brinjal crop, while the application of inorganic 50% NPK +50%VC significantly ( $P < 0.05$ ) increased chlorophyll content ( $\text{mg.g}^{-1}$ ), fruit length (cm), fruit weight (g), fruit yield  $\text{ha}^{-1}$  (q), Nitrogen and Phosphorus content in fruit (%), as compared to control but statistically at par with 25% inorganic NPK +75%VC. The application of boron (200ppm)

\*Corresponding author: E-mail: bcyadavhorti@gmail.com;

significantly ( $P < 0.05$ ) increased the plant height and plant spread at 30, 60, and 90 DAT(cm), chlorophyll content (mg/g), fruit length (cm), fruit weight (g), fruit yield hectare<sup>-1</sup>(q), NP content in fruit (%), as compared to control and boron (100 ppm).

**Keywords:** Brinjal; growth; NPK; quality; vermicompost.

## 1. INTRODUCTION

“Brinjal (*Solanum melongena* L.) which is commonly known as eggplant or aubergine belongs to the family *Solanaceae*. It is believed that brinjal is native to southern India and is one of the most popular vegetables grown throughout the tropics and subtropics” [1]. “Its green leaves are the main source of anti-ascorbic acid (vitamin C). Brinjal contains phytonutrients with antioxidant activity. They have anti-cancer, antiviral, antimicrobial, anti-inflammatory, laxative, antidiabetic, and anti-hypercholesterolemic activities. Eggplant fruits contain a phenolic compound, the most abundant source of chlorogenic acid” [2]. It has medicinal properties and is used as an appetizer, aphrodisiac, and “cardiotonic” in Ayurveda, and also used for the treatment of bronchitis, asthma, and dysentery. It is rich in dietary fiber, potassium, calcium, manganese, copper, and vitamins A & B [3].

“Brinjal being a long-duration crop requires good manure and fertilizer” [4]. Nitrogen is considered a building block in protein synthesis and is the main ingredient of protoplasm (Paul et al. 2017). Sat and Saimbhi [5] reported that increasing the nitrogen considerably can delay the flowering of eggplant and the time requirement for fruit setting increased significantly. “Phosphorus plays a key role in the formation of the plasma membrane, nucleic acids, coenzymes, organic molecules, many phosphorylated products, carbohydrates synthesis, and nutrient contents like Ca, Mg, N, K, and S” [6]. Crop productivity can be significantly increased with potassium use, as it imparts increased disease resistance. It regulates water balance within the plant cell through the process of anabolism, respiration, and transpiration and in turn, reduces the tendency to wilt. It improves the taste and quality of tomatoes through the formation of protein and chlorophyll by using available water.

“The technique of organic vegetable production plays an important role in producing high-value vegetables” [7]. “Vermicompost consistently encourages biological activity which can cause the plant to germinate seeds, produce a flower,

and provide a better yield than in commercially used media, independent of nutrient availability” [8]. “Vermicompost increases the surface area and provides strong absorbability of nutrients, and retains more nutrients for a longer period” [9]. “Vermicompost is slowly releasing organic manure which has most of the macro and micronutrients chelated from and fulfill the plant nutrient requirement of plant for a longer period, its use has been advocated in integrated nutrient management (INM) system in vegetable crops. Vermicomposts are characterized by high porosity, water-holding capacity, drainage, aeration, and microbial activity” [4].

“Boron (B) is a micronutrient critical to the growth, health, and carbohydrate metabolism of all crops. It is a component of the plant cell wall and reproductive structures. It is a mobile nutrient in the soil, meaning it is prone to movement in the soil. Boron role plays a diverse range of plant functions including maintenance of structural and functional integrity of biological membranes, cell wall formation and stability, movement of sugar, seed set, and pollination. In plants deficiency of boron nutrients results into delaying in pollen germination, pollen tube development, and halts flowering and fruit setting” [10].

## 2. MATERIALS AND METHODS

The field experiment was conducted at Horticulture farm, Department of Horticulture, S.K.N. COA, Jobner, Rajasthan, India from July to December 2017. The details regarding the use of experimental techniques, materials, and criteria opted for the assessment of treatments during the investigation are provided.

The soil texture in the experimental field was loamy sand, slightly alkaline, poor in organic carbon with low nitrogen (135.05 kg.ha<sup>-1</sup>), phosphorus (16.67 kg.ha<sup>-1</sup>), and medium in potassium contents (150.83 kg.ha<sup>-1</sup>).

In the experiment, Pant Rituraj variety of brinjal was used. Treatments were considered as six levels of INM & Vermicompost, viz., N<sub>0</sub>= Control, N<sub>1</sub>=100% RD of inorganic NPK, N<sub>2</sub>=75% RD of

inorganic NPK +25% through VC,  $N_3=50\%$  RD of inorganic NPK +50% through VC,  $N_4=25\%$  RD of inorganic NPK +75% through VC,  $N_5=100\%$  RD of NPK through vermicompost and three levels of boron viz.,  $B_0=$  Control,  $B_1=$  100 ppm,  $B_2=$  200 ppm.

The experiment was laid out in an RBD (Randomized Block Design) with three replications. Thirty days old seedlings were transplanted at the spacing of  $60 \times 45 \text{ cm}^2$  in the rainy season, 2017. The size of the experimental plots was  $2.4 \times 2.25 \text{ m}^2$ . Harvesting was carried out in December 2017. Data were collected from five randomly selected plants for each plot and the recorded parameters were plant height (cm), plant spread ( $\text{cm}^2$ ) for both directions (E-W & N-S), chlorophyll contents in leaves (mg/g), fruit length (cm), average fruit weight (g), Yield per hectare (q), nitrogen and phosphorus contents in fruit (%).

### 3. RESULTS AND DISCUSSION

#### 3.1 Plant Height

The significant maxi plant height at 30, 60, and 90 days after transplanting (19.15, 51.31, and 72.56 cm) was observed under the treatment  $N_2$ , which was statistically at par with the application of treatment  $N_3$  because inorganic fertilizer has more quantity of macronutrients that provide better vegetative growth. Nitrogen, phosphorous, and potassium are considered major nutrients required for the proper growth and development of the plant. This might be due to the better nutritional environment in the root zone for the growth and development of the plant by the application of NPK with vermicompost. Potassium fertilization improved overall crop growth in terms of plant height and number of branches per plant.

Whereas, minimum plant height at 30, 60, and 90 days after transplanting (12.80, 40.30, and 58.01 cm) were observed. The maximum plant height at 30, 60, and 90 days after transplanting (17.78, 49.68, and 69.98 cm) was recorded with 200 ppm, and minimum plant height at 30, 60, and 90 days after transplanting (15.66, 43.25, and 61.13 cm) was observed with  $B_0$ . "Growth attributes of brinjal increased with the application of boron at the initiation of flowering. Boron is also associated with the development of cell walls and results in improved shoot growth [11] and thus increased plant height, branches, and leaves per

plant [12] which greatly confirmed the present results".

#### 3.2 Plant Spread (E-W)

The maximum plant spread (East-West) at 30, 60, and 90 DAT (30.65, 56.22, and 81.29 cm) were observed under treatment  $N_2$ , which was statistically at par with treatment  $N_3$ . While, minimum plant spread (E-W) at 30, 60, and 90 DAT (22.45, 44.06, and 68.50 cm) was observed under control. "The plant spread (East-West) at 30, 60, and 90 DAT under the treatment  $N_2$  was registered at 36.52, 27.59, and 18.67% higher, respectively. Nitrogen is a main constituent of protoplasm, cell nucleus, amino acids, proteins, chlorophyll, and many other metabolic products. Phosphorus is a constituent of nucleic acid, phytin, and phospholipids. The beneficial influence of phosphorus in the early stages of growth may be explained by early stimulation of the root system through efficient translocation to the root or certain growth stimulation compounds formed on account of the protoplasmic activity of tops in phosphorus-fed plants when enhanced absorption of nitrogen and other nutrients and their utilization" (Parihar and Tripathi, 2003). Application of boron significantly ( $P < 0.05$ ) influenced the plant spread (E-W) at 30, 60, and 90 DAT (29.33, 54.28, and 79.90) in 200 ppm and minimum plant spread (E-W) at 30, 60, and 90 DAT (25.64, 46.85, and 70.85) was observed under control. The plant spread (E-W) at 30, 60, and 90 DAT under 200 ppm was noted 14.39, 15.85, and 12.87% more over control, respectively. "Boron is associated directly and indirectly with several plant functions, as it involves the growth of cells in newly emerging shoots and roots in some plants", [13].

#### 3.3 Plant Spread (N-S)

The maximum plant spread (N-S) at 30, 60, and 90 DAT (29.85, 58.15, and 85.50 cm) were observed under treatment  $N_2$ , which was statistically at par with the application of treatment  $N_3$ . The plant spread (N-S) at 30, 60, and 90 DAT under the treatment  $N_2$  was found 28.11, 28.45, and 17.42% higher than the control, respectively. "The response to potassium fertilization in terms of overall improvement in growth parameters is further supported by the fact that the leaching losses of potassium were more in light-textured soils. Most of these investigations have confirmed that manure compost and vermicompost usually have significant beneficial effects on plant growth",

[14]. Similarly, the application of different levels of boron also had significant increase in plant spread (N-S) at 30, 60, and 90 DAT (28.88, 55.86, and 84.46) in 200 ppm, and minimum plant spread (N-S) at 30, 60, and 90 DAT (24.86, 47.95, and 75.17) was noted under control. The plant spread (N-S) at 30, 60, and 90 DAT under 200 ppm was recorded at 16.17, 16.49, and 12.35% more over control, respectively. Foliar sprays of Boron increased the nitrogen content of the leaves. The leaf area was significantly ( $P < 0.05$ ) increased by nitrogen, possibly because nitrogen helps in greater assimilation of food material by the plant which resulted in greater meristematic activities of cells and accordingly the number of leaves, length, and width of the leaf. The findings are also in agreement with the findings of Solanki et al. [15] in brinjal.

### 3.4 Chlorophyll Content in Leaves

Maximum chlorophyll content in leaves at 45 DAT ( $1.315 \text{ mg.g}^{-1}$ ) was observed under the treatment  $N_3$ , which was statistically at par with the application of treatment  $N_4$ . Whereas, minimum chlorophyll content in leaves at 45DAT ( $1.014 \text{ mg.g}^{-1}$ ) was observed under control. The chlorophyll content in leaves under the treatment  $N_3$  was found 29.68% higher than the control. This might be because NPK with vermicompost helped in the expansion of leaf area and chlorophyll contents which together might have accelerated the photosynthetic rates and in turn increased the supply of carbohydrates to plants.

Potassium helps in chlorophyll formation ultimately the NPK is used for better vegetative growth. The effect of vermicompost on physicochemical properties imparts favorable soil structure for root growth which influenced better plant growth. The maximum chlorophyll contents in leaves at 45 DAT ( $1.270 \text{ mg/g}$ ) were recorded at 200 ppm and the minimum ( $1.130 \text{ mg.g}^{-1}$ ) in the control. The chlorophyll content in leaves under the treatment  $B_2$  (200 ppm) was recorded at 12.38% more than the control. These results point out that the spray of boron also helps in enhancing quality parameters such as the chlorophyll content of brinjal [16].

### 3.5 Fruit Length

The application of treatment  $N_3$  produced a significant ( $P < 0.05$ ) maximum fruit length (8.36 cm), which was statistically at par with the application of treatment  $N_4$ . While, the minimum fruit length (7.19 cm) was observed under control. The fruit length under the treatment  $N_3$  (50% NPK+50% VC) was registered 16.27% more as compared to the control and the application of boron produced the maximum fruit length (8.25 cm) in 200 ppm, which was minimum (7.41 cm) under control. Application of 200 ppm registered an increase of 11.33% higher fruit length over control. These results indicated that the spray of boron also helps in enhancing fruit size. Similar quality parameters were also reported by Selvi et al. [16].

**Table 1. Effect of NPK with vermicompost and boron on Plant height & Plant spread at different growth stages of brinjal**

Treatments	Plant height (cm)			Plant spread (cm)					
				(E-W)			(N-S)		
	30 DAT	60 DAT	At harvest	30 DAT	60 DAT	At harvest	30 DAT	60 DAT	At harvest
<b>INM levels</b>									
$N_0$ -Control	12.80	40.30	58.01	22.45	44.06	68.50	23.30	45.27	72.81
$N_1$ -100% NPK	16.85	46.56	65.52	28.11	51.31	74.79	27.19	52.15	79.31
$N_2$ -75% NPK+25% VC	19.15	51.31	72.56	30.65	56.22	81.29	29.85	58.15	85.50
$N_3$ -50% NPK+50% VC	17.88	48.61	68.99	29.36	52.89	79.77	27.88	54.69	84.34
$N_4$ -25% NPK+75% VC	17.57	47.29	65.78	28.70	51.67	75.22	27.45	52.57	79.46
$N_5$ -100% VC	15.43	44.79	63.50	25.82	48.56	74.49	25.73	49.96	78.56
<b>SEm±</b>	<b>0.44</b>	<b>1.15</b>	<b>1.90</b>	<b>0.72</b>	<b>1.47</b>	<b>1.77</b>	<b>0.73</b>	<b>1.56</b>	<b>2.00</b>
<b>CD (P=0.05)</b>	<b>1.28</b>	<b>3.32</b>	<b>5.47</b>	<b>2.06</b>	<b>4.21</b>	<b>5.09</b>	<b>2.09</b>	<b>4.47</b>	<b>5.73</b>
<b>Boron levels</b>									
$B_0$ -Control	15.66	43.25	61.13	25.64	46.85	70.85	24.86	47.95	75.17
$B_1$ -100 ppm	16.84	46.51	66.07	27.58	51.22	76.28	26.96	52.59	80.36
$B_2$ -200 ppm	17.78	49.68	69.98	29.33	54.28	79.90	28.88	55.86	84.46
<b>SEm±</b>	<b>0.31</b>	<b>0.82</b>	<b>1.35</b>	<b>0.51</b>	<b>1.04</b>	<b>1.25</b>	<b>0.51</b>	<b>1.10</b>	<b>1.41</b>
<b>CD (P=0.05)</b>	<b>0.90</b>	<b>2.34</b>	<b>3.87</b>	<b>1.45</b>	<b>2.98</b>	<b>3.60</b>	<b>1.47</b>	<b>3.16</b>	<b>4.05</b>

### 3.6 Fruit Weight

The maximum fruit weight (81.66 g) was recorded under treatment N<sub>3</sub>, which was statistically at par with the application of treatment N<sub>4</sub>. While the minimum fruit weight (70.53 g) was observed under control. The Fruit weight under the treatment N<sub>3</sub> was found 15.78% more as compared to the control. The application of boron gave a significant maximum fruit weight (80.38 g) in 200 ppm, and a minimum fruit weight (72.76 g) was observed under control. Application of 200 ppm registered an increase of 10.47% over control. The increased number of fruits due to the foliar spray of micronutrients might be attributed to enhanced photosynthetic activity, resulting in increased production and accumulation of carbohydrates and a favorable effect on vegetative growth and retention of flowers and fruits, which might have increased number and weight of fruits. Boron plays an important role in the uptake of calcium and the transport of carbohydrates. In case of B deficiency the rate of respiration increases and therefore more water loss. In boron-deficient plants, enzymatic activity increases which increases water loss from the fruit surface. Boron availability increases calcium synthesis which reduces respiration that might decrease the amount of moisture loss from the fruit surface Stanley et al. [17] also obtained similar results.

### 3.7 Fruit Yield

The maximum fruit yield ha<sup>-1</sup> (332.10 q.ha<sup>-1</sup>) was observed under treatment N<sub>3</sub>, which was statistically at par with the application of treatment N<sub>4</sub> (311.95 q.ha<sup>-1</sup>). Whereas, minimum fruit yield per hectare (212.49 q.ha<sup>-1</sup>) was observed under control. The percent increase in fruit yield per hectare under the treatment N<sub>3</sub> was found to be 56.28% more than the control. The application of treatment N<sub>3</sub> favored the metabolic and auxin activities in plants and ultimately resulted in increased fruit size, fruit weight, number of fruits per plant, yield per plant, and yield ha<sup>-1</sup>, [18]. The application of different boron levels also had a significant (P < 0.05) effect on fruit yield per hectare as compared to the control. The significant increase in fruit yield per hectare (328.52 q ha<sup>-1</sup>) was recorded in 200 ppm, and the minimum (236.11) was observed under control. The fruit yield per hectare under the treatment B<sub>2</sub> (200 ppm) was recorded at 39.13% more than the control. Boron deficiency commonly results in empty pollen grains, poor

pollen strength, and a reduced number of flowers per plant. The highest fruit set percentage might be due to optimum boron application, as boron plays important role in maintaining cell integrity, improving respiration, enhancing metabolic activities, and uptake of nutrients. Nonnecke [19] and Suganiya and Kumuthini [20] also obtained similar results.

### 3.8 Nitrogen Content in Fruit

The maximum nitrogen content in fruit (3.16%) was observed under the treatment N<sub>3</sub> (50% NPK+50% VC), followed by N<sub>4</sub> (25% NPK+75% VC) and followed by N<sub>2</sub> (75% NPK+25% VC). Whereas minimum nitrogen content in fruit (2.17%) was observed under control and the application of different levels of boron brought out a significant (P < 0.05) increase in nitrogen content in fruit as compared to control. Similar results were reported by Laxmi et al. [21]. The maximum nitrogen content in fruit (3.01%) was recorded at 200 ppm, which was 11.89% more than the control, and the minimum (2.69%) was observed under control. These results indicated that the boron spray, besides increasing the growth and yield also helps in enhancing the quality parameters of brinjal fruits. Similar parameters were also reported by Selvi et al. [16]. Adequate B is also required for effective nitrogen fixation and nodulation in legume crops. The results showed that the effect of different treatments on nutrient concentration and seed protein was significant (P < 0.05), [22].

### 3.9 Phosphorus Content in Fruit

The maximum phosphorus content in fruit (0.54%) was observed under the treatment N<sub>3</sub>, which was statistically at par with the application of treatment N<sub>4</sub>, whereas, the minimum phosphorus content in fruit (0.43%) was observed under control. The percent increase in phosphorus content in fruit under the treatment N<sub>3</sub> was found to be 25.58% over control. Among the various organic manures, the compost produced by earthworms (vermicompost), is a rich source of macro and micronutrients. It is evident from the data that phosphorus content in fruit with the application of boron 200 ppm was recorded as the maximum phosphorus content in fruit (0.52%) as compared to other treatments, while the minimum phosphorus content was recorded under control (0.45%). Boron deficiency adversely affects the quality and yield of many vegetables especially tomatoes [23,24].

**Table 2. Effect of NPK with vermicompost and boron on fruit length, fruit weight, fruit yield, chlorophyll content in leaves, nitrogen and phosphorous content in fruit of brinjal**

Treatments	Fruit length (cm)	Fruit weight (g)	Fruit yield (q ha <sup>-1</sup> )	Nitrogen content in fruit (%)	Phosphorous content in fruit (%)	Chlorophyll content in leaves (mg/g)
<b>INM levels</b>						
N <sub>0</sub> -Control	7.19	70.53	212.49	2.17	0.43	1.014
N <sub>1</sub> -100% NPK	7.78	76.12	284.11	2.92	0.48	1.204
N <sub>2</sub> -75%	7.85	76.30	286.13	2.95	0.49	1.214
NPK+25% VC						
N <sub>3</sub> -50%	8.36	81.66	332.10	3.16	0.54	1.315
NPK+50% VC						
N <sub>4</sub> -25%	8.21	79.39	311.95	3.06	0.52	1.275
NPK+75% VC						
N <sub>5</sub> -100% VC	7.68	75.97	281.00	2.89	0.46	1.198
<b>SEm±</b>	<b>0.17</b>	<b>1.41</b>	<b>11.07</b>	<b>0.05</b>	<b>0.01</b>	<b>0.025</b>
<b>CD (P=0.05)</b>	<b>0.48</b>	<b>4.05</b>	<b>31.81</b>	<b>0.14</b>	<b>0.03</b>	<b>0.073</b>
<b>Boron levels</b>						
B <sub>0</sub> -Control	7.41	72.76	236.11	2.69	0.45	1.130
B <sub>1</sub> -100 ppm	7.88	76.84	289.26	2.88	0.49	1.210
B <sub>2</sub> -200 ppm	8.25	80.38	328.52	3.01	0.52	1.270
<b>SEm±</b>	<b>0.12</b>	<b>1.00</b>	<b>7.83</b>	<b>0.04</b>	<b>0.01</b>	<b>0.018</b>
<b>CD (p=0.05)</b>	<b>0.34</b>	<b>2.86</b>	<b>22.49</b>	<b>0.10</b>	<b>0.02</b>	<b>0.051</b>

#### 4. CONCLUSION

The treatment 50% NPK+50% VC or treatment boron (200 ppm) was found significantly better in terms of growth, yield, and quality compared to other treatments. Thus, the application of 50% NPK+50% VC or Boron 200 ppm to brinjal is recommended. The results are only indicative and require further experimentation to arrive at a more consistent and final conclusion.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

- Chowdhury R, Tah J. Estimation of heritable components of variation and character selection in eggplant (*Solanum melongena* L.). Continental Journal of Biological Sciences. 2011;4:31–36.
- Butu M, Rodino S. Fruit and vegetable-based beverages—nutritional properties and health benefits. The Science of Beverages. 2019;13:303–338.
- Anonymous 2012. KAU-Agri. Infotech portal. Available: <http://www.celkau.in/Crop/Vegetables/Brinjal/brinjal.aspx> (accessed on 08.08.2014).
- Manimegala G, Gunasekaran G. Effect of vermicompost and NPK fertilizer on growth and yield components of eggplant (*Solanum melongena* L.). International Journal of Scientific and Technology Research. 2020;9(1):1388–1391.
- Sat P, Saimbhi MS. Effect of varying levels of nitrogen and phosphorus on earliness and yield of brinjal hybrids. Journal Reasearch Crops. 2003;4(2):217–222.
- Badiger MK, Subbareddy NP, Roselind M, Shivaraj B. Influence of phosphorus fertilizer on the yield and quality attributes of tomato. Journal of the Indian Society of Soil Science. 2006;43:123–129.
- Mishra VK, Kumar S, Pandey VK. Effect of organic manure and biofertilizers on growth yield and quality of brinjal (*Solanum melongena* L.). International Journal of Pure and Applied Bioscience. 2018;6(1):704–707.
- Arancon NQ, Edwards CA, Atiyeh RM, Metzger JD. Effects of vermicomposts produced from food waste on green house peppers. Bioresource Technology. 2004;93:139–144.
- Mamta AW, Khursheed Roa RJ. Effect of vermicompost on growth of brinjal plant (*Solanum melongena*) under field conditions. Journal of New Biological Reports. 2012;1(1):25–28.
- Jokanovi MB. Boron toxicity and deficiency in agricultural plants. International Journal

- of Molecular Science. 2020;21(4):1424.
11. Basavarajeshwari CP, Hosamani RM, Ajjappalavara PS, Naik BH, Smitha RP, Ukkund KC. Effect of foliar application of micronutrients on growth and yield components of tomato (*Lycopersicon esculentum*). Karnataka Journal of Agriculture Science. 2008;21(3):428–430.
  12. Oyinlola EY. Response of irrigated tomatoes (*Lycopersicon lycopersicum* Karst) to boron fertilizer: Growth and nutrient concentration. Nigerian Journal of Soil and Environmental Research. 2004;5:62–69.
  13. Miwa K, Takano J, Fujiwara T. Molecular mechanisms of boron transport in plants and its modification for plant growth improvement. Tanpakushitsu Kakusan Koso. 2008;53(9):1173–1179.
  14. Kashyap S, Kumar S, Maji S, Kumar D. Effect of organic manures and inorganic fertilizers on growth yield and quality of brinjal (*Solanum melongena* L.) cv. Pant Rituraj. International Journal Agriculture Science. 2014;10(1):305–308.
  15. Solanki MM, Solanki MS, Thakare G, Jogi PD, Sapkal RD. Effect of zinc and boron on growth of brinjal (*Solanum melongena* L.). International Journal of Plant Sciences. 2017;12(2):160–163.
  16. Selvi D, Thiageshwari S, Santhy P, Kannan BR. Fruit yield and nutrient uptake by brinjal due to integrated nutrient management in an Inceptisol. Journal of Maharashtra Agricultural University. 2004;29(2):220–223.
  17. Stanley DW, Bourne MC, Stone AP, Wismer WV. Low temperature blanching effects of chemistry firmness and structure of canned green beans and carrots. Food Science. 1995; (60):327–333.
  18. Kumar AB, Gowda NC. Effect of different organic manures and inorganic fertilizers on growth and yield of brinjal (*Solanum melongena* L.). Asian Journal Horticulture. 2010;5(2):444–449.
  19. Nonnecke IBL. Vegetable production. Avi Book Publishers. New York USA. 1989;200–229.
  20. Suganiya S, Kumuthini HD. Effect of boron on flower and fruit set and yield of raton brinjal crop. International Journal of Scientific Research and Innovative Technology. 2015;2(1):135–141.
  21. Laxmi PR, Saravanan S, Naik ML. Effect of organic manures and inorganic fertilizers on plant growth yield fruit quality and shelf life of tomato (*Solanum lycopersicon* L.) cv. PKM-1. International Journal of Agricultural Science and Research. 2015;5(2):7–12.
  22. Salih Hemn O. Effect of foliar fertilization of Fe B and Zn on nutrient concentration and seed protein of cowpea "*Vigna unguiculata*". IOSRJ. Agricultural Veterinary Science. 2013;6:42–46.
  23. Imtiaz M, Rashid P, Khan MY, Memon Aslam M. The role of micronutrients in crop production and human health. Pakistan Journal of Botany. 2010;42(4):2565–2578.
  24. Abusaleha Shanmulagavelu KG. Studies on the effect of organic and inorganic source of nitrogen on growth yield and quality of okra (*Abelmoschus esculentus*). Indian Journal of Horticulture. 1988;45(3-4):312–318.

© 2023 Yadav et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/97905>