



Effect of Different Mulching Material on Watermelon [*Citrullus lanatus* (Thunb.)] Hybrid under Prayagraj Agro-climatic Condition

Monojit Singha^{a++*}, Samir Ebson Topno^{a#}
and Anita Kerketta^{a#}

^a Department of Horticulture (Vegetable Science), Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj – 211007, Uttar Pradesh, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2023/v13i92567

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/104152>

Original Research Article

Received: 02/06/2023

Accepted: 04/08/2023

Published: 08/08/2023

ABSTRACT

The present field experiment was carried out during 2021-2022 at the Vegetable Research Farm, Department of Horticulture, Sam Higginbottom University of Agriculture Technology and Sciences, Prayagraj. The experiment was laid out in Randomized Block Design with three replications and nine treatments. The investigation was carried out with a view to study the effect of different mulching material on growth, yield and quality of watermelon. The results of the experiment revealed that plant growth was superior with silver plastic mulch with respect to days of germination, numbers of branches per vine and vine length. Plants without mulch (control) resulted poor growth. Watermelon plants mulched with silver plastic recorded maximum number of leaves;

⁺⁺ M.Sc. Scholar;

[#] Assistant Professor;

*Corresponding author: E-mail: monojtsingha115@gmail.com;

whereas, minimum was in control. The yield characters such as fruit length, fruit diameter, number of fruits per plant, average fruit weight and number of fruits per plot were recorded significantly highest in the mulch treatment with silver plastic mulch. Similarly, the highest fruit yield per plot (27.52kg) and fruit yield per hectare of watermelon (66.04 t/ha) was also observed with the same treatment. From the economic point of view, silver plastic mulch resulted in the highest net return and also highest cost ratio. The quality characters such as highest fruit pulp weight and lowest acidity were noted in mulching treatment silver plastic mulch. This treatment also recorded maximum vitamin C, TSS. It is concluded that application different mulches in which silver plastic mulch significantly superior in growth, yield and quality parameter. Similarly, with economic point of view, treatment silver plastic mulch resulted in the highest net return and cost: benefit ratio.

Keywords: Mulching; watermelon; silver plastic mulch.

1. INTRODUCTION

Watermelon is a warm, long-season crop. This fruit mostly cultivated for its fresh juice and sweet flesh. Commercial cultivation of watermelon takes place on larger scale in summer season in most of the Asian countries. Watermelon belongs to the family of "Cucurbitaceae" and genus of "Citrullus". Watermelons are native to the Kalahari Desert of southern Africa and the first recorded watermelon crop was found in Egypt. Farmers can make good profits from cultivation of watermelon, if proper cultivation methods and farm management practices are followed. Individual watermelon plants produce both male and female flowers. The fruit size varies from 2 to 15 kg, depending on variety. However, seedless varieties need pollinators. Watermelon can be successfully grown in pots, containers, indoors, greenhouses and poly houses [1-3].

Watermelon is believed to have originated from indigenous tropical Africa. Its growth is favoured by long period of warm, dry weather. A temperature of 25°C to 30°C is ideal for growth and 25°C is the best temperature for fruit setting of watermelon. Environment significantly influences the flavour and sweetness of watermelon.

The process of covering the open surface of the ground by a layer of some external material is called mulching & the material used for covering is called as 'Mulch.' Mulching is usually practiced when cultivating commercially important crops, fruit trees, vegetables, flowers, nursery saplings, etc [4-8]. Mulching in general is a beneficial practice for crop production. Mulch conserves soil moisture, retained heat as well as it suppresses weed growth [9]. The greatest benefit from plastic mulch is that the soil temperature in the planting bed is raised, promoting faster crop development and earlier

harvest. silver plastic mulch can give a harvest earlier by some 7-14 days, while black plastic may advance the harvest date by 21 days. Soil water loss is reduced under plastic mulch. As a result, a more uniform soil moisture is maintained and irrigation frequency can be reduced. The growth of plants on mulch can be twice that of plants in unmulched soil [10,11].

The edible product from a mulched crop is cleaner and less subject to rot, since no soil is splashed on to the plants or fruit [12,13]. To keep plants, clean, they should be grown in a raised bed that is firm and slightly convex, with the highest point down the center of the row, while the plastic should be stretched tight to encourage the run-off of water. Mulch film is nearly impervious to carbon dioxide, which is necessary for photosynthesis [14,15]. Research has shown that high levels of carbon dioxide may build up under the plastic. Because the film does not allow the gas to penetrate, it has to escape through the holes punched for the plants and a "chimney effect" is created, resulting in abundant CO₂ for the actively growing leaves. This study aimed to evaluate the effect of different mulches on growth, yield and quality of watermelon and to estimate the economics of various treatments under Prayagraj agro-climatic condition.

2. MATERIALS AND METHODS

The experiment was conducted at Vegetable Research Farm, Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology & Sciences, Prayagraj (UP) during Prayagraj during the *summer* season of 2022. Soils of the experimental site are classified as rich loam soil. The annual rainfall in the region is about 1042 mm. The experiment was laid out in randomized block design with three replications. There were total nine treatment of mulching materials viz. T₀

(control). T₁ (paddy straw), T₂ (paddy husk), T₃ (coconut husk), T₄ (dry leaves), T₅ (saw dust mulch), T₆ newspaper mulch) and T₇ (black plastic mulch), T₈ (silver plastic mulch).

2.1 Statistical Analysis

The Data recorded throughout the course of investigation was subjected to Statistical analysis by using analysis of variance (ANOVA) for randomized block design (RBD) by Fischer and Yates (1963). Whenever 'F' test was found significant for comparing the means of two treatments, a critical difference (C. D. at 5%) was worked out.

3. RESULTS AND DISCUSSION

Growth parameters: Data pertaining to growth parameters which are Days of germination, Number of leaves, Vine length (cm), Number of branches per vine.

3.1 Days of Germination

The data related to the days of germination are given in Table1. Results indicated that the effect of different type of mulching material on days of germination was found to be significant over control. Earliest days of germination (5.08) was recorded in treatment T₈ (Silver plastic mulch) followed by T₇ (Black plastic mulch) (5.17). While the treatment control recorded significantly late days of germination (6.58).

3.2 Number of Leaves

The data related to the number of leaves are given in Table1. The various mulching treatments have significant effect on no of leaves. At harvesting time, maximum no of leaves (33.33) was recorded in mulching treatment (T₈) (Silver plastic mulch), followed by (Black plastic mulch), However, minimum (27.67) no of leaves was recorded in (T₀) (control) which were significantly different.

3.3 Vine Length (cm)

The data related to the length of main axis of vine are given in Table1. The various mulching treatments have significant effect on length of main axis of vine. At harvesting time, maximum vine length (237.97 cm) was recorded in mulching treatment (T₈) (Silver plastic mulch), followed by (Black plastic mulch), However, minimum (166.05 cm) vine length was recorded in (T₀) (control) which were significantly different.

3.4 No of Branches per Vine

The results obtained on the effect of different mulching treatments on number of branches per vine have been presented in Table1. The results revealed that various treatments significantly influenced the number of branches per vine. The perusal of data revealed that all mulching treatments were found significantly superior in sense of a greater number of branches per vine as compared to control. The maximum number of branches per vine (14.25) was recorded in (T₈) silver plastic mulch treatment, followed by black plastic mulch (13.33). However, (T₀) control treatment recorded significantly the lowest number of branches per vine (9.08).

Flowering parameters: Data pertaining to flowering parameters which are Days of 1st male flower, Days of 1st female flower, Node at 1st male flower emergence, Node at 1st female flower emergence, No of male flowers, No of female flowers, Sex ratio.

3.5 Days to Emergence of 1st Male Flower

The data related to Days of 1st male flower is given in Table 2. The various mulching treatments have significant effect on Days of 1st male flower. Minimum days to emergence of first male flower were recorded in Silver plastic mulch (19.32) followed by black plastic mulch, while maximum days to, male of first male flower were recorded in No mulch (control) (23.85).

3.6 Days to Emergence of 1st Female Flower

The data related to Days of 1st female flower is given in Table 2. The various mulching treatments have significant effect on Days of 1st female flower. minimum days to emergence of first female flower were recorded in Silver plastic mulch (24.17) followed by Black plastic mulch (25.83), while maximum days to emergence of first female flower were recorded in No mulch (control) (30.08).

3.7 Node Number at which First Male Flower Emergence

The data related Node at first male flower emergence is given in Table 2. The various mulching treatments have significant effect on node at first male flower emergence. Earliest number of node at first male flower was recorded in No mulch (control) (8.48) followed by Newspaper mulch, while late number of node at

first male flower was recorded in silver plastic mulch (11.14).

3.8 Node Number at Which First Female Flower Emergence

The data related Node at first female flower emergence is given in Table 2. The various mulching treatments have significant effect on node at first female flower emergence. Earliest number of node at first female flower was recorded in No mulch (control) (12.04) followed by Newspaper mulch (13.12) while late number of node at first female flower was recorded in Silver plastic mulch (14.74).

3.9 No of Male Flowers

The data related to no of male flowers is given in Table 2. The various mulching treatments have significant effect on Days of 1st male flower. Maximum number of male flowers was recorded in Silver plastic mulch (35.85) followed by Black plastic mulch while minimum number of male flowers was recorded in No mulch (control) (16.44).

3.10 No of Female Flowers

The data related to no of female flowers is given in Table 2. The various mulching treatments have significant effect on Days of 1st female flower. Maximum number of female flowers was recorded in Silver plastic mulch (5.22) followed by Black plastic mulch while minimum number of female flowers was recorded in No mulch (control) (4.22).

3.11 Sex Ratio

The data related to Sex ratio is given in Table 2. The various mulching treatments have significant effect on sex ratio. Maximum number of sex ratio was recorded in Silver plastic mulch (6.86) followed by Black plastic mulch while minimum number of female flowers was recorded in No mulch (control) (3.89).

Yield parameters: Data pertaining to yield parameters which are Fruit length (cm), Fruit diameter (cm), Number of fruit per plant, Fruit weight (kg), Total fruit yield per plot (kg), Fruit yield per hectare (tones).

3.12 Fruit Length (cm)

The mean data on fruit length (cm) of watermelon affected by different treatments of

mulching are presented in Table 3. There was significant difference found in various treatments. The maximum fruit length (29.66 cm) was noted in the treatment T₈ (Silver plastic mulch), followed by (Black plastic mulch), while, the minimum fruit length (25.17 cm) was recorded in treatment T₀ (control).

3.13 Fruit Diameter (cm)

The mean data on fruit diameter (cm) of watermelon affected by different treatments of mulching are presented in Table 3. There was significant difference found in various treatments. The maximum fruit diameter (18.44 cm) was noted in the treatment T₈ (Silver plastic mulch), followed by (Black plastic mulch) while, the minimum fruit diameter (13.27 cm) was recorded in treatment T₀ No mulch (control).

3.14 Number of Fruits per Plant

The data presenting to number of fruits per plant as affected by various treatments of mulching are presented in Table 3. Results indicated that the effect of different type of mulching material on number of fruits per plant was found to be significant over control. Significantly highest number of fruits per vine (2.97) was recorded in treatment T₈ (Silver plastic mulch) followed by T₇ (Black plastic mulch) (2.80). While the treatment control recorded significantly lowest number of fruits per plant (1.69).

3.15 Fruit Weight (kg)

The data are presented in Table 3. revealed that various treatments of mulching exerted a significant effect on average fruit weight (kg) of watermelon. It is observed from the data that all the treatments of mulching were found significantly superior in recording higher average fruit weight (kg) over control. The maximum average fruit weight (2.41 kg) was recorded in treatment T₈ (Silver plastic mulch) followed by Black plastic mulch whereas the minimum average fruit weight was recorded in No mulch (control) (1.83 kg).

3.16 Total Fruit Yield per Plot (kg)

The mean data on fruit yield (kg/plot) of watermelon as affected by different type of mulching material are presented in Table 3. It is observed from the data that all the treatments of mulching were found significantly superior in recording higher fruit yield (kg/plot) over control.

Table 1. Effect of different mulches on yield of watermelon

Notation	Treatments	Days of germination	No of leaves (At harvest)	Vine length (cm) (At harvest)	Number of branches per vine
T ₀	No mulch (Control)	6.58	27.67	166.05	9.08
T ₁	Paddy straw	6.00	30.57	206.42	11.42
T ₂	Paddy husk mulch	6.17	28.55	196.26	10.08
T ₃	Coconut husk	5.92	30.45	224.79	12.17
T ₄	Dry leaves	6.00	29.71	189.28	9.50
T ₅	Saw dust mulch	6.25	29.54	223.12	13.33
T ₆	Newspaper mulch	5.33	28.72	184.19	9.33
T ₇	Black plastic mulch	5.17	32.91	230.71	13.33
T ₈	Silver plastic mulch	5.08	33.33	237.97	14.25
F-TEST		S	S	S	S
SE(d)±		0.22	0.16	0.46	0.42
CD 5%		0.46	0.34	0.98	0.89
C.V		4.55	0.65	0.27	4.49

Table 2. Effect of different mulches on flowering of watermelon

Notation	Treatments	Days to 1 st male flower	Days to 1 st female flower	Node at 1 st male flower emergence	Node at 1 st female flower emergence	No of Male flowers	No of Female flowers	Sex ratio
T ₀	No mulch (Control)	23.85	30.08	8.48	12.04	16.44	4.22	3.89
T ₁	Paddy straw	23.11	26.58	10.72	14.29	28.89	4.69	6.15
T ₂	Paddy husk mulch	22.82	28.25	9.79	13.36	26.56	5.13	5.18
T ₃	Coconut husk	22.07	27.25	10.55	14.12	32.19	5.45	5.90
T ₄	Dry leaves	21.65	29.58	9.55	13.37	19.45	4.08	4.76
T ₅	Saw dust mulch	20.04	25.67	10.22	13.79	33.53	5.89	5.69
T ₆	Newspaper mulch	21.22	28.58	9.55	13.12	29.55	5.85	5.04
T ₇	Black plastic mulch	20.11	25.83	10.97	14.54	33.24	5.19	6.40
T ₈	Silver plastic mulch	19.32	24.17	11.14	14.74	35.85	5.22	6.86
F-TEST		S	S	S	S	S	S	S
SE(d)±		0.18	0.38	0.38	0.09	0.31	0.09	0.13
CD 5%		0.37	0.80	0.80	0.19	0.66	0.20	0.28
C.V		1.00	1.69	4.57	0.81	1.33	2.27	2.87

The maximum yield per plot of watermelon (27.52 kg) was recorded in treatment T₈ (Silver plastic mulch) followed by Black plastic mulch Whereas, the minimum yield per plot was recorded in No mulch (control) (14.32 kg).

3.17 Fruit Yield per Hectare (tonnes)

The fruit yield (t/ha) of watermelon as affected by various treatments of mulching are presented in Table 3. Results indicated that the effect of different types of mulching material on fruit yield (t/ha) was found to be significant over Significantly the highest fruit yield per hectare of watermelon (66.04 t/ha) was recorded in treatment T₈(Silver plastic mulch) followed by Black plastic mulch while significantly lowest fruit yield per hectare (36.35 t/ha) was noted in the treatment T₀ No mulch (control).

Quality parameters: Data pertaining to Quality parameters which are Vitamin C (mg/100g), Acidity (%), TSS (Brix), Number of seed/fruits, Pulp weight (g).

3.18 Vitamin C (mg/100 g)

The analysis of variance presented in Table 4. showed significant differences among treatments and their interactions for Vitamin C (mg/100g) ranged from 7.57 to 8.12 The mean values for Vitamin C (mg/100g) revealed that the maximum Vitamin C was recorded in the Silver plastic mulch with (8.12 mg/100g)) and Followed by the Black plastic mulch with (8.07 mg/100g)) and Minimum was recorded in the No mulch (control) with (7.57 mg/100g)).

3.19 Acidity (%)

The data showed that the effect of various treatments of mulching on acidity (%) was found significant fruits of watermelon presented in Table 4. In The perusal of the data reveals that the lowest per cent of acidity was found in all the treatment of mulching as compared to control. Minimum acidity of watermelon fruit (0.31%) was observed in silver plastic mulch (T₈) followed by Black plastic mulch, while the maximum acidity (0.43%) was recorded in No mulch (control) (T₀).

3.20 TSS (°Brix)

The mean data on TSS as affected by different type of mulching material are presented in Table

4. Maximum TSS was recorded in the Silver plastic mulch with (9.61)) and Followed by the Black plastic mulch with (9.51) and Minimum was recorded in the No mulch (control) with (8.79).

3.21 Number of Seed/Fruits

The mean data on number of seeds per fruit in watermelon as affected by different type of mulching material are presented in Table 4. Minimum number of seed/fruits of watermelon (386.15) was observed in No mulch (control) followed by Newspaper mulch, while the maximum number of seed/fruits (485.83) was recorded in Silver plastic mulch (control) (T₈).

3.22 Pulp Weight (g)

The data are presented in Table 4 revealed that various treatments of mulching exerted a significant effect on average pulp weight (g) of watermelon. It is observed from the data that all the treatments of mulching were found significantly superior in recording higher average pulp weight (kg) over control. The maximum average pulp weight (1225.79 g) was recorded in treatment T₈ (Silver plastic mulch) followed by Black plastic mulch whereas the minimum average pulp weight was recorded in No mulch (control) (925.56 g).

3.23 Discussion

Effect of different colour of mulch on growth and yield attributing characters the results showed that different types of mulching materials significantly influenced the growth parameters of watermelon viz., number of lateral branches per vine and main vine length over control. The increase in growth parameters was attributed to sufficient soil moisture near root zone resulted from minimization of evaporation loss as well as reduced weed growth due to mulching. The extended retention of moisture and availability of moisture also lead to higher uptake of nutrient for proper growth and development of plants, resulted higher growth of plant, as compared to no mulch condition. The changes in soil temperature below plastic mulch could be attributed to different manners of heating and heat transfer to soil and also to heat accumulation during day and loss during night. The results indicated that the effect of different mulching material on fruit length and fruit diameter of watermelon is significant over

Table 3. Effect of different mulches on yield of watermelon

Notation	Treatments	Fruit length (cm)	Fruit Diameter (cm)	Number of fruits per plant	Fruit weight (kg)	Total fruit yield per plot (kg)	Fruit yield per hectare (tonnes)
T ₀	No mulch (Control)	25.17	13.27	1.69	1.83	14.32	36.35
T ₁	Paddy straw	26.53	15.61	2.22	2.01	18.18	43.62
T ₂	Paddy husk mulch	26.03	14.27	2.47	1.91	19.06	45.73
T ₃	Coconut husk	27.42	16.36	2.64	2.11	21.70	52.07
T ₄	Dry leaves	25.20	13.69	1.80	1.89	16.16	38.77
T ₅	Saw dust mulch	28.85	17.52	2.64	2.26	22.19	53.25
T ₆	Newspaper mulch	27.14	13.52	2.55	1.94	20.15	48.36
T ₇	Black plastic mulch	29.02	17.44	2.80	2.30	26.69	64.06
T ₈	Silver plastic mulch	29.66	18.44	2.97	2.41	27.52	66.04
F-TEST		S	S	S	S	S	S
SE(d)±		0.56	0.09	0.05	0.04	0.24	0.58
CD 5%		1.19	0.20	0.10	0.09	0.52	1.23
C.V		2.52	0.84	2.42	2.41	1.44	1.42

Table 4. Effect of different mulches on Quality of watermelon

Treatments	Treatments	Vitamin C (mg/100 g)	Acidity (%)	TSS (°Brix)	Number of seed/fruits	Pulp weight (g)
T ₀	No mulch (Control)	7.57	0.43	8.79	386.15	925.56
T ₁	Paddy straw	7.78	0.38	9.37	437.00	986.37
T ₂	Paddy husk mulch	7.61	0.41	9.33	403.64	981.59
T ₃	Coconut husk	7.83	0.36	9.40	410.79	1065.72
T ₄	Dry leaves	7.64	0.42	9.32	399.90	976.40
T ₅	Saw dust mulch	8.04	0.33	9.52	450.42	1155.97
T ₆	Newspaper mulch	7.69	0.41	9.31	407.03	1008.60
T ₇	Black plastic mulch	8.07	0.33	9.51	480.68	1182.23
T ₈	Silver plastic mulch	8.12	0.31	9.61	485.83	1225.79
F-TEST		S	S	S	S	S
SE(d)±		0.04	0.00	0.05	3.37	29.70
CD 5%		0.08	0.01	0.11	7.15	62.96
C.V		0.57	1.23	0.65	0.96	3.44

Table 5. Effect of different mulching material on economics of watermelon

Notation	Treatments	Gross return (Rs. /ha)	Cost of cultivation (Rs. /ha)	Net return (Rs. /ha)	B:C Ratio
T ₀	No mulch (Control)	363500	1,21,508	2,41,992	2.99
T ₁	Paddy straw	436200	1,23,008	3,13,192	3.55
T ₂	Paddy husk mulch	457300	1,26,058	3,31,242	3.63
T ₃	Coconut husk	520700	1,33,508	3,87,192	3.90
T ₄	Dry leaves	387700	1,21,858	2,65,842	3.18
T ₅	Saw dust mulch	532500	1,26,308	4,06,192	4.22
T ₆	Newspaper mulch	486300	1,27,508	3,56,092	3.79
T ₇	Black plastic mulch	640600	1,49,508	4,91,092	4.28
T ₈	Silver plastic mulch	660400	1,46,508	5,13,892	4.51

control. Maximum fruit length and diameter were found in silver plastic mulch, whereas the minimum fruit length and width of watermelon were noted in under no mulch condition. It was found that all treatments of mulching material were significantly increased the average fruit weight (kg) and fruit yield of watermelon than no mulch condition. Plants under polyethylene mulch (silver plastic mulch) produced larger fruit and gave higher fruit yield because of better plant growth due to favourable hydro-thermal regime of soil and complete weed free environment. Similar findings have also been obtained earlier [16].

3.24 Cost Benefit Analysis

Considering all the economics of different treatment It can be seen from the Table 5 that highest net returns were obtained in different mulches as compared to control or no mulch condition. Watermelon under silver plastic mulch recorded higher net monetary returns (Rs 513892) and lowest net monetary return under no mulch condition (Rs 241992). The maximum cost benefit ratio was obtained with treatment silver mulch (4.51) as compared to no mulch (2.99).

4. CONCLUSION

From the present investigation, it is concluded that, Silver plastic mulch (T8) performed best in terms of growth viz., vine length (237.97cm), yield (66.04 t/ha) and quality viz, Pulp weight (g) (1225.79).

The maximum benefit cost ratio was also observed in Silver plastic mulch (T8) with 4.51.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Ghorai AK, Bera PS. Note on effect of straw mulch and different levels of irrigation on yield and yield components of pointed gourd (*Trichosanthes dioica* L.). Indian Agriculturist. 1999;43(1/2):89-92.
2. Hallidri M. Comparison of the different mulching materials on the growth, yield and quality of cucumber (*Cucumis sativus* L.). Acta Hort. 2001;559:49-54.

3. Ibarra L. Flores J, Diaz Perez JC. Growth and response to plastic mulch and row covers. *Scientia Horticulturae*, 2001; 87(1/2):139-145.
4. Ansary SH, Roy DC. Effect of irrigation and mulching on growth, yield and quality of watermelon (*Citrullus lanatus* Thunb.). Environment and Ecology. 2005;23(Spl-1):141-143.
5. Alenazi M, Abdel-Razzak H, Ibrahim A, Wahb-Allah M, Alsadon A. Response of muskmelon cultivars to plastic mulch and irrigation regimes under greenhouse conditions. J. Anim. Plant. Sci. 2015;25(5): 1398-1410.
6. Battikhi AM, Ghawi I. Muskmelon production under mulch and trickle irrigation in the Jordan valley. Hort. Sci. 1987;22(4):578-581.
7. DOA (Department of Agriculture). Myanmar Horticultural Crops Production Report. Ministry of Agriculture and Irrigation; 2013.
8. Ghosh SN, Tarai RK. Effect of mulching on soil moisture, yield and quality of ber (*Ziziphus mauritiana* L.). Indian J. Soil Cons. 2007;35(3):246-248.
9. Ahmad, A. M., Iqbal, M. and Minhas, N.M. (2007) Effect of planting geometry and mulching on moisture conservation, weed control, and wheat growth under rainfed condition. Pakistan J. Bot., 39(4):1189-1195.
10. Maged AEN. Effect of mulch types on soil environmental conditions and their effect on the growth and yield of cucumber plants. J. Appl. Sci. Res. 2006;2(2): 67-73.
11. Sharma, H. G. and Agrawal, Narendra. Effect of different colour mulches on the growth and yield of tomato under drip irrigation. Plant Archives. 2004;4(1): 93-99.
12. Osiru T, Hahn J. Effect of mulching on the growth, yield and quality of yams. J. Plant Physiol. 1994;64(8):201:205.
13. Parmar HN, Polara ND, Viradiya RR. Effect of mulching material on growth, yield and quality of watermelon (*Citrullus lanatus* thunb) cv. Kiran. Univers. J. Agric. Res. 2013;1(2):30-37.
14. Lament WJ Jr. Plastic mulches for the production of vegetable crops. Hort. Sci. 1993;3(1):35-39.
15. Maughan T, Drost D. Use of Plastic Mulch for Vegetable Production. Utah State University. USA; 2016.

16. Dean B, Zanic K, Dumicic G, Culjak TG, Ban SG. The type of polythene mulch impacts vegetative growth, yield and aphid populations in watermelon production. J. Food, Agri. and Envi. 2004;7(3-4): 543-550.

© 2023 Singha et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/104152>