



Evaluation of the Phytochemical Composition of Some Commonly Sold Male Herbal Fertility Supplements in Port Harcourt, Rivers State, Nigeria

Arit Okechukwu Nwogu ^{a*},
Davies G. Tamuno-Emine ^a, Adline E. Ben-Chioma ^a
and Ebirien-Agana Samuel Bartimaeus ^a

^a Department of Medical Laboratory Science, Rivers State University, Port Harcourt, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJMAH/2023/v21i10891

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/103176>

Original Research Article

Received: 22/05/2023
Accepted: 26/07/2023
Published: 18/08/2023

ABSTRACT

Plants are major sources used in the production of pharmaceutically active drugs and they contain diverse phytochemicals which give these herbal supplements their biological and pharmacological effects. This study evaluated the phytochemical constituents of some commonly sold male herbal fertility supplements (Libron Herbal, Mascum Herbal Pride and Energy 3000) in Port Harcourt, Nigeria. The qualitative and quantitative analysis of the herbal supplements was done using standard procedures. The GC 7890B 5977A and Mass Spectrometry Device (MSD) Model was used for the GC/MS Study. Results of the qualitative phytochemical analysis of the supplements shows that flavonoid, protodiocin, tannins, phenols and alkaloids were present in libron herbal supplement. Protodiocin, saponins, tannins, phenols, alkaloids and cardiacglycosides were present

*Corresponding author: E-mail: arietta457@gmail.com;

in mascum herbal supplement while flavonoids, protodiocin, tannins, alkaloids, anthroquinones and terpenoids were present in Energy 3000 herbal supplement. For the quantitative phytochemical analysis of the herbal supplements, libron herbal supplement contained Flavonoid $0.24 \pm 0.00\%$, protodiocin $0.29 \pm 0.01\%$, tannins $0.37 \pm 0.02\%$, phenols $0.18 \pm 0.00\%$ and alkaloid $0.28 \pm 0.10\%$. Mascum herbal supplement were seen to contain protodiocin $0.27 \pm 0.07\%$, saponins $0.48 \pm 0.08\%$, tannis $0.38 \pm 0.06\%$, phenols $0.21 \pm 0.07\%$, alkaloids $0.18 \pm 0.10\%$, and cardiaglycosides $0.27 \pm 0.07\%$ while Energy 3000 was made up of flavonoid $0.10 \pm 0.00\%$, protodiocin $0.29 \pm 0.04\%$, tannis $0.64 \pm 0.10\%$, phenols $0.28 \pm 0.01\%$, alkaloids $0.26 \pm 0.01\%$, anthroquinones $0.27 \pm 0.10\%$ and terpenoids $0.25 \pm 0.00\%$. The GC/MS study was seen to contain various chemical compounds. The three herbal supplements assayed contain diverse phytochemicals that may enhance male fertility as well as some chemical compounds that could alter hormonal functioning of some biochemical systems in the albino rats.

Keywords: *Plants; male fertility; herbal supplements; phytochemical constituents; libron herbal; mascum herbal pride; energy 3000.*

1. INTRODUCTION

Plants are major sources used in the production of pharmaceutically active drugs by pharmaceutical industries and a very important branch of medicine; as such this active components are being employed in the production of herbal supplements with a general belief that the raw form which is used in the production of herbal supplements are more potent. They have been proven to possess therapeutic potentials, and are widely used by traditionalists in the treatment of various diseases since antiquity [1].

Plants contain diverse phytochemicals and have been in use by humans from ancient time in many parts of the world including Nigeria for the treatment and prevention of various illnesses [2]. Even in the emergence of complementary medicine, herbs are considered one of the most important branches of traditional medicine and as such they play important roles in health care system especially in rural areas and developing countries due to the level of poverty in this part of the world. About 80% of the world's population depends on herbal medicines as their primary source of health care [3]. Herbs and other natural products represent 50% of all currently utilized medication worldwide [4].

The use of plant products in pharmaceutical industries has given herbal product a place beside complementary medicine [5]. Herbal substance is considered medicinal when it possesses the ability to treat or prevent disease [6]. Due to the global usage of herbal supplement, the market is infiltrated with many new products as such public health issues and concerns about its safety demands the knowledge of its components as many of them

remain untested and unmonitored in the open market. Hence the need to evaluate its phytochemical components. Phytochemicals in plant give them their biological and pharmacological effect in the treatment of various illnesses and also play many roles contributing to the plant colour, texture, aroma, and flavour and also protect the plant cells from environmental hazards such as drought, ultraviolet exposure, pathogens and environmental pollution and also protects humans that eat it in significant amount [7]. Commonly used herbal supplements in this study were libron herbal supplement composed of *Allum sativum* (Garlic), *Panax ginseng* (Red Ginseng) *Nymphaea alba* (waterlily), *Rosa gallica* (French rose) *Monarda didyma* (Bee Balm) and *Aframomum melegueta*. Mascum herbal pride composed of *Hepo createa pollens*, (Pollen) *Xylopiya aethiopicia*, (Guinea pepper) *Medicago sativa* (Alfafa or Lucerne) *Tetrapleura tereaptera*, (Aidan fruit) *Urtica dioica*. (Stinging nettle) and Energy 3000 supplement made up of *Citrus aurantifolia* (Lime) (0.5%), *Psidium guajava* (Guava) (0.5%), *Xylopiya aethiopicia* (Guinea pepper) (1%), *Sesamum indicum* (Sesame) (1%), *Magnifera indica* (Mango) (2%) and water. The study was aimed at evaluating the phytochemical composition of some commonly sold male herbal fertility supplements manely Libron herbal, Mascum herbal pride, and Energy 3000 in Port Harcourt, Nigeria.

2. MATERIALS AND METHODS

The three herbal supplements were obtained from local herbal stores in mile III and mile IV of Rumueme, Port Harcourt, Rivers State, Nigeria. Phytochemical screening of all the supplements were carried out using standard method by Trease & Evans, 1989. A Spectro-UV-Visible 2500 was used for the quantitative

phytochemical analysis while a GC 789OB 5977A and Mass Spectrometry Device (MSD) Model was used for the GC/MS Study.

2.1 Quantitative Phytochemical Analysis of the Herbal Supplements (Libron, Mascum and Energy 3000)

The following herbal products (Libron capsule and Mascum herbal pride) were properly washed with deionized water to remove dirt. They were all dried under room temperature and pulverized using warring blender. The extraction was done using Soxhlet extractor in a continuous extraction Process for 72 hours using Methanol as the extracting solvent. A weighed portion of the pulverized samples (200 g) was used against 1000 ml of absolute methanol (BDH Chemicals) for 72 hours. The extracts including Energy 3000 were concentrated using rotary evaporator set at 60 °C. The extract was then prepared using distilled water for quantitative phytochemical analysis. The extract was read using spectrophotometric methods by Spectro-UV-Visible 2500, manufactured by Labo Med Inc., USA via scanning method at different wave lengths in accordance with ISO 17025.

The absorbance obtained was compared with phytochemical reference standard and the results indicated.

2.2 Gas Chromatography and Mass Spectrometry (GC/MS)

The GC/MS study of the herbal drugs was done using a GC 789OB 5977A and Mass Spectrometry Device (MSD) Model. Gas chromatography-mass spectrometry (GC-MS) is a quick and accurate approach for examining the constituents in complicated mixtures. It enables the identification of several chemicals in modest amounts of plant materials. It provides the structure and weight of phytocompounds present in a plant sample [8].

3. RESULTS

3.1 Qualitative Phytochemical Test Results of Libron Herbal

The result of the qualitative phytochemical analysis of libron herbal shows that it contains various chemical constituents such as flavonoid, protodiocin, tannis, phenols and alkaloids. They were present in large amount as shown in Table 1.

Table 1. Qualitative screening result for phytochemical components in Libron Herbal

Phytochemicals	Result
Flavonoid	+
Protodiocin	++
Saponins	-
Tannins	++
Phenols	+
Alkaloids	+
Terpenes	-
Cardiaglycosides	-
Quinones	-
Steroids	-
Terpenoids	-

+ = present
- = absent

3.2 Qualitative Phytochemical Test Results of Mascum Herbal Pride

The result of the qualitative phytochemical analysis of mascum herbal pride is shown to contain various chemical constituents, including Protodiocin, saponins, phenols, alkaloids and cardiaglycosides in large quantity as shown in Table 2.

Table 2. Qualitative screening result for phytochemical components in Mascum Herbal pride

Phytochemicals	Results
Flavonoid	-
Protodiocin	+
Saponin	++
Tannins	++
Phenols	+
Alkaloids	+
Terpenes	-
Cardiaglycosides	+
Quinones	-
Steroids	-
Terpenoids	-

+ = present
- = absent

3.3 Qualitative Phytochemical Test Results of Energy 3000

The result of the qualitative phytochemical analysis of Energy 3000 is shown to contain various chemical constituents such as flavonoids, protodiocin, tannis, phenols, alkaloids, anthroquinones and terpenoids. These were present in large quantity as shown in Table 3.

Table 3. Qualitative screening result for phytochemical components in energy 3000

Phytochemicals	Results
Flavonoid	+
Protodiocin	++
Saponins	-
Tannins	++
Phenols	+
Alkaloids	+
Terpenes	-
Cardiaglycosides	-
Anthroquinones	+
Steroids	-
Terpenoids	+

+ = present
- = absent

3.4 Quantitative Phytochemical Test Results of Libron Herbal

The results of the quantitative phytochemical analysis of Libron herbal shows that it contains flavonoid $0.24 \pm 0.00\%$, protodiocin $0.29 \pm 0.01\%$, tannins $0.37 \pm 0.02\%$, phenols $0.18 \pm 0.00\%$ and alkaloid $0.28 \pm 0.10\%$ as seen on Table 4.

3.5 Quantitative Results of the Phytochemical Analysis of Mascum Herbal Pride

The result of the quantitative phytochemical analysis of Mascum herbal pride shows that it contains protodiocin $0.27 \pm 0.07\%$, saponins $0.48 \pm 0.08\%$, tannins $0.38 \pm 0.06\%$, phenols $0.21 \pm 0.07\%$, alkaloids $0.18 \pm 0.10\%$, and cardiaglycosides $0.27 \pm 0.07\%$ as shown on Table 5.

Table 4. Quantitative screening result for phytochemical constituents in Libron Herbal

Phytochemicals	Percentage (%) constituents in Libron Herbal
Flavonoid	0.24 ± 0.00
Protodiocin	0.29 ± 0.01
Tannins	0.37 ± 0.02
Phenols	0.18 ± 0.00
Alkaloid	0.28 ± 0.1

3.6 Quantitative Results of the Phytochemical Analysis of Energy 3000

The result of the quantitative phytochemical analysis of Energy 3000 shows that it contains

flavonoid $0.10 \pm 0.00\%$, protodiocin $0.29 \pm 0.04\%$, tannins $0.64 \pm 0.10\%$, phenols $0.28 \pm 0.01\%$, alkaloids $0.26 \pm 0.01\%$, anthroquinones $0.27 \pm 0.10\%$ and terpenoids $0.25 \pm 0.00\%$ as shown on Table 6.

Table 5. Quantitative screening result for phytochemical constituents in Mascum Herbal

Phytochemical	Percentage (%) constituents in Mascum Herbal
Protodiocin	0.23 ± 0.10
Saponins	0.48 ± 0.08
Tannins	0.38 ± 0.06
Phenols	0.21 ± 0.07
Alkaloids	0.18 ± 0.10
Cardiaglycosides	0.27 ± 0.07

3.7 Qualitative Results of Gas Chromatographic and Mass Spectrometry of the Three Supplements

3.7.1 Qualitative results of gas chromatographic and mass spectrometry of Libron Herbal

The result of the qualitative Gas chromatographic and mass spectrometry result of Libron capsule shows that it contains various chemical compounds. Cyclohexanol, 1, 5-cyclooctadiene, carvone, 2-cyclohexene-1-ol, beta-famesene, beta capaene, bicyclogermacrene, cyclohexane, deodecanoic acid, carotol, Apiol, methyl tetradecanoate, hexadecanoic acid, 9, 12-octadecadienoic acid, methyl stearate, 4,4,8-trimethyl-non-5-enal and 9,octadecenoic acid.

3.7.2 Qualitative results of gas chromatographic and mass spectrometry of Mascum Herbal pride

The result of the qualitative gas chromatographic and mass spectrometry analysis of Mascum herbal pride shows that it contains a large number of chemical constituents. Salicyclic acid, methyl- 8-methyl decanoate, nonanoic acid, benzoic acid, carvone, benzene, cyclohexasiloxane, decanoic acid, cyclo heptasiloxane, dodecanoic acid, 2,5-dihydroxybenzoic acid, methyl tetra decanoate, hexadecanoic acid, 9-octadecanoic acid, methyl stearate, mercaptoacetic acid, diisocetyl phthate, cis-vaccenic acid and oleic acid is seen to be present in large quantity.

Table 6. Quantitative screening result for phytochemical constituents in energy 3000

Phytochemical	Percentage (%) constituents in energy 3000
Flavonoid	0.10 ± 0.00
Protodiocin	0.29 ± 0.04
Tannins	0.64 ± 0.10
Phenols	0.28 ± 0.01
Alkaloids	0.26 ± 0.01
Anthroquinones	0.27 ± 0.10
Terpenoids	0.25 ± 0.00

3.7.3 Qualitative results of gas chromatographic and mass spectrometry of energy 3000

The result of the qualitative gas chromatography and mass spectrometry analysis of Energy 3000 shows that it contains numerous chemical compounds. Endo-Borneol, cis-dihydrocarvone, cyclohexanol, benzoic acid, carvone, cyclohexasiloxane, decanoic acid, trans-carveyl acetate, benzene, cyclopenta cis-beta-farnesene, cis-muurolo-4-5-diene, cycloheptasiloxane, gamma-muuroloene, Napthalene, bicycle-5-2 non-1-ene, carotol, epicubenol, Apiol, dihydroxybenzoic acid, alpha-cadinol, methyl tetradecanoate, hexadecanoic acid, 9, 12 octadecadienoic acid, methyl stearate, 1,3-octadecenal, oleic acid and 1-5-9-undecatriene.

4. DISCUSSION

The qualitative and quantitative phytochemical components present in Libron herbal tablets, Mascum herbal pride and Energy 3000 herbal supplements were demonstrated in this study. Several plants have been proven to contain various phytochemicals with these compounds playing essential roles. Miyoung et al. [9]; and Kahkonen et al. [10] discovered that phytochemicals in plants serve as antioxidants, antibacterial, antiviral agents as well as possessing anti-inflammatory activities because they act as free radical scavengers as they are potential reducing agents protecting organs from oxidative damage due to the presence of hydroxyl groups. In this study, it was observed that these herbal supplements contain numerous phytochemicals such as flavonoids, anthocyanins and phenolic acid as such they are able to serve as good antioxidants possessing anti-inflammatory activities. Somayeh et al. [11] revealed that *Rosa gallica* (french rose) a component of Libron herbal supplement possesses flavonoids, anthocyanins and

phenolic acid thereby exerting its anti-inflammatory effects on the seminiferous tubules. This is similar to studies by Muhammad et al. [12] in which *Allum sativa* (garlic) another component of libron was seen to contain flavonoids, saponins, tannins, phenols, alkaloids and terpenoids. Though another study by Ebomoyi & Ahumibe, [13] gives contradicting results.

Protodiocin are bioactive components and have therapeutic role as such increases the concentration of androgen receptors in cells, thereby causing the organism to become more sensitive to androgens like testosterone as seen in a study carried out by Lakshmi et al. [14]. This study reveals the presence of protodiocin in all the three herbal supplement hence an improvement in testosterone levels is expected upon its usage.

From this study, Mascum herbal pride demonstrates qualitative and quantitative phytochemical components. Shoushtari et al. [15] in his study discovered that Hepo-cretea pollens (Pollen) one of the component of mascum herbal pride contains flavonoid, phenols and saponins and exert anti-inflammatory and anti-oxidative effect. Studies by Abedi et al. [16] also reveals that extract of pollen significantly increased sperm motility, LH, FSH and testosterone levels. *Xylopi aethiopica* (Guinea pepper) another component of mascum herbal pride has been reported to contain flavonoids, tannins, phlobotannins, phenol, anthroquinones and saponins and reduces sperm count, motility, testosterone, FSH and LH levels in a study by Yusuf et al. [17].

Ali et al. [18] in his study reveals that *Medicago sativa* (alfafa or lucerne), another component of mascum herbal pride supplement contains saponins, lignin, tannins, alkaloids, triterpenes, glycosides, flavons and sterols as such serves as anti-inflammatory and antioxidant agents. Also *Tetrapleura tetraptera* (aidan fruit), another component in mascum herbal pride was reported to contain tannins, flavonoid, saponin, triterpenoid and coumarin thereby serving as anti-inflammatory, anti-coagulant and anti-oxidant agents in a study by Adusei et al. [19]. Adalakun et al. [20] in his study reveals that this plant improves spermatogenic activity and maintains testicular integrity. Phytochemicals such as saponins, flavonoid, tannins, phenols, alkaloids, terpenes and protodiocin were identified in mascum herbal pride so the observed phytochemicals may be responsible for

the increase testicular function attributable to this supplement as seen in a study by Okam et al. [21] upon sub-chronic administration of Mascum herbal pride on male albino rats for 90 days. He observed an increase in sperm parameters and recommended a phytochemical screening in further studies. This study has been able to look at the phytochemical analysis of Mascum herbal pride.

For Energy 3000, its components, *Citrus aurantifolia* (Lime), contains flavonoids, flavones, triterpenoid which serves as anti-inflammatory, antioxidants and antibacterial thus protecting against bone marrow toxicity caused by lead acetate and also protects the liver and kidney tissue in a study by Aprioku & Obianime [22]. Idris et al. [23] has a contrary result in which *Citrus aurantifolia* (Lime), causes testicular damage and azoospermia.

Psidium guajava (guava) another component of Energy 3000 contains large number of antioxidants that gives it its antispasmodic, anti-inflammatory, antimicrobial effect in a study by Oluwole et al. [24]. Also Neeta et al. [25] reveals that *Sesamum indicum* (sesame) another component of Energy 3000 contain flavonoids, phenolic acid, alkaloids, tannins, saponins, steroids and terpenoids this gives it its anti-inflammatory and antioxidant effects. Also, Jing et al. [26] reveals that *Magnifera indica* (mango), another component of Energy 3000, contains tannins, polyphenols, flavonoids and triterpenoids as such it serves as a powerful antioxidant. Energy 3000, therefore, was thus reported to should possess anti-inflammatory and antioxidant effect since it possesses these various phytochemicals.

Apart from the phytochemical studies carried out, GC/MS studies is fundamental in understanding the response that could be seen in different organs as the individual herbal supplement contains some phytochemicals that may be protective or harmful to organs. For libron herbal supplement, some of its components such as cyclohexanol-1-ol, cyclohexane and Apoil were harmful to the reproductive organ as seen in a study by Dixit et al. [27] in which he administered 25 mg/kg of cyclohexanol-1-ol to rabbits for 40 days. It was seen to inhibit the process of spermatogenesis at the spermatocyte and spermatid levels, it reduced sialic acid concentration but there was no effect on the liver. Another study by Tyagi et al. [28] reveals severe reduction of the testes and seminiferous tubules

upon subcutaneous injection of cyclohexanol. Seminiferous tubules and Leydig cell nuclei were seen to be shrunken, sialic acid concentration dropped as such testosterone was reduced but no effect on the thyroid, kidney and liver cells was reported.

Mascum herbal pride was also observed to contain benzene, cyclohexasiloxane, cycloheptasiloxane, diisocetyl and thioglycolic acid following the GC/MS analysis. Studies by Micheal et al. [29] shows that siloxane compound is harmful to the liver whereas benzene was discovered to be harmful to the lungs and could cause malignant lymphoma and multiple myeloma in studies by Aksoy, [30]. Similar reports were seen by Fayed et al. [31] as he discovered significant increase in liver enzymes. Diisocetyl phthalate seen in mascum herbal pride acts as endocrine disruptors and alters liver functions in a study by Manisha & Ravinder [32].

5. CONCLUSION

All the three herbal supplements assayed contain diverse phytochemicals and chemical component with potentials to enhance male fertility. However, they also contain chemical component which activities could alter the normal functioning of the male reproductive system and liver functions after prolonged exposure.

6. RECOMMENDATION

Further studies on the effect of these chemicals on various organs should be carried out.

ACKNOWLEDGEMENT

Department of Pharmacognosy and Drug development, faculty of pharmaceutical sciences, Ahmadu Bello University, Zaria, Kaduna State for GC/MS studies.

Department of Biochemistry, faculty of science, Rivers State University for phytochemical analysis.

CONSENT AND ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Umeaku CN, Chris-Umeaku CI, Emmyegbe IO, Ukoha CC, Uzor UC, Agbo UJ. Proximate, Phytochemical and Antibacterial Analysis of *Persea Americana* Obtained from Nigeria. *Journal of Diseases and Medicinal Plants*. 2018;4(3):89-95.
2. Saxena M, Saxena J, Nema R, Singh D, Gupta A. Phytochemistry of medicinal plants. *Journal of Pharmacognosy and Phytochemistry*. 2013;1(6):168-182.
3. Bodeker C, Bodeker G, Ong CK, Grundy CK, Burford G, Shein K. *WHO Global Atlas of Traditional, Complementary and Alternative Medicine*. Geneva, Switzerland: World Health Organization [Google Scholar]; 2005.
4. Kunle OF, Egharevba HO, Ahmadu PO. Standardization of herbal medicines – A review. *International Journal of Biodiversity and Conservation*. 2012;4(3):101-112.
5. Sarvnarinder K, Muskaan S, Aniqqa Aniqqa, Shilpa S. Selenium attenuates bisphenol A incurred damage and apoptosis in mice testes by regulating mitogen-activated protein kinase signaling. *First international journal of andrology*. 2021;53(3):8-10. DOI:https://doi.org/10.1111/and.13975
6. Nasri H, Baradaran A, Shirzad H, Rafieian-Kopaei M. New concepts in nutraceuticals as alternative for pharmaceuticals. *International Journal of Preventive Medicine*. 2014;5(12):1483-1487.
7. Salisu A, Nura K. Phytochemical screening and antimicrobial properties of ethanolic extract of *Nymphaea lotus* L. stem. *Bayero Journal of Pure and Applied Sciences*. 2022;13(1):22-26.
8. Keke CO, Nsofor WN, Kumabia FKR, Iloabuchi GC, Ejiolor JC, Osuagwu OL. GCMS and FTIR analysis of ethanol and methanol leave extract of *Urena lobata* (Caesar weed) for bioactive phytochemical constituents. *JDDT*. 2023;13(1):99-115.
9. Miyoung P, Hyunnho C, Hana J, Heejae L. Antioxidant and anti-inflammatory activities of tannin fraction of the extract from Black Raspberry seeds compared to Grape seeds. *Journal of Food Biochemistry*. 2013;38(3):9–11.
10. Kahkonen MP, Hopia AI, Vuorela HJ, Rauha JP, Pihlaja K, Kujala KS. Antioxidant activity of plant extracts containing phenolic compounds. *Journal of Agriculture and Food Chemical*. 1999; 47:3954–3962.
11. Somayeh H, Tahoori S, Aliakbar H. *Rosa damascena* Mill. Essential oil has protective effect against testicular damage in diabetic rats. *Journal of Dietary Supplements*. 2018;15(3):311-317.
12. Muhammad A, Ali A, Idris S, Ibrahim I. Phytochemical screening and proximate analysis of Garlic (*Allium Sativum*). *Archives Organic and Inorganic Chemical Science*. 2019;4:1–7.
13. Ebomoyi M, Ahumibe K. Poster communications: Serum testosterone and testicular morphology in garlic-fed wistar rats. *Procedure Physiology Society*. 2014; 31:172-174.
14. Lakshmi JN, Babu AN, Kiran SS, Nori LP, Hassan N, Ashames A, Shaik AB. Herbs as a source for the treatment of polycystic ovarian syndrome: A systematic review. *Biotechnology*. 2023;12(1):4.
15. Shoushtari MS, Majid A, Nejadstari T, Moin M, Kardar GA. Novel report of the phytochemical composition from *Fraxinus excelsior* pollen grains. *Journal of Applied Botany and Food Quality*. 2018;91: 310-320.
16. Abedi A, Parvis M, Karimian SM, Rodsari HRS. The effect of aqueous extract of *Phoenix dactylifera* pollen grain in sexual behaviour of male rats. *Journal of Physiology Pharmaceutical Advancement*. 2012;2:235-242.
17. Yusuf AA, Lawal B, Yusuf MA, Omonije YO, Adejoke AO, Rij FH, Wenawo DL. Free radical scavenging, antimicrobial activities and effect of sub-acute exposure to Nigerian *Xylopiya aethiopica* seed extract on liver and kidney functional indices of albino rat. *Iran Journal Toxicity*. 2018; 12(3):51-58.
18. Ali S, Khan MR, Sajid M, Zahra Z. Phytochemical investigation and antimicrobial appraisal of *Parrotiopsis jacquemontiana* (Decne) Rehder. *BMC Complementary Alternative Medicine*. 2018;18(1):43.
19. Adusei S, Prince O, Richard QM Emmanuel M. Phytochemical analysis, antioxidant and metal chelating capacity of *Tetrapleura tetraptera* National library of medicine. National Center for Biotechnology Information. 2019;5(11): 27-62.
20. Adelakun SA, Ogunlade B, Olawuyi TS, Ojewale AO. Aqueous extract of *Tetrapleura tetraptera* fruit peels influence copulatory behavior and maintain testicular

- integrity in sexually mature male Sprague-Dawley rats: Pro-fertility evaluation and histomorphometry evidence. *Current Research Physiology*. 2021;19(4):7-16.
21. Okam PC, Okam CF, Obi E, Uneke PC. Effect of sub-chronic administration of mascum herbal pride on sperm quality of male Albino rats. *British Journal of Pharmaceutical Research*. 2016;11(4):1-5.
 22. Aprioku JS, Obianime AW. Evaluation of the effects of *Citrus aurantifolia* (lime) Juice in lead –induced Haematological and Testicular Toxicity in Rats. *Pharmacologia*. 2014;5(1):36-41.
 23. Idris BM, Dikko AAU, Yarube IU, Salim MA, Saleh MIA, Alhassan AW. Disturbances in calcium and zinc homeostasis during testicular damage induced by citrus aurantifolia juice in wistar rats. *Nigerian Journal of Physiological Sciences*. 2018;33(2):201-209.
 24. Oluwole A, Oluwafunmike SA, Olufunke D. Ethanol extract of the leaves of *Psidium guajava* Linn enhances sperm output in healthy wistar rats. *African Journal of Medicine and Medical Sciences*. 2007; 36(2):137-140.
 25. Neeta MP, Mukta N, Kulkarni B. Comparative qualitative phytochemical analysis of *Sesamum indicum* L. *International Journal of Curr. Microbiology in Applied Science Special*. 2015;2: 172–181.
 26. Jing P, Xiaomin Y, Shujuan Z, Jun C, Yihai W, Chunyu L, Xiangjiu H. Bioactive phenolics from mango leaves (*Mangifera indica* L.). *Industrial Crops Production*. 2018;111:400-406.
 27. Dixit V, Gupta P, Santosh R, Kumar S, Joshi BC. Reversible chemical sterilization: Effects of cyclohexanol administration on the testes and epididymides of male rabbit. *Indian Journal Physiology Pharmacology*. 1980;24(4):278-289.
 28. Tyagi A, Joshi BC, Kumar S, Dixit VP. Antispermato-genic activity of cyclohexanol in gerbil (*Meriones hurrianae* Jerdon) and house rat (*Rattus rattus Rufescens*). *Indian Journal Experimental Biology*. 1979; 17(12):1305-1307.
 29. Michael W, Lieberman L, Ernest D, Lykissa-Roberto R, Barrios S, Ching G, Nan-Ou U, Geeta-Kala K, Subbarao-Kala V. Cyclosiloxanes produce fatal liver and lung damage in mice environmental health perspectives. *Environmental Health Perspectives*. 1999;107(2):161-165.
 30. Aksoy M. Benzene Carcinogenicity. Boca Raton: CRC Press; 2017.
 31. Fayed MHM, Sanaa SA, Samira SA, Mohamed EE, El-Shahat S, Yasser AA. Phenotype analysis of Lymphocytes in workers with chronic benzene exposure. *Immunology Letter*. 2017;42(4):161–168.
 32. Manisha SO, Ravinder S. Toxic effects of sub lethal concentrations of dioctyl phthalate on the histology of liver of *Clarias batrachus* (Linn.). *International Journal of Bioassays*. 2016;5(9):4874–4877.

© 2023 Nwogu et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/103176>