



Cultural Variability among *Ceratocystis fimbriata* Ell. and Halst. Isolates Causing Wilt Disease of *Punica granatum* L.

Raja ^{a*}, Gururaj Sunkad ^b and Amaresh Y. S. ^b

^a Department of Plant Pathology, Agriculture College, Bheemarayanagudi-585287, UAS, Raichur, India.

^b Department of Plant Pathology, Agriculture College, UAS, Raichur-584104, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2023/v35i193752

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/105920>

Original Research Article

Received: 26/06/2023

Accepted: 01/09/2023

Published: 12/09/2023

ABSTRACT

Punica granatum L (Pomegranate) is one of the important fruit crops cultivated all over the world, particularly in the tropical and sub-tropics. It is affected by several diseases of which one of the most important disease is wilt caused by *Ceratocystis fimbriata*. In recent years, several orchards of farmers have been severely infected by wilt and were removed in Karnataka state. This may be due to changes in the pathogenic characteristics of the fungus. Very little work has been carried out on the cultural variability of *C. fimbriata* associated with *Punica granatum* L (Pomegranate) wilt in Karnataka. Moreover, cultural variability is the property of an organism to change its character from one generation to the other. Therefore, there is a need to study on cultural variability of *C. fimbriata*. Cultural variability showed variation among the *C. fimbriata* isolates. On the basis of colony color, type of colony growth, type of margin, margin color, and colony growth, fifty isolates were categorized.

*Corresponding author: E-mail: rajarc8888@gmail.com;

Keywords: Pomegranate; *Ceratocystis fimbriata*; cultural variability; isolates; colony color.

1. INTRODUCTION

Punica granatum L (Pomegranate) is an ancient fruit, that belongs to the family Lythraceae. Pomegranate is native to Iran, where it was first cultivated in about 2000 BC and spread to the Mediterranean countries. It is cultivated in India, Iran, China, Turkey, USA, Spain, Azerbaijan, Armenia, Afghanistan, Uzbekistan, the Middle East, Pakistan, Tunisia, Israel, dry regions of Southeast Asia, Peninsular Malaysia, the East Indies, and tropical Africa. Pomegranate cultivation is expanding globally because to its hardiness, adaptability, drought tolerance, increased output levels with outstanding keeping quality, and remunerative prices in both domestic and export markets. It thrives well in dry tropics and sub-tropics and comes up very well in soils of low fertility status as well as in saline soils. India is the world's leading country in pomegranate production. It is one of the most adaptable subtropical fruit crops. In India it is regarded as a "vital cash crop", extensively grown in Maharashtra, Karnataka, Andhra Pradesh, Telangana, and Gujarat, and is picking up fast in Himachal Pradesh, Rajasthan, and Madhya Pradesh. Small areas are under cultivation in Tamil Nadu, Mizoram, Odisha, Nagaland, Lakshadweep, Jharkhand and Jammu Kashmir. total area under pomegranate in India is 1,80,640 ha out of which 1,28,650 ha is in Maharashtra only. The total production in India is 17,89,310 metric tons and 11,97,710 metric tons in Maharashtra. In Karnataka, the total area is 23,230 ha with a production of 2,61,820 metric tonnes (<http://nhb.gov.in>)

"In Karnataka, the crop has spread across different districts viz., Vijayapura, Bagalkot, Koppal, Yadgir, Raichur, Ballari, Chitradurga, Tumakuru and Hassan. The most popular varieties suitable for processing and table use are Ganesh, Mridula, Arakta, Bhagwa (Kesar), G-137 and Khandar. Successful cultivation of pomegranates in recent years has been threatened by different pests and diseases. Bacterial blight, wilt, anthracnose, leaf spot, and root-knot nematode are important diseases. Among them, wilt caused by *Ceratocystis fimbriata* Ell. and Halst. is an emerging threat. At present the crop is severely affected by wilt pathogen and day by day the wilting severity is increasing at a faster rate. It was first noticed in some areas of Vijayapur districts of India during 1990" [1]. "By 1993, the rapid spread of this

disease was observed in the entire Vijayapura district. The cause was not identified until 1995; however, in 1996 the fungus *C. fimbriata* was isolated from discolored stem, root, and branch tissues on wilting plants. Disease is characterized by initial symptoms of yellowing and wilting of leaves on one to several branches leading to the death of affected plants in a few weeks. Cross sections of diseased plants revealed brown discoloration in the outer xylem from the roots to the main trunk" [1].

"The disease is prevalent in parts of Maharashtra, Karnataka, Telangana, Gujarat, and Tamil Nadu states" [2]. "Despite many factors conducive to the high severity, seedlings' selection for planting, soil-borne nature, and also an association with shot hole borer and plant parasitic nematodes is noticed. This might be the reason for the current rampant spread of the disease in south Indian states. Several agents are known to cause wilt in pomegranate, but *C. fimbriata* is the major cause" [3,4], Hence, emphasis given on *C. fimbriata* very little work has been carried out on the cultural variability of *C. fimbriata* associated with *Punica granatum* L (Pomegranate) wilt in Karnataka. Moreover, cultural variability is the property of an organism to change its character from one generation to the other. Therefore, there is a need to study on cultural variability of *C. fimbriata*.

2. MATERIALS AND METHODS

2.1 Cleaning of Glassware

Borosil and Corning glassware were used for all the laboratory experimental studies. They were kept for a day in a cleaning solution, prepared by dissolving 60 g of potassium dichromate ($K_2Cr_2O_7$), and 60 ml of concentrated sulphuric acid (H_2SO_4) in one liter of water. Each of these chemicals dissolved separately in 500 ml of water and finally mixed. Then glassware was cleaned by washing it with a detergent solution followed by tap water and finally rinsing in distilled water.

2.2 Sterilization

All the glassware used in the study wrapped were sterilized in an autoclave at 15 p.s.i pressure for 20 minutes and kept for drying in a hot air oven at 60°C for two hours. Both solid and liquid media were sterilized at 15 p.s.i pressure for 15 minutes.

2.3 Isolation of the Pathogen

"*Ceratocystis fimbriata*, associated with wilt was isolated from the infected roots of the pomegranate plant which were collected from Ganjalli field. The sliced pieces of collected stem portions with characteristic symptoms of vascular staining were surface sterilized with 1 per cent NaHCO₃ (sodium hypochlorite) for about 2 minutes and washed in alcohol (70%) and twice with sterile water to remove traces of NaHCO₃. Pathogen isolation was made using the carrot bait technique in which, stems were placed in between the carrot disks kept in a humid chamber, and incubated at 25 ± 2°C under a 12-hour photoperiod" [5]. "After perithecium formation, a portion of the fungi was transferred to freshly prepared PDA and oat meal agar media to allow the full development of fungi. In order to confirm the identity of the fungus, the ascospores, aleuroconidia, endoconidia, and perithecia were observed under the high power (40x) microscope from Raichur isolates the pure culture. The identification of studies of pathogens has been done as explained" by Sharma et al. [4].

2.4 Hyphal Tip Isolation

"This method was followed to maintain pure culture. Hyphal tip isolation was done on water plates. Dilute spore suspension of the pathogen was prepared in sterilized distilled water containing eight to ten spores per ml from a 15-day-old culture. One ml of such suspension was spread uniformly on two per cent solidified water agar plates and observed for spores under the microscope. Single spore was marked with a marker on the backside of the Petri plate and it was allowed to germinate. Such plates were periodically observed for spore germination under the microscope. The hyphae growing from each cell of the single spore were traced and marked with a marker. The tip of the hyphae was cut carefully transferred to PDA plates and incubated at 25 ± 2°C for 15 days. Later, mycelial bits of the fungus were transferred to the center of Petri plates containing PDA and incubated at 25 ± 2°C for 15 days. Saltation or sectoring was observed in the culture to confirm the pure culture of the fungus" [6].

2.5 Maintenance of the Culture

The hyphal tip cultures of the fungus were sub-cultured on potato dextrose agar slants and kept in the laboratory at 25 ± 2°C for 15 days. Such

mother culture slants were preserved at 5°C in the refrigerator. Further, these cultures were sub-cultured once in a month and used for future studies.

2.6 Morphological Characters *C. fimbriata*

"*C. fimbriata* was characterized by the production of aleurioconidia, endoconidia, ascospore and perithecia. For this, the growth of Cf-26 was selected from 21 days old pure culture and kept on a clean sterile glass slide using a sterilized needle. With the help of a fluorescent microscope, the length and breadth of aleurioconidia, endoconidia, ascospore, and perithecia in µm were measured. Three observations were recorded from the pure culture of the fungus. Ten aleurioconidia, endoconidia, ascospores, and perithecia were picked up randomly to determine the diameter, and *C. fimbriata* (Cf-26) was characterized for colony color and growth pattern on oat meal agar. The mycelial disc of 5 mm diameter was cut from the periphery of the actively growing culture of Cf-26 and transferred aseptically to a 90 mm Petri dish containing 20 ml of oat meal agar and incubated for a period till the fungal growth covered the complete petri plate in the media at 26 ± 2°C. The colony was characterized for phenotype and growth pattern. Different morphotypes of colony color, type of colony, type of margin, margin color, and colony growth were observed *in vitro*" [6].

2.7 Cultural Characters of *C. fimbriata* on Oat Meal Agar

The cultural characteristics of *C. fimbriata* were studied on oat meal agar. The composition and preparation of the above-mentioned media were obtained from Ainsworth and Bisby's "Dictionary of the Fungi" by Hawksworth et al. [7]. The composition and preparation of the media are as follows

Oat meal agar:

Oat meal powder : 40.00 g
Agar - agar : 20.00 g
Distilled water : 1000 ml (volume to make up)

The oat meal powder was dissolved in 500 ml of distilled water and Agar-agar was melted in 500 ml of distilled water separately. Both solutions were mixed thoroughly. The volume was made up to one liter and sterilized.

Twenty ml of each medium was poured into the Petri dishes for solidification. Five mm discs of *C. fimbriata* were placed at the centre of the plate. Each set of experiments was replicated thrice and plates were incubated at $26 \pm 2^\circ\text{C}$. Observations were taken on parameters such as growth type, mycelial color, type of margin, and radial growth (mm) when the fungus covered a complete Petri plate in the media. The results were analyzed statistically.

Studies on cultural variability among the isolates of *C. fimbriata* were carried out during the study. Fifty samples were collected from nine pomegranate-growing districts of Karnataka during the survey. The isolates were obtained by tissue isolation using the carrot bait technique followed by inoculation on oat meal agar. Fifty isolates were obtained from such samples and designated as Cf-1 to Cf-50 for variability studies (Table 1).

Table 1. Designation of *C. fimbriata* isolates of pomegranate wilt collected from different districts of Karnataka

Sl. No.	Name of the Place		Designation of the isolate
	District	Village	
1	Viajyapura	Kumtagi	Cf-1
2		Babaleshwar	Cf-2
3		Hittinhalli	Cf-3
4		Jumnal	Cf-4
5		Kannollo-1	Cf-5
6		Devara hippargi-1	Cf-6
7		Bandal	Cf-7
8	Bagalkot	Devanal	Cf-8
9		Govindkoppa	Cf-9
10		Kaladgi-1	Cf-10
11		Lokapur-1	Cf-11
12	Koppal	Mahalingapur-1	Cf-12
13		Kalkbandi	Cf-13
14		Kamanur	Cf-14
15		Kustgi	Cf-15
16		Maladgatti-1	Cf-16
17		Kodkera	Cf-17
18	Yadgir	Gogi K	Cf-18
19		Wandurga-1	Cf-19
20		Tumkur	Cf-20
21		Heggandoddi-1	Cf-21
22	Raichur	Chincholi-1	Cf-22
23		Yatgal	Cf-23
24		Chandrabanda	Cf-24
25		Karekal	Cf-25
26		Ganjhalli-1	Cf-26
27		Kurkihalli	Cf-27
28		Benkal	Cf-28
29	Ballari	Arkera-1	Cf-29
30		Kampli	Cf-30
31		Lakshmipura	Cf-31
32		Khondanhalli	Cf-32
33	Chitradurga	Thambrahalli	Cf-33
34		Basarkodu	Cf-34
35		Sirana hatti-1	Cf-35
36		Ramajjanahalli	Cf-36
37		Nagayana hatti-1	Cf-37
38		Maskal-1	Cf-38

Sl. No.	Name of the Place		Designation of the isolate
	District	Village	
39		Seerana katte-1	Cf-39
40		Shrarianagar	Cf-40
41	Tumakur	Madana kunte-1	Cf-41
42		Karekyatana halli	Cf-42
43		Chikka halikute-1	Cf-43
44		Thogargunte-1	Cf-44
45		Hosahali	Cf-45
46	Hassan	Mylanahalli-1	Cf-46
47		Nadakhalli	Cf-47
48		Chika bidane-1	Cf-48
49		Haranhalli-1	Cf-49
50		Goran koppal-1	Cf-50

3. RESULTS

3.1 Pathogen

On carrot bait followed by oat meal agar, the growth of *C. fimbriata* started after 3-4 days and the mycelium was a whitish-grey color which changed to brown colour. As the growth progressed, the production of endoconidia, aleurioconidia, and perithecium was observed. The black-colored perithecia with a globose base were observed, exuding small, hyaline, and hat-shaped ascospores from the apex of the perithecium neck. The endoconidia were hyaline, cylindrical, and formed endogenously in hyphae and aleurioconidia were thick-walled ellipsoidal or pyriform, golden-brown in colour. They are borne singly or in chains.

3.2 Isolation and Identification

Standard tissue (Carrot bait technique followed by oat meal agar) isolation was followed to isolate *Ceratocystis fimbriata* culture from a diseased sample of infected root with the typical symptom of dark grayish-brown streaks on splitting of root portion, collected from pomegranate field. Within 3-4 days after on carrot bait the white cottony growth was observed. Later 5-6 days black color perithecia were observed when carrot the culture was transformed on oat meal agar. The pure culture was maintained on oat meal agar at $28 \pm 2^\circ\text{C}$. Sub-culturing was done at every fortnight interval. The fungus isolated was confirmed as *C. fimbriata* based on their cultural and morphological characteristics.

Table 2. Cultural characteristics of different isolates of *C. fimbriata* on oat meal agar

Sl. No.	Isolate	Colony color	Type of colony growth	Type of margin	Margin color	Colony growth (mm)
1	Cf-1	Grayish	Flat	Regular	Light gray	90
2	Cf-2	Grayish	Flat	Regular	Light gray	90
3	Cf-3	Brown	Flat	Regular	Brown	88
4	Cf-4	Grayish	Flat	Regular	Light gray	90
5	Cf-5	Grayish	Flat	Regular	Light gray	83
6	Cf-6	Grayish	Flat	Regular	Light gray	90
7	Cf-7	Light gray	Flat	Regular	Light gray	90
8	Cf-8	Brown	Flat	Regular	Brown	85
9	Cf-9	Grayish	Fluffy	Irregular	Light gray	90
10	Cf-10	Brown	Flat	Regular	Brown	80
11	Cf-11	Grayish	Fluffy	Regular	Light gray	89
12	Cf-12	Light gray	Flat	Regular	Light gray	81
13	Cf-13	Grayish	Flat	Irregular	Light gray	90
14	Cf-14	Grayish	Flat	Regular	Light White	90
15	Cf-15	Light gray	Flat	Regular	Light white	70
16	Cf-16	Light gray	Fluffy	Irregular	Light gray	70
17	Cf-17	Grayish	Flat	Regular	Light gray	90

Sl. No.	Isolate	Colony color	Type of colony growth	Type of margin	Margin color	Colony growth (mm)
18	Cf-18	Grayish	Flat	Regular	Light gray	85
19	Cf-19	Grayish	Flat	Regular	Light White	90
20	Cf-20	Grayish	Flat	Regular	Light gray	88
21	Cf-21	Light gray	Flat	Regular	Light gray	90
22	Cf-22	Brown	Flat	Regular	Brown	81
23	Cf-23	Grayish	Flat	Regular	Light gray	90
24	Cf-24	Grayish	Fluffy	Irregular	Light gray	90
25	Cf-25	Grayish	Flat	Regular	Light white	90
26	Cf-26	Brown	Flat	Regular	Brown	90
27	Cf-27	Grayish	Flat	Regular	Light gray	82
28	Cf-28	Light gray	Flat	Regular	Light gray	90
29	Cf-29	Light gray	Flat	Regular	Light gray	83
30	Cf-30	Brown	Fluffy	Irregular	Light gray	90
31	Cf-31	Grayish	Fluffy	Irregular	Light gray	70
32	Cf-32	Grayish	Flat	Regular	Light gray	90
33	Cf-33	Brown	Flat	Regular	Brown	86
34	Cf-34	Grayish	Flat	Regular	Light gray	90
35	Cf-35	Grayish	Flat	Regular	Light white	90
36	Cf-36	Light gray	Flat	Regular	Light gray	81
37	Cf-37	Grayish	Flat	Regular	Light gray	85
38	Cf-38	Grayish	Flat	Regular	Light gray	70
39	Cf-39	Grayish	Fluffy	Irregular	Light gray	90
40	Cf-40	Grayish	Flat	Regular	Light gray	84
41	Cf-41	Grayish	Flat	Regular	Light white	90
42	Cf-42	Brown	Flat	Regular	Brown	90
43	Cf-43	Grayish	Flat	Regular	Light gray	79
44	Cf-44	Grayish	Fluffy	Irregular	Light gray	70
45	Cf-45	Brown	Flat	Regular	Light gray	90
46	Cf-46	Grayish	Flat	Regular	Light gray	78
47	Cf-47	Grayish	Flat	Regular	Light gray	90
48	Cf-48	Grayish	Flat	Regular	Light gray	90
49	Cf-49	Light gray	Flat	Regular	Light gray	90
50	Cf-50	Light gray	Flat	Regular	Light gray	90

3.3 Cultural Variability of Different Isolates of *C. fimbriata*

To study the variability, cultural characters such as colony color, type of colony growth, type of margin, margin color, and colony growth of fifty isolates were assessed on Oat meal agar as described in 'Material and Methods' and results are presented in Table 2, Table 3 and Plate 1.

Diversity in cultural characters on oat meal agar at room temperature showed variation among the *C. fimbriata* isolates, such as colony color (grayish, brown, and light gray), type of colony growth (flat and fluffy), type of margin (regular and irregular) margin color (light gray and brown) and colony growth in mm were

closely observed in 50 isolates of *C. fimbriata*. Most of the isolates showed a grayish color with a flat type of colony growth and regular margin. The margin color was light gray in many isolates and colony growth ranged from 70 to 90 mm.

3.4 Grouping of *C. fimbriata* Isolates Based on Cultural Characteristics

Diversity in cultural characters such as colony color (grayish/brown/light gray), type of colony growth (flat/fluffy), type of margin (regular/irregular) margin color (light gray/brown), and colony growth in mm were closely observed in 50 isolates of *C. fimbriata* and categorized as described in Table 3a and Table 3b.

Table 3a. Categorization of isolates based on colony color, type of colony growth and type of margin

i. Based on colony color

Colony color	Isolate number	Total
Grayish	Cf-1, Cf-2, Cf-4, Cf-5, Cf-6, Cf-9, Cf-11, Cf-13, Cf-14, Cf-17, Cf-18, Cf-19, Cf-20, Cf-22, Cf-24, Cf-25, Cf-26, Cf-27, Cf-31, Cf-32, Cf-34, Cf-35, Cf-37, Cf-38, Cf-39, Cf-40, Cf-41, Cf-43, Cf-44, Cf-46, Cf-47, Cf-48	32
Light gray	Cf-7, Cf-12, Cf-15, Cf-16, Cf-21, Cf-28, Cf-29, Cf-36, Cf-49, Cf-50	10
Brown	Cf-3, Cf-8, Cf-10, Cf-23, Cf-30, Cf-33, Cf-42, Cf-45	08
Total		50

ii. Based on type of colony growth

Topography	Isolate number	Total
Fluffy growth	Cf-9, Cf-11, Cf-16, Cf-25, Cf-30, Cf-31, Cf-39, Cf-44	8
Flat	Cf-1, Cf-2, Cf-3, Cf-4, Cf-5, Cf-6, Cf-7, Cf-8, Cf-10, Cf-12, Cf-13, Cf-14, Cf-15, Cf-17, Cf-18, Cf-20, Cf-21, Cf-22, Cf-23, Cf-24, Cf-26, Cf-27, Cf-28, Cf-29, Cf-32, Cf-33, Cf-34, Cf-35, Cf-36, Cf-37, Cf-38, Cf-40, Cf-41, Cf-42, Cf-43, Cf-45, Cf-46, Cf-47, Cf-48, Cf-49, Cf-50	42
Total		50

iii. Based on type of margin

Type of margin	Isolate number	Total
Irregular	Cf-9, Cf-13, Cf-16, Cf-25, Cf-30, Cf-31, Cf-39, Cf-44	8
Regular	Cf-1, Cf-2, Cf-3, Cf-4, Cf-5, Cf-6, Cf-7, Cf-8, Cf-10, Cf-11, Cf-12, Cf-14, Cf-15, Cf-17, Cf-18, Cf-19, Cf-20, Cf-21, Cf-22, Cf-23, Cf-24, Cf-26, Cf-27, Cf-28, Cf-29, Cf-32, Cf-33, Cf-34, Cf-35, Cf-36, Cf-37, Cf-38, Cf-40, Cf-41, Cf-42, Cf-43, Cf-45, Cf-46, Cf-47, Cf-48, Cf-49, Cf-50	42
Total		50

Table 3b. Categorization of isolates based on margin color and rate of growth

i. Based on margin color

Color	Isolate number	Total
Light gray	Cf-1, Cf-2, Cf-4, Cf-5, Cf-6, Cf-7, Cf-9, Cf-11, Cf-12, Cf-13, Cf-16, Cf-17, Cf-18, Cf-20, Cf-21, Cf-22, Cf-24, Cf-25, Cf-27, Cf-28, Cf-29, Cf-30, Cf-31, Cf-32, Cf-34, Cf-36, Cf-37, Cf-38, Cf-39, Cf-40, Cf-43, Cf-44, Cf-45, Cf-46, Cf-47, Cf-48, Cf-49, Cf-50	38
Brown	Cf-3, Cf-8, Cf-10, Cf-23, Cf-33, Cf-42	6
Light white	Cf-14, Cf-15, Cf-19, Cf-26, Cf-35, Cf-41	6
Total		50

ii. Based on rate of growth

Rate of growth	Isolate number	Total
I-Slow growing (<50 mm growth)	-	0
II-Moderately growing (51-70 mm growth)	Cf-15, Cf-16, Cf-31, Cf-38	4
III-Fast growing (71-90 mm growth)	Cf-1, Cf-2, Cf-3, Cf-4, Cf-5, Cf-6, Cf-7, Cf-8, Cf-9, Cf-10, Cf-11, Cf-12, Cf-13, Cf-14, Cf-17, Cf-18, Cf-19, Cf-20, Cf-21, Cf-22, Cf-23, Cf-24, Cf-25, Cf-26, Cf-27, Cf-28, Cf-29, Cf-30, Cf-32, Cf-33, Cf-34, Cf-35, Cf-36, Cf-37, Cf-39, Cf-40, Cf-41, Cf-42, Cf-43, Cf-44, Cf-45, Cf-46, Cf-47, Cf-48, Cf-49, Cf-50	46
Total		50




Plate 1. Cultural characteristics of different isolates of *C. fimbriata* on oat meal agar
1=Cf-1, 2=Cf-2, 3=Cf-3, 4=Cf-4, 5=Cf-5, 6=Cf-6, 7=Cf-7, 8=Cf-8, 9=Cf-9, 10=Cf-10, 11=Cf-11, 12=Cf-12, 13=Cf-13, 14=Cf-14, 15=Cf-15, 16=Cf-16, 17=Cf-17, 18=Cf-18, 19=Cf-19, 20=Cf-20, 21=Cf-21, 22=Cf-22, 23=Cf-23, 24=Cf-24, 25=Cf-25




Plate 1. Cultural characteristics of different isolates of *C. fimbriata* on oat meal agar (Contd.)

26=Cf-26, 27=Cf-27, 28=Cf-28, 29=Cf-29, 30=Cf-30, 31=Cf-31, 32=Cf-32, 33=Cf-33, 34=Cf-34, 35=Cf-35, 36=Cf-36, 37=Cf-37, 38=Cf-38, 39=Cf-39, 40=Cf-40, 41=Cf-41, 42=Cf-42, 43=Cf-43, 44=Cf-44, 45=Cf-45, 46=Cf-46, 47=Cf-47, 48=Cf-48, 49=Cf-49, 50=Cf-50

4. DISCUSSION

In nature, new strains may arise by mutation, hybridization, differential cytoplasmic inheritance [8] heterokaryosis [9], and parasexual life cycle [10]. The study of pathogenic variability is essential for breeding disease resistance in crop improvement programs. A potential pathogen is often blessed with biodiversity within its population. Basically, variation in the pathogen is a desirable trait for its existence in nature. This variability among the pathogens underlies their diverse nature and ability to withstand the host environment. The variability of pathogens was studied with cultural, morphological, and molecular to focus on the existence of variation in *C. fimbriata* collected from different locations.

In the present investigation, diversity in cultural characters showed variation among the *C. fimbriata* isolates. The variation was observed with respect to colony color (ranged from grayish, brown to light gray), type of colony growth (flat and fluffy), type of margin (regular to irregular) margin color (light gray to brown), and colony growth were closely observed in fifty isolates of *C. fimbriata*.

In the present investigation, based on colony color, isolates were categorized into three groups: grayish, light gray, and brown. Thirty-two isolates showed grayish, ten isolates light gray and eight isolates showed brown color. Similar results with respect to variation in margin color were reported by several workers [11,12]. Lal and Kandhari [13] reported that “six isolates were light brown, five isolates were found yellowish brown, four isolates were whitish brown in color, six isolates were dark brown and four isolates were very pale brown in the case of *R. Solani*”. “Further, nineteen isolates of *Fusarium oxysporum* f.sp. *zingiberi* causal organism of Fusarium yellows in ginger was collected from different ginger growing areas of Himachal Pradesh and designated as I1 to I19. Morphological variations with respect to mycelial color, conidial size, formation of chlamydospores, and pathogenic variation in terms of disease incidence among different isolates were studied in these isolates. The mycelial color varied from white to dull white with a slightly pinkish tinge” [11].

Based on colony growth, isolates were categorized into two groups: fluffy and flat in the present study. Forty-two isolates showed flat growth and colonies of eight isolates fluffy

growth. Similar results with respect to variation in colony growth were reported by several workers [11,14,15,16,17,12]. Prasad et al. [18] studied the variation among the isolates of *S. rolfsii* collected from groundnut and reported that out of 20 isolates, colonies of 11 isolates showed fluffy growth, whereas 9 isolates were flat. Ravi Chandran and Reddi Kumar [19] studied “the variation among the isolates of *Fusarium solani* (Mart.) Sacc., collected from citrus and reported that *F. solani* isolates, AFS1, AFS2, AFS4, CFS9, and CFS13 grew more than 85 mm after 7 days of inoculation and considered as a fast-growing category. Among all the isolates of *F. solani* CFS9 isolate showed significant variation in radial growth (90 mm) on the PDA medium”.

Based on the type of margin, isolates were categorized into two groups: irregular and regular, in the present study. Forty-two isolates showed the regular type of margin, and eight isolates were irregular. Similar results with respect to variation in the type of margin were reported by Manashi Debbarma and Pranab Dutta [15] in the case of variability *R. solani* in which they reported that Colony size, colony growth, color and texture (smooth or rough) varied in six isolates they studied.

With respect to diversity in margin color, isolates were categorized into three groups: light gray, brown, and light white. Thirty-eight isolates showed light gray color, eight isolates brown color, and six isolates light white color. Similar results with respect to variation in margin color were reported by several workers [11,12]. Further, Lal and Kandhari [13] reported that “six isolates were light brown, five isolates were found yellowish brown, four isolates were whitish brown in color, six isolates were dark brown and four isolates were very pale brown in the case *R. Solani*”.

The rate of growth was closely observed in fifty isolates of *C. fimbriata*. Isolates were found fast-growing, moderate, and slow-growing. Forty-six isolates were fast-growing with an average growth rate (71-90 mm), four intermediate-growing isolates with an average growth rate (51-70mm), and none of the isolates were slow-growing with an average growth rate of <50 mm. The findings are in accordance with Prabhu and Patil [20] who reported that among twelve isolates of *S. rolfsii* 11 are fast growing and one isolate is moderate in growth. Further, Hussain et al. [21] classified the isolates of *S. rolfsii* based on morphological variation into fast-growing, moderately, and slow-growing and reported that

the isolates AT-1, AT-2, and RW-2 represented significantly fast-growing, isolates SR-1, CH-1 and DL-2 intermediate and SR-2, CH-2, CH-3, DL-1, AT-3, and RW-1 under slow radial colony growth and Saxena et al. [22] also classified “the isolate based on the growth rate was considered the main criterion for grouping the *Fusarium oxysporum* f.sp. *lentis* isolates. Fast-growing isolates comprised fourteen groups, those having large macroconidia formed six groups, while small size conidia were in eight groups. In the medium growth rate, there were seventeen groups, out of these eight groups had large macro conidia, while in nine groups macro conidia were small. In the slow growth rate, there were nineteen groups, those having large macro conidia were eight, while in eleven groups small macro conidia were present”.

5. CONCLUSIONS

Cultural variability studied on oat meal agar showed variation among *C. fimbriata* isolates. On the basis of colony color, type of colony growth, type of margin, margin color, and colony growth, fifty isolates were categorized viz., based on colony color of mycelia among fifty isolates *C. fimbriata*, thirty-two isolates were found grayish color, ten isolates, light gray color, and brown colony color in remaining eight isolates. Based on the type of colony growth, eight isolates showed fluffy growth and fifty-two isolates showed the flat type of colony growth. With respect to the type of margin, eight isolates were irregular and forty-two isolates showed the regular type of margin. Based on margin color, three categories of isolates, light gray in thirty-eight isolates, brown in six isolates, and light white in the remaining six isolates, and based on the rate of growth, four isolates were found moderately growing (51-70 mm), forty-six were fast growing (71-90 mm) and none of the isolates were slow growing (< 50 mm) covering the oat meal agar within 13-16 days.

ACKNOWLEDGEMENT

The authors wish to thank financial contributions from the University Grants Commission-Rajiv Gandhi National Fellowship (UGC-RGNF), Farmer, and the University of Agricultural Sciences, Raichur (UASR).

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Somasekhara YM, Wali SY. Survey of incidence of pomegranate (*Punica granatum* Linn) wilt *Ceratocystis fimbriata* (Ell & Halst). Orissa J. Hort. 2000;28:84-89.
2. Jadhav VT, Sharma KK. Integrated management of disease in pomegranate. Paper Presented In: 2nd Inter. Symp. Pomegranate and minor including Mediterranean Fruits, Univ. Agric. Sic. Dharwad. June 23-27 2009;48-52.
3. Sharma KK. Vascular wilt of pomegranate caused by *Ceratocystis fimbriata* Ellis and Halsted and its control. 5th International conference on Plant Pathology in the globalized Era, at IARI, New Delhi: Nov 10-13 2009:240.
4. Sharma KK, Sharma J, Jadhav VT. Etiology of pomegranate wilt and its management. In: Fruit, vegetable, cereal science and biotechnology. Global Science Books. 2010;4(2):96-101.
5. Moller WJ, Devay JE. Carrot as a species selective isolation medium for *Ceratocystis fimbriata*. Phytopathol. 1968;58:123.
6. Raja SG, Amaresh YS, Yenjerappa ST, Amaregouda A, Shreenivas AG. Morphological characteristics of *Ceratocystis fimbriata* Ell. and Halst. causing wilt in pomegranate. Int. J. Pure Appl. Biosci. 2017;5:285-9.
7. Hawksworth DL, Sutton BC, Ainsworth GC. Ainsworth & Bisby's Dictionary of the Fungi, 7th edn. Commonwealth Agricultural Bureaux, Slough; 1983.
8. Worthington CJ, Horn BW, Moore GG, Monacell JT, Singh R, Stone EA. Hybridization between *Aspergillus flavus* and *Aspergillus parasiticus*, in Fungal Genetics Conference Proceedings, Pacific Grove, CA. 2011;82-84.
9. Milgroom MG, Sotirovski K, Risteski M, Brewer MT. Heterokaryons and parasexual recombinants of *Cryphonectria parasitica* in two clonal populations in southeastern Europe. Fungal Genet. Biol. 2009;46:849-854.
10. Rosada LJ, Franco CCS, San Anna JR, Kaneshima EN, Gonçalves-Vidigal, MC, Castro-Prado, MAA. Parasexuality in race 65 *Colletotrichum lindemuthianum* isolates. J. Eukaryot. Microbiol. 2010;57:383-384.
11. Gupta Meenu, Jarial Kumud, Vikram Amit. Morphological, cultural, pathological and molecular variability among *Fusarium*

- oxysporum* f.sp. *zingiberi* isolates. International Journal of Bio-resource and Stress Management. 2014;5(3):375-380.
12. Sonyal S, Madhu, Mahesh HS, Palanna B, Manjunath SH, Papachan A. Cultural characteristics of *Ceratocystis fimbriata* Ell. and Halst. causing wilt in pomegranate. Int. J. Pure App. Biosci. 2015;3(4):23-27.
 13. Lal M, Kandhari J. Cultural and morphological variability in *Rhizoctonia solani* isolates causing sheath blight of rice. J. Mycol. Pl. Pathol. 2009;39(1):77-81.
 14. Kumar M, Madhavi Santhoshi, Giridhara Krishna M, T, Raja Reddy K. Cultural and morphological variability of *Sclerotium rolfsii* isolates infecting groundnut and its reaction to some fungicides. Int. J. Curr. Microbiol. App. Sci. 2014;3(10):553-561.
 15. Manashi Debbarma, Pranab Dutta. Cultural and morphological variability in *Rhizoctonia solani* isolates of different hosts of Assam. Indian J. Appl. Res. 2015; 5(2):878-884.
 16. Poornima, Investigations on stem and pod rot of groundnut caused by *Sclerotium rolfsii* Sacc. Ph. D. Thesis, Univ. Agril. Sci., Raichur. 2015;186-221.
 17. Rasu T, Sevugapperumal N, Thiruvengadam R, Ramasamy S. Morphological and genomic variability among *Sclerotium rolfsii* populations. The Bioscan. 2013;8(4):1425-1430.
 18. Prasad SD, Basha ST, Reddy NPGE. Molecular variability among the isolates of *Sclerotium rolfsii* causing stem rot of groundnut by RAPD, ITS-PCR and RFLP. Eur. Asia J. Bio. Sci. 2010;4:80-87.
 19. Ravi Chandran M, Reddi Kumar M. Studies on cultural, morphological variability in isolates of *Fusarium solani* (Mart.) Sacc., incitant of dry root-rot of Citrus. Curr. Biotica. 2012;6(2):152-162.
 20. Prabhu K, Patil I. Morphological, biochemical and pathogenic variation among *Sclerotium rolfsii* isolates of soybean, Karnataka J. Agric. Sci. 2005;18(4):990-994.
 21. Hussain M. Zakir, Rahman MA. Mohammad Nurul Islam, Latif MA, Bashar MA. Morphological and molecular identification of *Fusarium oxysporum* SCH isolated from guava wilt in Bangladesh. Bangladesh J. Bot. 2012;4(1):49-54.
 22. Saxena DR, Moly Saxena, Neha Tiwari. Morphological and cultural variability in *Fusarium oxysporum* f.sp. *lentis* causing wilt of lentil. Indian Phytopathol. 2019;72: 665–673.

© 2023 Raja et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/105920>