



Agronomic Efficiency of Hairy Woodrose (*Merremia aegyptia* L.) Mixed with Cattle Manure in the Productivity of Coriander Cultivars in the Semiarid Region of Brazil

**Paulo César Ferreira Linhares^{a++*},
Maria Elisa da Costa Souza^{a#},
Domingos Severino de Souza Júnior^{a#},
Thiffany Louhanne Alves Dantas^{a#},
Maria Eduarda Pereira dos Santos^{a#},
Adma Larissa da Silva^{a#},
Antônia Vilma de Andrade Ferreira Amâncio^{b†},
José Ferreira Amâncio^{c‡},
Roberto Pequeno de Sousa^{c^},
Eudes de Almeida Cardoso^{c^},
Walter Martins Rodrigues^{c^},
Joelma de Assis França^{d##}
and Isaac Reinaldo Pinheiro de Lima^{d##}**

^a Jitirana Research Group, Federal Rural University of the Semi-Arid, Mossoró, RN, Brazil.

^b Federal Rural University of the Semi-Arid and Member of the Jitirana Research Group, Mossoró, RN, Brazil.

^c Federal Rural University of the Semi-Arid, Mossoró, RN, Brazil.

^d Federal University of Rio Grande do Norte (UFERN), Natal, RN, Brazil.

⁺⁺ PhD Researcher, Leader;

[#] Agronomy Student and Member;

[†] Master, Researcher;

[‡] Agricultural Engineer, Specialization in Plant Protection;

[^] Professor;

^{##} Master of Science in Education;

*Corresponding author: E-mail: paulolinhares@ufersa.edu.br;

Authors' contributions

This work was carried out in collaboration among all authors. The execution of activities, such as planning, planting, conducting the experiment in the field, as well as writing, had the participation of all authors and was of great importance for the conclusion of the article. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJRCS/2023/v8i4223

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/108303>

Original Research Article

Received: 27/08/2023

Accepted: 02/11/2023

Published: 03/11/2023

ABSTRACT

Aims: Ecologically-based agriculture is widely applied by family farmers who grow vegetables in the semi-arid region, contributing to a great extent to the increase in production.

Place of Study: The experiment was carried out at the Rafael Fernandes Experimental Farm, belonging to the Universidade Federal Rural do Semiárido-UFERSA, with geographic coordinates of 5°03'37"S and 37°23'50"W Gr, with an altitude of 72 m, with the objective of agronomic efficiency of hairy woodrose (*Merremia aegyptia* L.) mixed with cattle manure in the productivity of coriander cultivars in the semiarid region of Brazil.

Study Design and Methodology: The experimental design used was complete randomized blocks in a 5 x 3 factorial scheme, with three replications. The first factor consisted of four amounts of the mixture of hairy woodrose plus cattle manure (0; 1.0; 2.0; 3.0 and 4.0 kg m⁻² of bed) and the second factor consisted of three cultivars of coriander (verdão, Super-verdão and tabocas) in single cultivation. The spacing used was 0.1 x 0.05 m with five plants pit⁻¹, corresponding to 1000 plants m⁻² of bed, density used by coriander producers.

Agronomic Characteristics of Coriander: Plant height, expressed in centimeter plant⁻¹; number, expressed in plant⁻¹ units; productivity, expressed in grams m⁻²; number of sauces, expressed unit m⁻² and dry mass, expressed in grams m⁻² of area.

Conclusions: The best agronomic performance was obtained in the amount of 4.0 kg m⁻², for the verdão cultivar, with an average value of 1585 g m⁻² and 31.7 units m⁻² for productivity and number of bunches, respectively.

The mixture of fertilizers of plant and animal origin is extremely important for increasing coriander production and productivity.

Keywords: Organic fertilizer; vegetables; spontaneou specie; *Coriandrum sativum* L.

1. INTRODUCTION

Ecologically-based agriculture is widely applied by family farmers who grow vegetables in the semi-arid region, contributing greatly to the increase in production.

Among vegetables, coriander stands out, a species widely cultivated by ecologically based family farmers who produce for consumption and commercialization in the semi-arid region of

Brazil, being of paramount importance for those who work in this activity.

In the semi-arid region of Brazil, this species is harvested by producers between 25 and 32 days after sowing, when the species has a good amount of leaves, which characterizes the commercial aspect. Since it is a leafy crop, coriander requires nitrogen fertilization, which increases production costs, as it demands

financial resources that are not available to farmers.

“Within this context, the semi-arid region of Brazil develops spontaneous species with the potential to be used as green manure, such as the hairy woodrose (*Merremia aegyptia* L.) which appears during the period of rainfall, with production of green and dry phytomass on the order of 40000 and 6000 kg ha⁻¹, nitrogen content of 25.4 g kg and carbon-nitrogen ratio of 23/1” [1 and 2].

When growing short-cycle vegetables, it is of great importance to use alternative sources that allow greater availability of nutrients to the soil, to be absorbed by the plants. In this sense, the mixture of fertilizers of vegetable (hairy woodrose) and animal (cattle manure) origin is extremely effective, considering that the hairy woodrose provides the greatest amount of nitrogen to the soil, due to its concentration in the matter (24.5 g kg⁻¹), in relation to phosphorus and potassium, cattle manure provides the greatest amount of these nutrients, due to its concentration (622.3 mg dm⁻³ of phosphorus and 2177.2 mg dm⁻³ of potassium), thus ensuring a better supply of nutrients for coriander cultivation.

Given the importance of mixing alternative sources of nutrients to the soil, the aim was to agronomically evaluating cultivars of coriander fertilized with green manure with hairy woodrose (*Merremia aegyptia* L.) plus cattle manure incorporated into the soil.

2. MATERIALS AND METHODS

2.1 Geographic Location of the Experiment

The experiment was conducted at the Rafael Fernandes Experimental Farm, belonging to the Universidade Federal Rural do Semi-Árido (UFERSA), in the semi-arid region of Brazil with geographic coordinates of latitude between 5° 03' 37”S and longitude between 37° 23' 50”W Gr,

with altitude above sea level of 72 m [3]. According to [4] and the Köppen classification, “the local climate is BSwH', dry and very hot, with the dry season usually from June to January, and the rainy season from February to May, with average annual precipitation of 673.9 mm and the average relative humidity is 68.9%”.

2.2 Experimental Design and Treatments

“The statistical design used was complete randomized blocks in a 5 x 3 factorial scheme, with three replications, consisting of fifteen treatments with 45 experimental plots. The first factor consisted of five amounts of hairy woodrose as green manure plus cattle manure (0.0; 1.0; 2.0; 3.0 and 4.0 kg m⁻²), the second factor consisting of three cultivars of coriander (Verdão; Super-verdão and tabocas). The spacing used was 0.10 x 0.05 m, with five plants pit⁻¹, making a population of 1000 plants m⁻², which makes it economically viable” [5].

Soil preparation consisted of harrowing, then the beds were surveyed using an agricultural machine, using a rotary tiller. During the experiment, manual weeding was carried out to keep the crop free from weed competition. The irrigations carried out were carried out by sprinkling, throughout the crop cycle, in two shifts, morning and afternoon, lasting 40 minutes each.

When the experiment was set up, the hairy woodrose was harvested in an area with predominance of the species to compose the treatments, being incorporated into the soil. Four samples were taken and sent to the soil laboratory, whose chemical composition was: 550 g kg⁻¹ of carbon (C); 24.5 g kg⁻¹ of nitrogen (N); 14.6 g kg⁻¹ of phosphorus (P); 20 g kg⁻¹ potassium (P); 10.2 g kg⁻¹ of calcium (Ca); 11.7 g kg⁻¹ of magnesium (Mg) and a carbon-nitrogen ratio of 22.4 (C/N) (Fig. 1).



Fig. 1. Hairy woodrose (*Merremia aegyptia* L.) in full flowering stage (A) and vegetative stage (B) in the semi-arid region of Brazil. Photograph: Researcher PhD Paulo César Ferreira Linhares

Cattle manure came from lactating cows raised in an intensive system with the following chemical characteristics: 5.42 g kg⁻¹ of nitrogen (N); 158.39 g kg⁻¹ of organic matter (MO); 622.3 mg dm⁻³ of phosphorus (P); 2177.2 mg dm⁻³ of potassium (K⁺); 429.9 mg dm⁻³ of sodium (Na⁺); 8.10 cmolc dm⁻³ of Ca²⁺; 0.50 cmolc dm⁻³ of Mg²⁺ and 16,04 cmolc dm⁻³ of cation exchange ability (CTC). Planting was carried out thirty days after incorporation, as recommended by [6].

At thirty-three days after sowing, the analyzes of the agronomic characteristics of the coriander crop were carried out, with the assessment of plant height being carried out in the field, measuring thirty plants in plot⁻¹, being measured with a ruler millimeter with results expressed in centimeter plant⁻¹. After harvesting in the field, other characteristics were evaluated: number of stems (by counting the leaves of a sample of thirty plants, expressed in plant⁻¹ units); productivity (weighing of all plants within the useful area of the plot was performed on a 1.0g precision scale, expressed in grams m⁻² of area); number of sauces (to determine this characteristic, the weight of the green pasta was divided by 50g, a value referring to a coriander sauce unit) and dry mass (a sample of thirty plants per experimental plot was used, weighed on a 1.0g precision scale, placed in an air circulation oven at 65°C until constant mass, the results being expressed in grams m⁻² of area).

2.3 Statistical Analysis

Statistical analysis was performed according to conventional methods of analysis of variance [7], using the ESTAT statistical software [8]. The procedure for adjusting the response curve was performed using the ESTAT software [8].

3. RESULTS AND DISCUSSION

It was observed that there was no interaction between the factors amounts of jitirana plus cattle manure and coriander cultivars for any of the characteristics evaluated. However, there was an isolated effect for the characteristics of plant height, number of stems, productivity, number of bunches and dry mass of coriander at a probability level of p< 0.01 (Figs. 2 to 5).

For the plant height characteristic, the amount of 4.0 kg m⁻² of jitirana plus cattle manure was what promoted the greatest increase, with a maximum value of 21.84 cm plant⁻¹ (Fig. 2). This characteristic is greatly influenced by the

availability of nitrogen in the soil, this element being responsible for leaf expansion. For the coriander cultivar characteristic, the verdão cultivar was statistically superior to the others, with values of 23.5; 21.4 and 20.2 cm plant⁻¹ for verdão, Super-verdão and tabocas respectively (Table 1).

Andrade et al. [9], studying “the yield of coriander intercropped with beetroot and arugula under green manure, found a plant height of 12.96 cm plant⁻¹ in the amount of 31.5 t ha⁻¹ at a density of 50% coriander + 50% arugula and 50% carrot, being lower than the present work”. [10], studying the agro-economic efficiency of using hairy woodrose (*Merremia aegyptia* L.) mixed with cattle manure in vegetable intercropping, found a height of 20.7 cm plant⁻¹ in the amount of 3.8 kg m⁻² of the mixture of hairy woodrose plus cattle manure, being lower than the aforementioned research.




Fig. 2. Height of coriander plant under green manure with hairy woodrose (*Merremia aegyptia* L.) plus cattle manure incorporated into the soil

For the number of stems there was an average increase of 4.59 units plant⁻¹ between the smallest quantity (0 kg m⁻²) and the largest quantity of green manure plus cattle manure (4.0 kg m⁻²) with a maximum value of 7.43 plant⁻¹ units (Fig. 3). Regarding coriander cultivars, there was no statistical difference, with average values of 7.0; 6.8; and 6.8 for verdão, super-verdão and tabocas, respectively (Table 1). The greater the number of plant⁻¹ stems, the more appreciated the coriander is by the consumer, being of great relevance for producers who sell their production.

Linhares et al. [11], studying “the agro-economic efficiency of using hairy woodrose (*Merremia aegyptia* L.) mixed with cattle manure in vegetable intercropping, found a number of

stems of 8.0 plant⁻¹ units in the amount of 4.0 kg m⁻² of the mixture of hairy woodrose plus cattle manure, being higher than the aforementioned research". [12], evaluating "quantities and decomposition times of hairy woodrose on the agronomic performance of coriander in single cultivation, found a number of stems of 10.2 plant⁻¹ in the amount of 15.6 Mg ha⁻¹, a value higher than that of the present work".




Fig. 3. Number of stems per plant under green manure with hairy woodrose (*Merremia aegyptia* L.) plus cattle manure incorporated into the soil

For coriander productivity, the data obtained were adjusted to a linear equation, with a





Fig. 4. Productivity (A) and number of sauces (B) of coriander under green manure with hairy woodrose (*Merremia aegyptia* L.) plus cattle manure incorporated into the soil




Fig. 5. Coriander dry mass under green manure with hairy woodrose (*Merremia aegyptia* L.) plus cattle manure incorporated into the soil

maximum value of 1470 g m⁻², corresponding to 29.4 units of bunches m⁻² when applying 4.0 kg m⁻² of the cilantro mixture hairy woodrose mixture as green manure plus cattle manure (Fig. 4A and 4B). For coriander cultivars, there was a statistical difference, with the verão cultivar being statistically superior to the Super-verdão and tabocas cultivars with values of 1700, 1400 and 1450 g m⁻², corresponding to 34, 28 and 29 units of bunches m⁻², respectively (Table 1).

According to [12] the number of bunches is an extremely important characteristic for coriander producers, as it determines the way it is sold, generating income for the vegetable grower. [13] evaluating coriander yield as a function of green manure incorporation of hairy woodrose (*Merremia aegyptia* L.), rooster tree (*Calotropis procera*) and kills pasture (*Senna uniflora* L.) in a semi-arid region of Brazil found coriander productivity yield of 807.83; 648.20 and 698.40 g m⁻² of bed for amounts of 1.2; 1.6 and 1.6 kg m⁻² of bed, corresponding to 16.0; 13.0 and 14.0 m⁻² units coriander sauces for hairy woodrose, rooster tree and kills pasture, respectively, lower than the aforementioned research.

Table 1. Plant height (cm plant⁻¹) and number of stems (mean terms), productivity (g m⁻²), number of bunches (units m⁻²) and dry mass (g m⁻²) of coriander depending on coriander cultivars

Coriander cultivars	Characteristics of coriander				
	Height	Stems	Productivity	Numbers of sauces	Dry mass
Verdão	23.5a	7.0a	1700a	34a	540a
Super green	21.4b	6.8a	1400b	28b	550a
Tabocas	20.2b	6.8a	1450b	29b	480a
CV (%)	10.5	12.2	10.8	10.7	11.3

* Means followed by the same letter in the column are not different from each other using the Tukey test, at a 5% probability level

In relation to dry matter, there was an increase between the highest quantity and the treatment without fertilizer with a maximum value of 103.5 g m⁻² in the amount of 4.0 kg m⁻² of the mixture of hairy woodrose and cattle manure (Fig. 5). No statistical difference was observed in relation to coriander cultivars, in terms of dry mass, with average values of 540, 550 and 480 g m⁻², for verdão, super-verdão and tabocas, respectively (Table 1). Dry matter is a characteristic that evidences plant growth [14], being influenced by organic fertilization.

4. CONCLUSION

The best agronomic performance was obtained in the amount of 4.0 kg m⁻², for the verdão cultivar, with an average value of 1585 g m⁻² and 31.7 units m⁻² for productivity and number of bunches, respectively.

The mixture of fertilizers of plant and animal origin is extremely important for increasing coriander production and productivity.

ACKNOWLEDGEMENTS

To the Jitirana research group (CNPQ-National Council for Scientific and Technological Development, Brazil), a pioneer in Brazil in the use of spontaneous species from the semi-arid region as green manure in the agroecological production of leafy and root vegetables, and the Universidade Federal Rural do Semi-Árido-UFERSA, for the physical structure and granting of scientific initiation scholarships for the development of research.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Linhares PCF, Maracajá PB, Liberalino Filho J, Assis JP, Sousa RP, Medeiros AC. Hairy woodrose (*Merremia aegyptia* L. Urban) [electronic book]: Potential use as a spontaneous species in the semi-arid region in the green manuring of vegetables In: Linhares PCF, Cunha LMM, Silva NV, Neves AM, Medeiros BBM and Paiva AC. Green and dry phytomass, levels and accumulation of macronutrients in hairy woodrose (*Merremia aegyptia* L. Urban) at different phenological stages – Nova Xavantina, MT: Ed. Pantanal. 96p. 2021;Cap. 2:24-45. Available:<https://doi.org/10.46420/9786588319901>
2. Linhares PCF, Bezerra Neto F, Maracaja PB, Duda GP, SÁ JR. Phytomass production and macronutrient contents of jitirana in different phenological stages. *Caatinga*. 2008;21(4):72-78. Available:<https://periodicos.ufersa.edu.br/caatinga/article/view/844>
3. Rêgo LGS, Martins CM, Silva EF, Silva JJA, Lima RNS. Pedogenesis and classification of soils in an experimental farm in Mossoró, Rio Grande do Norte, Brazil. *Revista Caatinga*,2016;29(4):1036-1042. Available:<https://www.scielo.br/j/rcaat/a/9M7XntrrQsNQpHsFYMLZDqy/abstract/?lang=pt>
4. Carmo Filho F and Oliveira OF. Mossoró: a municipality in the semi-arid northeast, climatic characterization and floristic aspect. Mossoró: ESAM, (Mossoroense Collection, Series B). 1995;62.
5. Linhares PCF, Maracajá PBM, Pereira FS, Assis JP and Sousa RP. Rooster tree (*Calotropis procera*) under different amounts and periods of incorporation on yield of coriander. *Revista Verde de*

- Agroecologia e Desenvolvimento Sustentável. 2014;9(3):07-12.
Available: <https://www.gvaa.com.br/revista/index.php/RVADS/article/view/2779>
6. Linhares PCF, Pereira MFS, Assis JP and Bezerra AKH. Amounts and times of decomposition of jitirana in the agronomic performance of coriander. *Ciência Rural*. 2012;42(2):243-248.
Available: <https://doi.org/10.1590/S0103-84782012000200010>
 7. Banzatto DA, Kronka SN. Experimentação agrícola. 3.ed. Jaboticabal: FUNEP. 1995;245.
 8. Barbosa JC, Malheiros EB, Banzatto D. A. ESTAT: A system for statistical analysis of agronomic trials. Jaboticabal: Unesp, Version 2.0; 1992.
 9. Andrade Filho FC, Bezerra Neto F, Moreira JN, Oliveira EQ. Yield of coriander intercropped with beetroot and arugula under green manure. VIII Brazilian Congress of Agroecology – Porto Alegre/RS – 25 to 28/11/2013.
 10. Ramalho WB, Linhares PCF, Maracajá PB, Alves LS. Agroeconomic efficiency of using jitirana (*Merremia aegyptia* L.) mixed with cattle manure in vegetable intercropping. Nova Xavantina-MT: Editora Pantanal. 2023;52.
Available: <https://doi.org/10.46420/9786581460846>
 11. Linhares PCF, Pereira MFS, Assis JP and Bezerra AKH. Amounts and times of decomposition of jitirana in the agronomic performance of coriander. *Ciência Rural*. 2012;42(2):243-248.
Available: <https://doi.org/10.1590/S0103-84782012000200010>
 12. Linhares PCF, Pereira MFS, Assis JP and Bezerra AKH. Amounts and times of decomposition of jitirana in the agronomic performance of coriander. *Ciência Rural*. 2012;42(2):243-248.
Available: <https://doi.org/10.1590/S0103-84782012000200010>
 13. Linhares PCF, Sousa JdaS, Maracajá PB, Medeiros AC, Alves LdeS, da Silva U L, Carlos KGdaS & de Souza Júnior DS. Coriander yield as a function of green manure incorporation of hairy woodrose (*Merremia aegyptia* L.), rooster tree (*Calotropis procera*) and kills pasture (*Senna uniflora* L.) in a semi-arid region of Brazil. *DELLOS: DESARROLLO LOCAL SOSTENIBLE*. 2023;16(46):2370–2385.
Available: <https://doi.org/10.55905/rdelosv16.n46-024>
 14. Taiz L, Zeiger E. Plant physiology, 3. ed. Porto Alegre: Artmed. 2017:719.

© 2023 Linhares et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/108303>