



# A Comparative Study of the Anti-Candidal Activity of Four Medicinal Plants Used in Côte d'Ivoire

Yapo Y. C. V.<sup>a</sup>, Kra A. K. M.<sup>b</sup>, Konan G. K. N. A.<sup>b\*</sup>,  
Kple T. K. M.<sup>b</sup>, Money Y. U.<sup>b</sup>, Kouame C. C.<sup>b</sup>,  
Bolou G. E. K.<sup>c</sup>, N'guessan J. D.<sup>b</sup> and Djaman A. J.<sup>b,d</sup>

<sup>a</sup> UFR des Sciences et Techniques, Université Alassane Ouattara, BPV 18 Bouaké 01, Côte d'Ivoire.

<sup>b</sup> Laboratoire Biologie et Santé, Université Félix Houphouët-Boigny, 22 BP 582 Abidjan 01, Côte d'Ivoire.

<sup>c</sup> Centre National Floristique, 22 BP 582 Abidjan 22, Côte d'Ivoire

<sup>d</sup> Institut Pasteur of Côte d'Ivoire, 01 BP 490 Abidjan 01, Côte d'Ivoire.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/IJBCRR/2023/v32i8834

### Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/108669>

**Original Research Article**

**Received: 02/09/2023**

**Accepted: 08/11/2023**

**Published: 15/11/2023**

## ABSTRACT

Some plants groups are used by traditional practitioners to treat infectious diseases. The ongoing development of antibiotic resistance continues to have a lasting impact on public health. This study aimed to assess the potential activity of the medicinal plants found in Ivorian soil. Spectrometric measurements was employed in addition to the moss test, revealed the presence of sterols and polyterpenes, polyphenolic compounds, saponins, quinones, alkaloids, and tannins in the extracts of the four plants studied. The use of the double dilution method in tilted tubes revealed the Anti-candida potential of the plant extracts. Results revealed showed that the hydroethanolic extract of

\*Corresponding author: E-mail: [angepsn@gmail.com](mailto:angepsn@gmail.com);

*T. mantaly* produced the lowest MFC (3.125 mg/mL) and IC<sub>50</sub> (0.3 mg/mL) values. It was followed by *Z. gillettii* (MFC = 3.125 mg/mL; IC<sub>50</sub> = 0.62 mg/mL) and *T. catappa* (MFC = 3.125 mg/mL; IC<sub>50</sub> = 0.68 mg/mL). No significant inhibition was observed with the extracts of *A. occidentale*. This study contributes to the valorization of medicinal plants and the floral heritage of Côte d'Ivoire.

**Keywords:** Anticandidosis activity; extraction; medicinal plants.

## 1. INTRODUCTION

The *Candida albicans* has become a significant public health concern over the past two decades [1,2,3]. The spectrum of diseases caused by this species ranges from vaginal infections, affecting up to 75% of women at least once in their lifetime, to deep-seated infections in hospitalized patients, leading to high rates of morbidity and mortality. It may also play a role in the persistence or exacerbation of certain chronic inflammatory bowel diseases.

Candidiasis is highly recurrent, affecting between 138 and 140 million women worldwide every year [4,5], with 70-75% of them experiencing at least one episode of vulvovaginal candidiasis in their lifetime [6,7]. Incidence has doubled over the past 20 years, unlike gonococcal and *Trichomonas vaginitis*, which have seen a parallel decrease [8]. *C. albicans* remains the most common species (77-95%), followed by non-albicans *Candida* (20-30%), with the most common being *C. glabrata* [9].

The prevalence of candidiasis varies from one country to another. It was 35.52% in Cameroon [10], 32.6% in Senegal [11], and 38.9% in Benin [12]. In Côte d'Ivoire, *Candida albicans* is responsible for 16.3% of otomycoses [13], 59.6% of vaginal candidiasis [14], and is found in 13.5% of HIV-positive patients suffering from chronic diarrhea. Consequently, various herbal preparations are recommended in the market to treat candidiasis.

The resistance of some commercially used antifungal agents has made it necessary to seek naturally effective alternatives. Plants have been

used in traditional phytotherapy for years. In some parts of the world, plants and herbs are still the primary sources of remedies used in disease treatment [15]. Additionally, in low- and middle-income countries, approximately 80% of people rely on medicinal plants for their primary healthcare needs [16,17]. In order to contribute to the management of candidiasis infections, this study aims to evaluate the antifungal activity of four medicinal plants species through *in vitro* growth of *C. albicans*.

## 2. MATERIALS AND METHODS

The plant material used consists of various parts of four different plants (Table 1), collected separately. These various plant parts were harvested in Côte d'Ivoire. A herbarium was prepared for each plant to identify the different plants collected at the National Center Floristics (CNF).

The *C. albicans* isolates were provided to us by the mycology laboratory of the Pharmacodynamic Biochemical Research and Teaching Unit. The choice of this clinical strain was motivated by its confirmed pathogenicity in humans and its resistance to Fluconazole, Voriconazole, and Itraconazole.

### 2.1 Plant Collection and Packaging

The organs of each plant species were harvested in large quantities and transported to the biochemistry laboratory for sorting. For each sample of the medicinal plant species, these were thoroughly washed and cleaned to remove any contaminations such as mold or parasites and infested parts of the medicinal plants.

**Table 1. The medicinal plants**

Plants	Part use	Place of harvest	Plant types
<i>Anacardium occidentale</i> L. (Anacardiaceae)	Trunk bark	Abengourou	Tree
<i>Terminalia catappa</i> L. (Combretaceae)		Agboville	Tree
<i>Terminalia mantaly</i> H. (Combretaceae)		Agboville	Tree
<i>Zanthoxylum gillettii</i> De Wild. (Rutaceae)		Abengourou	Tree

Except for the bark, the other plant components were rinsed, drained by shaking, and cut into small pieces. These plant pieces were then air-dried away from direct sunlight at room temperature (25-30°C) for three weeks. The drying process was carried out in a well-ventilated area. After drying, the plant components (for each plant species taken separately) were finely ground using an IKA-Labortechnik electric grinder (Type MFC/Janke & Kunkel). The different powders were packaged separately in non-translucent glass jars. These containers were then sealed, labeled, and stored (in the laboratory, under a hood) away from moisture and heat. The obtained powders were stored for future use.

## 2.2 Extraction Method

Aqueous and hydroethanolic extracts were prepared following the method of Zirih [18], using distilled water as a solvent and an ethanol-water mixture (70/30, v/v). To do this, one hundred grams (100 g) of plant powder were dissolved in one liter of solvent, resulting in a 1:10 (w/v) ratio. The mixture was vigorously homogenized using a blender. The homogenate was squeezed through a percale cloth and then successively filtered three times using hydrophilic cotton and filter paper. The filtrate was evaporated at 45°C using a Venticell® ventilated drying oven for 24 hours for the 70% ethanol extract and 48 to 72 hours for the aqueous extract [19].

## 2.3 Phytochemical Sorting

Phytochemical sorting is a method for characterizing the main chemical groups, such as sterols, polyterpenes, alkaloids, tannins, phenolic compounds, flavonoids, quinones and saponins. These compounds were highlighted using appropriate reagents that reacted with chemical compounds to give specific colors or precipitates that would attest to the presence or absence of the desired molecules in the extracts. This study was carried out with the total extracts (aqueous and hydroethanolic) of each plant, following the protocol of Békro [20].

## 2.4 Determination of Antifungal Activity of Extracts

### 2.4.1 Preparation of Agar

The amount of agar needed for the test was prepared according to the supplier's instructions. Forty-two grams (42 g) of powder were

homogenized in 1 L of distilled water. The mixture was stirred and heated on an IKA MAG hotplate for 5 minutes until complete homogeneity was achieved.

### 2.4.2 Incorporation of Extract into Agar

Tests were conducted separately for each extract and for each fungal species in order to determine antifungal parameter values. The comparison of these parameters would lead to the selection of the most active plant extract. The double dilution method in slanted tubes was used to incorporate the extracts into the agar. Precooked agar was poured into 10 test tubes numbered from 1 to 10, with 20 mL in tube No. 1 and 10 mL in the other tubes (No. 2 to No. 10) in each series.

Among these 10 tubes, 8 were test tubes containing the plant extract, and 2 were control tubes without plant extract: one serving as a control for germ growth (GC), and the other without germs serving as a control for culture medium sterility (SC). Overall, and according to the test series, concentrations ranged from 1000 µg/mL to 0.38 µg/mL. For the 8 test tubes in each series, concentrations varied geometrically by a factor of ½, from tube No. 1 to tube No. 8. After incorporating the extract, all 10 tubes in each series were sterilized in an autoclave at 121°C for 15 minutes and then slanted with a small base at room temperature to allow cooling and agar solidification.

### 2.4.3 Inoculum Preparation

The inoculum was prepared from young cultures of *C. albicans*, which were 48 hours old. A well-isolated colony was taken with a loop and homogenized in 10 mL of sterile distilled water. This resulted in the 10<sup>0</sup> concentrated mother suspension of 10<sup>6</sup> cells/mL. From this suspension, a 10<sup>-1</sup> suspension was prepared by diluting the first suspension 1:10 by transferring 1 mL of the suspension into 9 mL of sterile distilled water. This final suspension contained 10<sup>5</sup> cells/mL.

### 2.4.4 Tube Inoculation

For each test tube in each series, except for the SC tube, *C. albicans* culture was performed on the agar media prepared earlier by streaking 10 µL of the 10<sup>-1</sup> suspension until the entire droplet was exhausted. The spreading was done over the entire surface of the agar. These 10 µL corresponded to 1000 cells. The cultures thus

prepared were incubated in an incubator at 30°C for 72 hours. The tests were conducted under strict sterile conditions. The experiments were repeated 5 times for each plant extract [19,21].

### 2.4.5 Counting of Colonies

At the end of the incubation period, the colonies of each fungal species were counted by direct counting using a colony counter. Subsequently, the growth in the 8 experimental tubes of each series was evaluated as a percentage of survival calculated relative to 100% survival in the control growth tube. The percentage of survival of fungal germs in the experimental tubes was expressed by the given formula:

$$S = n/N \times 100$$

S: Germ survival (expressed as a percentage)  
 N: Number of colonies in growth control tube  
 n : number of colonies in experimental tube

Processing of the experimental data enabled us to plot the survival curves that helped determine the antifungal parameter values. These are:

- Minimum Fungicidal Concentration (MFC): this is the lowest concentration of extract in the tube that gives 99.99% inhibition compared with the growth control tube. Conversely, it is the concentration that gives 0.01% survival compared with the growth control tube.

- Concentration for 50% Inhibition (CI50): this is the concentration giving 50% estimated inhibition in relation to the number of colonies counted in

the growth control tube. This parameter is determined graphically from the survival curve [18].

## 3. RESULTS

### 3.1 Extraction Yields

The different extraction yields is shown in Fig. 1. Aqueous extracts gave the highest yields, with values ranging from 27.12 to 19.27%. Hydroethanol extracts generated values ranging from 12.18 to 20.14%. The best aqueous yield was obtained with *T. catappa* and the best hydroethanolic yield was obtained with *T. mantaly*.

### 3.2 Phytochemical Study

The screening allowed us to identify the main chemical groups present in the various extracts (Table 2). The plus (+) sign indicates the presence of the metabolite, while the minus (-) sign indicates its absence. Thus, we found all the targeted groups in the aqueous extracts of each plant, with the exception of saponins; the hydroethanolic extracts contained the other six secondary metabolites we were searching for.

### 3.3 Anticandidal Activity of Plant Extracts

The results reveal that the four medicinal plants have different activities. The parameter values obtained for the aqueous and hydroethanolic extracts is presented in Table 3.




Fig. 1. Total extract yield

**Table 2. The bioactive compounds of the four medicinal plant extracts**

Plants	Extracts	Chemical groups						
		Sterols and polyterpenes	Phenolic compounds	Flavonoids	Tannins	Quinones	Alkaloids	Saponosides
<i>Anacardium occidentale</i>	aqueous	+	+	+	+	+	+	+
	Ethanol 70%	+	+	+	+	+	+	-
<i>Terminalia catappa</i>	aqueux	+	+	+	+	+	+	+
	Ethanol 70%	+	+	+	+	+	+	-
<i>Terminalia mentaly</i>	aqueux	+	+	+	+	+	+	+
	Ethanol 70%	+	+	+	+	+	+	-
<i>Zanthoxylum gillettii</i>	aqueux	+	+	+	+	+	+	+
	Ethanol 70%	+	+	+	+	+	+	-

**Table 3. The Aqueous and Hydroethanolic medicinal plant extracts**

Plants		<i>Candida albicans</i>	
		Antifungal parameters (mg/mL)	
		IC50	MFC
<i>Anacardium occidentale</i>	aqueous	ND	ND
	Ethanol 70%	ND	ND
<i>Terminalia catappa</i>	aqueous	1,19	12,5
	Ethanol 70%	0,68	3,125
<i>Terminalia mentaly</i>	aqueous	0,35	6,25
	Ethanol 70%	0,3	3,125
<i>Zanthoxylum gillettii</i>	aqueous	1,1	12,5
	Ethanol 70%	0,62	3,125

#### 4. DISCUSSION

The antifungal properties of four plants used in traditional medicine were studied. Extractions were carried out using distilled water and 70% ethanol on various plant parts. The extractions in this study were performed under similar cold conditions and for a relatively equal duration. The data shows that, overall, the yields of aqueous extractions range from 27.12% to 19.57%, while those of hydroethanolic extractions range from 20.14% to 12.18%. A comparison indicate that the best yields were obtained with distilled water, which can be attributed to the polarity of distilled water. These results differ from those of Kra [21], who found that hydroalcoholic extraction yielded better results.

To determine the nature of the chemical groups present in the extracts, phytochemical screening was conducted on the plant-derived extracts. Secondary metabolites in medicinal plants form the material basis of their clinically curative effects and are important indicators to assess the quality of medicinal materials. Therefore, it is necessary to identify them in the obtained extracts.

The data showed the presence of Sterols and Polyterpenes, Phenolic Compounds, Flavonoids, Tannins, Quinones, Alkaloids, and Saponins in the aqueous extracts of all the plants. Except for Saponins, these metabolites were also found in the hydroethanolic extracts. The results obtained with *A. occidentale* are consistent with those obtained by Kikakedimau [22]. The metabolites found in *Z. gillettii* by Abiodun [23] are the same as those found in this study. Regarding the *Terminalia* species, the results of the phytochemical screening of *T. cattapa* and *T. mantaly* are identical to the data from the works of Yayé [24] and Bérenger [25], respectively. In addition to Sterols, Polyterpenes, Phenolic Compounds, Quinones, Alkaloids, and Saponins, Bérenger [25] also found other compounds such as protoanthocyanins and anthraquinones. Abiodun [23] also found leucoanthocyanins and cardiac glycosides, which were not investigated in our study.

The presence of these phytochemical substances in various plants could justify their use in the treatment of microbial infections mentioned in the literature.

Regarding antifungal activity, the analysis of the data on the evaluation of the anticandidal activity of the eight total extracts from four plants reveals

that the hydroethanolic extracts exhibited better antifungal activities than the aqueous extracts from all the plants. The antifungal test data shows that, for each plant, the hydroethanolic extracts generated lower MFC values than their aqueous counterparts during the tests. These results are similar to those of Yayé [24], Ouattara [26], Bagré [27]. According to these authors, this superior performance of the hydroethanolic extracts is attributed to their higher qualitative richness in secondary metabolites (SM).

The anticandidal activity carried out with the total extracts of *A. occidentale* did not show any confirmed activity. In fact, the colony counts in the various experimental tubes indicated a reduction in candidal load compared to the growth control tubes. Therefore, no antifungal parameters (IC50 and MFC) could be determined. This lack of clear activity could be due to the resistance profile of the *C. albicans* species used. The absence of activity in *A. occidentale* in this study does not refute its traditional use. Indeed, a study conducted by Batista [28] on several bacterial species resulted in MFC values ranging from 50 to 200 mg/mL. This study justifies the traditional use of this plant in the case of infectious diseases.

The analysis of the activities of the extracts from *Z. gillettii*, *T. cattapa*, and *T. mantaly* corroborates their traditional use. Indeed, all these extracts showed MIC values. The values obtained with the hydroethanolic extracts were lower than those generated by the aqueous extracts. Based on the MFC values, the hydroethanolic extracts of each plant were at least two times lower than their aqueous equivalents. From the comparative analysis of the extracts from different plants, it is evident that *T. mantaly* has more active extracts than those of *T. cattapa* and *Z. gillettii*. The MFC values generated by *T. cattapa* and *Z. gillettii* are identical for some extracts. When comparing IC50 values, it is evident that *Z. gillettii* is more active than *T. cattapa*.

The parameters obtained from the aqueous and hydroethanolic extracts of *Z. gillettii*, *T. cattapa*, and *T. mantaly* align with the interest of traditional medicine in the treatment of infections. These values also conforms the presence of terpenoid, phenolic, and alkaloid metabolites with antimicrobial activity.

#### 5. CONCLUSION

This study aimed to evaluate the anticandidal potential or antifungal activity of the four

medicinal plants of hydroalcoholic and aqueous extracts from four medicinal plants widely used in African pharmacopoeia. The hydroethanolic extract of *T. mantaly* showed the highest activity against the multi-resistant strain of *C. albicans* tested. The extracts of *Z. gilletii* and *T. cattapa* also exhibited inhibitory effects on *C. albicans*. A marked lack of activity was observed with *A. occidentale* against this multi-resistant yeast. The level of activity observed with these extracts suggests that medicinal plants have an inhibitory effect on *C. albicans*. However, the use of medicinal plants should be regulated, considering the results obtained with the extracts of *A. occidentale*. This study contributes to the valorization of medicinal plants and the floral heritage of Côte d'Ivoire.

### COMPETING INTERESTS

Authors have declared that no competing interests exist.

### REFERENCES

- Rai LS, Wijlick LV, Bougnoux ME, Bachellier-Bassi S, d'Enfert C. Regulators of commensal and pathogenic life-styles of an opportunistic fungus-*Candida albicans*. *Yeast*. 2021;38(4):243-250. DOI: 10.1002/yea.3550
- Poulain D. *Candida albicans*, plasticity and pathogenesis. *Crit Rev Microbiol*. 2015; 41(2):208-217. DOI: 10.3109/1040841X.2013.813904.
- Arkowitz RA & Bassilana M. Recent advances in understanding *Candida albicans* hyphal growth. *F1000Res*. 2019; 8:80-89. DOI:10.12688/f1000research.18546.1.
- Denning DW, Kneale M, Sobel JD, Rautemaa-Richardson R. Global burden of recurrent vulvovaginal candidiasis: a systematic review. *Lancet Infect Dis*. 2018;18(11):339–347. DOI :10.1016/S1473-3099(18)30103-8
- Willems HME, Ahmed SS, Liu J, Xu Z, Peters BM. Vulvovaginal Candidiasis: A Current Understanding and Burning Questions. *J Fungi*. 2020;25;6(1):27. DOI :10.3390/jof6010027
- Rosati D, Bruno M, Jaeger M, ten Oever J, Netea MG. Recurrent Vulvovaginal Candidiasis: An Immunological Perspective. *Microorganisms*. 2020-21; 8(2):144-156. DOI : 10.3390/microorganisms8020144
- Wu X, Zhang S, Li H, Shen L, Dong C, Sun Y, et al.. Biofilm Formation of *Candida albicans* Facilitates Fungal Infiltration and Persister Cell Formation in Vaginal Candidiasis. *Front Microbiol*. 2020;5;11: 1117. DOI : 10.3389/fmicb.2020.01117.
- Pizzorno JE, Murray MT, Joiner-Bey H. *The Clinician's Handbook of Natural Medicine*. Elsevier: Chronic candidiasis. 2016;206–212.
- Hashemi SE, Shokohi T, Abastabar M, Aslani N, Ghadamzadeh M, Haghani I. Species distribution and susceptibility profiles of *Candida* species isolated from vulvovaginal candidiasis, emergence of *C lusitanae*. *Curr Med Mycol*. 2019;5(4):26–34. DOI: 10.18502/cmm.5.4.2062.
- Kechia FA, Dohbit JS, Kouotou EA, Iwewe SY, Dzoyem JP, Mbopuwouo NM, et al. Epidemiological and Etiological Profile of Vulvo-Vaginal Candidiasis in Pregnant Women in Yaoundé (Cameroun) *Health Sci Dis*. 2015;16(4):12-21.
- Sylla K. Vulvo-vaginal candidiasis at the parasitology-mycology laboratory of the Centre Hospitalier Universitaire de Fann, Dakar (Senegal). *Rev Afr Malgache Rech Sci Santé*. 2020;5(2):56-68.
- Ogouyèmi-Hounto A, Adisso S, Djamaal J, Sanni R, Amangbegnon R, Biokou-Bankole B, et al. Place of vulvovaginal candidiasis in the lower genital tract infections and associated risk factors among women in Benin. *Med Trop*. 2014;78(1):49-52. DOI :10.1016/j.mycmed.2014.01.003.
- Yavo W, Kassi RR, Kiki-Barro PC, Bamba A, Kplé T, Menan EI, Ehouo F, Koné M. Prevalence and risk factors for otomycosis treated in the hospital setting in Abidjan (Ivory Coast). *Med Trop*. 2014;64(1):39-42.
- Djohan V, Angora KE, Vanga-Bosson AH, Konaté A, Kassi KF, Kiki-Barro PCM, Bedia-Tanoh AV. et al. Recurrent vulvovaginal candidiasis in Abidjan (Côte d'Ivoire): Aetiology and associated factors. *J Mycol Med*. 2019;29(2):127-131.
- Therizol-Ferly PM, Tagliante-Saracino J, Kone M, Konan A, Ouhon J, Assoumou A, Aka K, Assale G. Chronic diarrhea and parasitoses in adults suspected of AIDS in the Ivory Coast. *Bull Soc Pathol Exot Filiales*. 2016;82(5):690-693.
- Bishop FL, Lewith GT. Who uses CAM? A narrative review of demographic

- characteristics and health factors associated with CAM use. Evidence-based complementary and alternative medicine. 2010;7(1):11-28.
17. Pathak K., Das RJ. Phytotherapy-a rational approach to the healthcare system. International Journal of Herbal Medicine. 2013;1 (3):86-89.
  18. Zirihi GN, Datté JY, Kra-Adou KM, Grellier P. Phytochemical and pharmacological studies of the alcoholic extract (MFA) of *Fagara macrophylla* Oliv. Engl. (Rutaceae): the chemical structure of the active compound inducing antipaludic activity. Journal of Chinese Clinical Medicine. 2003;2:205-210.
  19. Yomeh CVY, N'Dri GK, Kiyinlma C, Djeneb C, Guédé NZ. Botanical study, evaluation of antifungal activity on *in vitro* growth of *C. albicans* and toxicity on HFF cells of *mallotus opposifolius*. Journal of Animal & Plant Sciences. 2016;28(1):4330-4339.
  20. Békro YA, Békro JAM, Boua BB, Tra BFH, Ehilé EE. Ethnobotanical study and phytochemical screening of *Caesalpinia benthamiana* (Baill.) Herend.et Zarucchi (Caesalpinaceae). National Science Review. 2007;4 (2): 217-225.
  21. Kra AKM, Ahon GM, Djo-Bi D, Ouattara S, Coulibaly A, Djaman AJ. Antifungal activities of medicinal plants extracts of Ivorian pharmacopoeia. Journal of Intercult Ethnopharmacology. 2014;3:159-166.
  22. Kikakedimau Nakweti R, Muganza MD, Nsendo Muangu Kabi O, Doumas P, Kahambu Mbugheki S, Taba KM, Luyindula NS. Correlation between antioxidants and antiradical activities with *in vitro* antimalarial activity of *Phyllanthus odontadenius*. Société américaine de spectrométrie de masse. 2019;3(7):144–154.
  23. Abiodun EA, Oluwabusola IO, Abimbola O. Phytochemical screening and *in vitro* antibacterial activity of *Justicia secunda* Vahl leaf extracts on selected clinical pathogens. Micro Medicine. 2020;8:46-54.
  24. Yayé YG, Kra AKM, Ackah AAB, Djaman AJ. Evaluation of the antifungal activity and purification tests of the active principles of extracts of *Terminalia mantaly* H. Perrier, a combretaceae on the *in vitro* growth of *Candida albicans*. Bulletin de la Société Royale des Sciences de Liège. 2011;80: 953-964.
  25. Bérenger L, Salomé K, Hyacinthe AJG, Bénédicte KK, Eric M, Marie FH, Joanne B, Yvan L, Joëlle QL, Mansourou M. Chemical composition and nutritional properties of *Terminalia catappa* L. oil and kernels from Benin. Comptes Rendus Chimie. 2019;19(7):876-883.
  26. Ouattara S, Kporou KE, Kra AKM, Zirihi GN, N'Guessan JD, Coulibaly A, Djaman AJ. Antifungal activities of *Terminalia ivorensis* A. Chev. bark extracts against *Candida albicans* and *Aspergillus fumigatus*. Journal of Intercultural Ethnopharmacology. 2013;2:49-52.
  27. Bahi CI, Ouattara K, Guede NZ, Djaman AJ, Coulibaly A, N'Guessan JD. Botanical study and exploration of the antifungal activity of *Morinda morindoides* (Baker) Milne-Redh on the *in vitro* growth of *Cryptococcus neoformans*. Phytothérapie, 2011;9:136-141.
  28. Baptista A, Gonçalves RV, Bressan J, Pelúzio MDCG. Antioxidant and Antimicrobial Activities of Crude Extracts and Fractions of Cashew (*Anacardium occidentale* L.), Cajui (*Anacardium microcarpum*), and Pequi (*Caryocar brasiliense* C.): A Systematic Review. Oxid Med Cell Longev; 2018. DOI:10.1155/2018/3753562.

© 2023 Yapo et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:  
The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/108669>