



Effect of Irrigation Scheduling and Different Sowing Dates on Growth and Yield of Wheat (*Triticum aestivum* L.)

Pravesh Kumar ^{a*}, Vineet Dheer ^b, Pradeep Kumar ^b,
Jaykar Singh ^b, Krishna Kumar Singh ^c,
Yogesh Kumar ^d and Anil Kumar Singh ^{a++}

^a Department of Agronomy, Acharya Narendra Deva University of Agriculture and Technology, Ayodhya, India.

^b Department of Agronomy, Chandra Shekhar Azad University of Agriculture and Technology, Kanpur, India.

^c Department of Soil Science and Agricultural Chemistry, Acharya Narendra Deva University of Agriculture and Technology, Kanpur, India.

^d Department of Soil Conservation and Water Management, Chandra Shekhar Azad University of Agriculture and Technology, Kanpur, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2024/v14i13820

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/110922>

Original Research Article

Received: 29/10/2023

Accepted: 05/01/2024

Published: 07/01/2024

ABSTRACT

The study was conducted at Acharya Narendra Deva University of Agriculture and Technology, Ayodhya (U.P.) Agronomy Research Farm in rabi season 2021-22. Twelve main plot treatments included 15th November, 25th November, and 5th December sowing dates, while four sub plot treatments included irrigation at 0.6, 0.8, 1.0, and 1.2 IW/CPE ratios. Split plot design was used for

⁺⁺ Assistant Professor;

^{*}Corresponding author: E-mail: praveshkumar18072000@gmail.com;

three replications. Under 15th November sowing, all growth, yield, and characteristics rose dramatically. Irrigating at 1.0 IW/CPE ratio increased wheat shoot m-2, plant height (cm), dry matter accumulation (g m-2), yield characteristics, grain and straw yield (q ha-1) considerably. Wheat yields were highest when sown on November 15. Under 15th November planting, water use efficiency was highest (9.85 kg ha-1mm-1). Out of all treatments, 15th Nov planting had the highest biological (128.21 q ha-1) and seed yield (51.43 q ha-1) and irrigation on 1.0 IW/CPE had the highest biological (127.18 q ha-1) and seed yield (50.83 Maximum Harvest Index was 40.11 percent with 15th November planting and 1.0 IW/CPE ratio irrigation (39.96%).

Keywords: *Wheat; irrigation scheduling; moisture regimes; IW; CPE and leaf area index.*

1. INTRODUCTION

One of the most important cereal crops, wheat (*Triticum aestivum* L.), is a member of the Poaceae family. Two billion people, or 36 percent of the global population, follow this diet as their primary way of eating. Wheat is the source of 20% of the caloric intake for 55% of the world's population. Wheat, sometimes known as the "King of cereals" has the ability to self-pollinate. The country's second most important source of grain after rice. India comes in at number two in terms of wheat output. Wheat is the cereal that is grown around the globe the more than any other type. In 1967 and 1968, dwarf Mexican wheat completely altered the production of wheat in India [1,2].

Wheat grows on all continents. China, India, Russia, Ukraine, the US, France, Canada, Germany, Pakistan, and Australia produce the most wheat. Wheat is the second most produced crop after maize with 219 million ha, 760 million metric tonnes, and 3390 kg ha⁻¹ productivity. India grows 29.8 million hectares of it, generating 109 million tonnes at 3424 kg ha⁻¹. Anonymous, [3]. Uttar Pradesh, Punjab, Haryana, Madhya Pradesh, Rajasthan, Bihar, and Maharashtra grow the most wheat by area and output. Uttar Pradesh leads in acreage (9.21 million ha) and output (24.51 million tonnes), but its productivity (2.7 tonnes ha⁻¹) is much lower than Punjab and Haryana. Anonymous, [3].

More than a billion people throughout the world consume wheat in various forms. It is the second most significant crop grown as a staple meal in India, behind rice. Wheat is consumed as "chapatis" in regions where it is the main source of cereal food. Wheat is consumed in the form of "puris" or "upma" in regions where rice is the main grain crop. Additionally, wheat is used in a variety of different dishes like "Dalia," "halwa," etc. The consumption of baked leavened bread, flakes, cakes, biscuits, etc. is rising quickly

throughout the majority of the nation's metropolitan centers.

India has ample land and ideal weather for agriculture cultivation. Thus, wheat output ranks second globally. Many factors contribute to this country's poor wheat production. Environmental variables like late planting reduce wheat output. Another issue is the unavailability of improved varieties with quick maturity and appropriate for late sowing due to the crop's shorter growth cycle. Late-sown cultivars vary in yield and nutrient absorption.

According to more recent and thorough studies on the consequences of climate change, temperatures in South Asia are expected to rise by up to 3–4°C by the end of the century (DEFRA, 2005) and rainfall is expected to increase significantly. Reduced grain yield is one of the predicted implications on wheat output, with the worst effects occurring in low-potential locations like the eastern plains.

Physiological growth stage, climatic (IW/CPE ratio), and soil moisture depletion are the three fundamental techniques for scheduling wheat irrigation. The climatological technique is scientific and practical, and scientists and researchers worldwide acknowledge it. It is generally established that complete crop cover evapotranspiration is linked to open pan evaporation. Crop irrigation schedule is based on the ratio of fixed irrigation water (IW) to CPE. Soil measurements and crop monitoring inform irrigation schedule. Irrigation scheduling involves choosing the time and amount of water. Knowing the plant's initial soil water allows intelligent scheduling. This allows determining the earliest date for subsequent irrigation for optimal irrigation using the system before water stress impairs crop performance. Improved irrigation timing lowers expenses and boosts crop quality. The scientific and beneficial climatological irrigation scheduling technique is generally

acknowledged by scientists and researchers. This IW/CPE method is worth considering for its simplicity and great water efficiency.

2. MATERIALS AND METHODS

The current experiment contained twelve treatment combinations: main plot (15 November, 25 November, and 5 December) and sub plot (0.6, 0.8, 1.0, and 1.2 IW/CPE ratio irrigation). Three replications utilised split plots. Acharya In Rabi 2021-22, Narendra Deva University of Agriculture & Technology, Ayodhya, sponsored the experiment at its Agronomy Research Farm in Wheat cv. PBW 343 with 20 cm row spacing in 5.0m x 6.0m plots. The experimental site is at 26°47' N latitude and 82°12' E longitude, 113 metres above sea level. It is part of the Indo-Gangetic alluvial plains (IGP), which have a semi-arid subtropical climate and alluvial calcareous soil. About 1002 mm of precipitation fall on average each year, 80 to 85 percent of which fall between June and September during the monsoon season. 49.6 mm of rain fell altogether during the trial. The winter months are chilly, and there may be some frost at this time. The experimental soil was sandy loam with a pH of 8.1, EC of 0.32 dSm⁻¹, organic carbon content of 0.34%, nitrogen availability of 156.10 kg ha⁻¹, P₂O₅ availability of 15.13 kg ha⁻¹, and potassium availability of 280.42 kg ha⁻¹. The test location was primed for germination. The field was fertilised with 120 kg N, 60 kg P₂O₅, and 40 kg K₂O ha⁻¹ before planting [4]. Quality PBW 343 seed was seeded at 100 kg ha⁻¹ on November 15, 25, and December 5, 2021. The best pre-emergence herbicide application method was a Knapsack sprayer with a flat fan nozzle and 500 litres of water per hectare. To grow a good harvest, several agronomical and plant protection measures were employed [5].

The IW/CPE ratio are calculated by the formula of

$$\frac{IW}{CPE} = \frac{\text{Irrigation water depth (mm)}}{\text{Cumulative pan evaporation (mm)}}$$

Each plot has five randomly selected plants tagged. Height (cm), tillers, and leaves were measured at 30, 60, 90 DAS and maturity mean values were calculated. Yield traits like from planting to 50% blooming, wheat plant productive tillers per m², spike length, spikelet count, grain

count, and test weight of tagged plants from each plot replication wise were recorded.

3. RESULTS AND DISCUSSION

3.1 Initial Plant Population (m⁻²)

Dates of sowing and moisture regimes statistically affect plant population (m⁻²). Planting dates and moisture regimes did not have affect initial plant population (m⁻²). However, wheat sown on November 15th had the most plants (157.75), followed by November 25th and December 5th. The moisture regime with the highest plant population was 1.0 IW/CPE ratio (159.33), followed by 0.8 IW/CPE ratio (156.87), 1.2 IW/CPE ratio (154.4), and 0.6 IW/CPE ratio (154.2).

3.2 Plant Height (cm)

Beginning on November 15th, when the wheat crop is sown, plant height rises at every stage of crop growth. The date of seeding had no effect on the plant's height at 30 DAS, according to the data. On November 15th, the wheat plant reached its maximum height of 83.98 cm, which corresponded to November 25 at 60 and 90 DAS. The plant height with the lowest value was observed when the planting date was set for December 5. The moisture regime did not appear to have a substantial impact on plant height at 30 DAS. When it came to plant height, the response to irrigation was better at 1.0 IW/CPE ratio, similar at 0.8 IW/CPE ratio, and much better at 60 and 90 DAS [6,7].

3.3 Number of Tillers (m⁻²)

Tillers rose gradually up to 60 DAS, when they began to decline in spite of treatments. At the 30-day mark, there was no appreciable difference in the influence of different sowing dates on the growth of wheat crops. However, at the 60, 90, and harvest stages, the differences in sowing dates were significant and equivalent to the sowing date of November 25. Variable moisture regimes had significant impacts on wheat crop growth stages at 30, 60, and 90 days later as well as at harvest; these effects were equivalent to 0.8 and 1.2 IW/CPE ratios. It might be accounted for by the fact that better planting dates and increased moisture availability in well-managed plots produced more shoots (m⁻²) than in other treatments [7].

Table 1. Effect of dates of sowing and moisture regimes on growth of wheat at different growth stage

Treatments	Initial Plant population m ⁻²	Plant height (cm)			No. of tillers m ⁻²			Leaf area index			Dry matter accumulation (g m ⁻²)				
		30 DAS	60 DAS	90 DAS	30 DAS	60 DAS	90 DAS	At harvest	30 DAS	60 DAS	90 DAS	30 DAS	60 DAS	90 DAS	At harvest
Date of sowing															
D ₁ : 15 November	157.75	24.45	73.03	83.98	189.30	349.53	356.21	359.81	1.75	4.90	5.04	94.65	630.81	1051.35	1282.13
D ₂ : 25 November	156.20	23.98	69.38	79.78	187.44	326.15	332.38	335.74	1.71	4.65	4.79	93.72	595.13	991.89	1209.62
D ₃ : 05 December	154.65	23.48	63.55	73.08	185.58	312.09	318.06	321.27	1.68	4.27	4.39	92.79	551.74	920.19	1121.66
SEm±	3.93	0.60	1.67	1.92	4.72	8.12	8.27	8.35	0.043	0.112	0.116	2.36	14.34	19.86	30.17
CD at 5%	NS	NS	6.57	7.56	NS	31.88	32.49	32.82	NS	0.441	0.454	NS	56.31	77.98	118.47
Moisture regime															
I ₁ : 0.6 IW/CPE ratio	154.20	23.37	64.33	73.98	185.04	314.33	320.34	323.58	1.67	4.31	4.45	92.52	562.46	937.43	1143.21
I ₂ : 0.8 IW/CPE ratio	156.87	24.30	71.33	82.03	188.24	335.10	341.51	344.96	1.74	4.78	4.92	94.12	610.36	1017.26	1227.64
I ₃ : 1.0 IW/CPE ratio	159.33	24.70	73.13	84.10	191.20	345.98	352.60	356.16	1.76	4.90	5.05	95.60	619.23	1042.91	1271.84
I ₄ : 1.2 IW/CPE ratio	154.40	23.50	65.80	75.67	185.28	321.61	327.75	331.07	1.68	4.42	4.55	92.64	578.20	953.63	1175.19
SEm±	4.16	0.64	1.86	2.14	5.01	8.87	9.04	9.13	0.046	0.125	0.129	2.50	14.69	25.85	30.00
CD at 5%	NS	NS	5.43	6.26	NS	25.90	26.40	26.67	NS	0.366	0.377	NS	42.89	75.45	87.58

3.4 Leaf Area Index

Except for the 30 DAS growth stage, sowing and watering regimes affected wheat leaf area index. The leaf area index decreased after 30 days but grew and peaked after 60 and 90 days of crop development. A small study found that 30 DAS sowing dates did not affect leaf area index. The greatest leaf area index was 4.90 at 60 DAS and 5.04 at 90 DAS on November 15. November 25 and December 5 readings were comparable. Different moisture regimes did not significantly impact the leaf area index at 30 DAS, but at 60 and 90 days into wheat crop growth, they did, reaching 1.0 IW/CPE ratio. The greatest leaf area index (5.05) at 90 DAS was achieved with moisture regime 1.0 IW/CPE ratio, equal to irrigation at 0.8 at 60 and 90 DAS [7].

3.5 Dry Matter Accumulation (g m^{-2})

The dry matter output of the wheat crop rose with growth, in line with growth parameters. Wheat crops had the greatest dry matter content during harvest. Dry matter rose more quickly between 30 and 60 DAS, then between 60 and 90 DAS, and finally from 90 until harvest. Table 1 scans reveal that on November 15, November 25, and December 5, respectively, there was a rise in dry matter accumulation in the latter phases (60, 90 DAS, and harvest). At 60, 90 DAS, and harvest stage, the IW/CPE ratio under irrigation was 1.0, which was noticeably better than 0.8, 1.2, and 0.6. When it came to dry matter production at harvest stage, irrigation at 1.0 IW/CPE ratio generated the most (1271.84 g m^{-2}), whereas

irrigation at 0.6 produced the least (1143.21 g m^{-2}) [8].

3.6 Biological Yield (q ha^{-1})

Irrigation at 1.0 IW/CPE ratio (I3) produced the maximum biological production (127.18 q ha^{-1}), which was followed by 0.8 (I2), 1.2 (I4), and 0.6 (I1). In comparison to 25 November and 5 December, which produced biological yields of 120.962 and $112.583 \text{ q ha}^{-1}$, respectively, the 15 November sowing produced the greatest yield ($128.213 \text{ q ha}^{-1}$) [9,10].

3.7 Grain Yield (q ha^{-1})

Statistics on average grain production are significantly impacted by planting dates and moisture regimes. The data indicates that grain yield is significantly impacted by moisture regime. The highest grain yield (51.43 q ha^{-1}) was recorded on November 15, and it was subsequently reached on November 25, and December 5. Irrigation at 1.0 IW/CPE ratio was the moisture regime with the maximum grain production (50.83 q ha^{-1}), which was considerably superior than I2 (0.8), I4 (1.2), and I1 (0.6) [9,10].

3.8 Straw Yield (q ha^{-1})

The Table 2. demonstrates how different moisture regimes greatly increased straw production. The highest straw output was recorded on November 15, 25, and December 5 at 76.79 q ha^{-1} . The straw yield with I3 (1.0 IW/CPE ratio) was the greatest, at 76.35 q ha^{-1} [9,10].

Table 2. Effect of dates of sowing and moisture regimes on yield of wheat

Treatments	Grain yield (q ha^{-1})	Straw yield (q ha^{-1})	Biological yield (q ha^{-1})	Harvest index (%)
Dates of sowing				
D1: 15 November	51.43	76.79	128.21	40.11
D2: 25 November	48.28	72.69	120.96	39.91
D3: 05 December	44.93	67.66	112.58	39.71
SEm\pm	1.19	1.65	1.03	1.01
CD at 5%	4.68	6.47	2.94	NS
Moisture regime				
I1: 0.6 IW/CPE ratio	45.57	68.75	114.32	39.85
I2: 0.8 IW/CPE ratio	49.53	74.52	124.05	39.92
I3: 1.0 IW/CPE ratio	50.83	76.35	127.18	39.96
I4: 1.2 IW/CPE ratio	46.90	69.88	116.78	39.90
SEm\pm	1.20	1.81	2.08	1.06
CD at 5%	3.51	5.28	4.41	NS

3.9 Harvest Index (%)

Harvest index was highest on November 15, sowing (40.11%), and lowest on December 5, sowing (39.71%). The harvest index for moisture regimes ranged from the lowest (40.05%) at 0.6 to the greatest (40.16%) at 1.0 IW/CPE ratio [6].

4. CONCLUSIONS

Based on the overall findings, it can be said that November 15th was the best day to sow wheat for growth and production. Wheat responded best to irrigation at a 1.0 IW/CPE ratio (6 irrigations), according to research on moisture regimes. No correlation was seen between the dates of sowing and the moisture regime.

CONFERENCE DISCLAIMER

Some part of this manuscript was previously presented in the conference: "International Conference on Emerging Trends in Agriculture & Allied Sector for Sustainable Developments" organized by Faculty of Agricultural Sciences & Allied Industries, Rama University, Kanpur Nagar, U.P., India on 8th and 9th December, 2023. Web Link of the proceeding: <https://www.ramauniversity.ac.in/news-rama-university-hosts-successful-international-conference-on-emerging-trends-in-agriculture-12-49-5706>.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Pal S, Malik N, Singh S, Pandey V, Yadav R, Dheer V, Sachan DS, Gangwar P. Effect of foliar feeding of nutrients, salicylic acid and sea weed extract on yield and yield attributes of wheat varieties under Tarai region. *International Journal of Plant & Soil Science*. 2023;35(20):574–581. Available:<https://doi.org/10.9734/ijpss/2023/v35i203841>
2. Ahmad A, Kumar R, Effect of irrigation scheduling on the growth and yield of wheat genotypes. *Agric. Sci. Digest*. 2015; (3):199-202.
3. Anonymous. Agriculture statistics at a glance, Directorate of Economics and Statistics, Department of Agriculture and Co-operation, Ministry of Agriculture, Government of India, New Delhi; 2020.
4. Yadav AK, Dheer V, Singh J, Yadav A, Singh KK, Kumar P, Singh V. Effect of FYM, Vermi-compost, azotobacter inoculation and chemical fertilizers on growth and yield in wheat (*Triticum aestivum* L.). *International Journal of Environment and Climate Change*. 2023; 13(12):1312–1316. Available:<https://doi.org/10.9734/ijecc/2023/v13i123797>
5. Kumar P, Singh J, Kumar P, Dheer V, Yadav R, Singh KK, Singh R, Singh AK. Effect of irrigation scheduling and different sowing dates on water productivity and economics of wheat (*Triticum aestivum* L.). *International Journal of Environment and Climate Change*. 2023;13(12):1292–1297. Available:<https://doi.org/10.9734/ijecc/2023/v13i123794>
6. Aslam H, Ansari MA, Baloch SK, Baloch SU, and Baloch AS. Effect of irrigation scheduling on the growth and harvest index of wheat (*Triticum aestivum* L.) varieties. *Persian Gulf Crop Protection*. 2014;3(2):15-29.
7. Ngwako S, Mashiqa PK. The effect of irrigation on the growth and yield of winter wheat (*Triticum aestivum* L.) cultivars. *Int. J. of Agric. and Crop Sci*. 2013;5(9):976-982.
8. Kumar S, Dhindwal AS, Arya RK. Dry matter and straw yield in wheat as influenced by preceding crops, planting methods and irrigation levels. *Forage Research*. 2013;39(2):88-92.
9. Dhaka AK., Bangarwa AS, Pannu RK, Malik RK. Garg R. Phenological development, yield and yield attributes of different wheat genotypes as influenced by sowing time and irrigation. *Agricultural Science Digest*. 2006;26(3):174-177.

10. Throat TN, Bhan M, Agrawal KK. Effect of sowing date on weed infestation and yield of wheat (*Triticum aestivum* L.) varieties under different irrigation schedules. *Jawaharlal Nehru Krishi Vishwa Vidyalaya Research Journal*. 2015;49(1):41-45.

© 2024 Kumar et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/110922>