



Effect of Vitamin D₃ Enrichment by Ultra Violet Blue (UVB) Radiation and Dietary Zinc Supplementation on Egg Production and Feed Consumption in Layer Birds

P. S. Megha^{a++*}, V. Ramnath^{b#} and K. V. Vishnudas^{c++}

^a Department of Veterinary Physiology, Poyyara House, P. O. Naduvilkara, Vatanappally, Thrissur, Kerala-680 614, India.

^b Department of Veterinary Physiology, College of Veterinary and Animal Sciences, Mannuthy, Thrissur, Kerala-680651, India.

^c Department of Animal Science, College of Agriculture and Life Sciences, Texas A&M University, College Station, Texas, Pin-77840, USA.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/ACRI/2024/v24i4665

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/114856>

Original Research Article

Received: 16/01/2024

Accepted: 19/03/2024

Published: 22/03/2024

ABSTRACT

Nutritionists have recently been involved in enriching or altering the nutritional profile of poultry products, such as meat and eggs, in relation to increased consumer interest in the nutritional value of foods. Newer technologies like development of designer eggs were exploited beyond their traditional food value. Designer eggs are those that have modified contents from standard egg

⁺⁺ Ph. D Scholar;

[#] Professor and Head;

*Corresponding author: Email: meghaps4748@gmail.com, meghaps7@gmail.com;

according to the consumers' preference or according to current market demand. The present study to evaluate the influence of vitamin D₃ enrichment by UVB radiation and dietary zinc supplementation on egg production and feed consumption in layer birds was conducted at Department of Veterinary Physiology, College of Veterinary and Animal Sciences, Mannuthy, for a period of 12 weeks from 01 February 2020 to 01 May 2020. A total of sixty-four laying hens, belonging to crossbred (White Leghorn N strain and Desi) housed in animal house attached to the department were used in the experiment. The treatments of this study included a control diet (according to BIS 2007) fed group (group I), a group with zinc supplementation alone (group II), a group with UVB exposure without zinc supplementation (group III) and a group with both UVB exposure and zinc supplementation (group IV). The four treatments were tested in 28-week-old crossbred hens in a completely randomised experimental design. Each of the experimental treatments had four replicates with four birds with all standard managemental conditions. The statistical analysis showed that egg production was not significantly affected by UVB light exposure and dietary zinc supplementation. However, the effect of UVB exposure and zinc supplementation via feed was highly significant ($p \leq 0.01$) for feed consumption. The highest overall mean feed intake (113.18 ± 0.14 g/day) was recorded with those birds exposed to UVB and fed with inorganic zinc. While the lowest value (107.17 ± 0.24 g/day) was recorded in group of birds fed with zinc supplementation in diet without UVB exposure. As a future prospective we can study the egg production rate and feed consumption rate in layer chicken that are intended for different types of designer egg production.

Keywords: Egg production; Feed consumption; *ultra violet blue*; vitamin D3 enrichment; zinc enrichment; designer egg production.

1. INTRODUCTION

Eggs, irrespective of table or fertilized, are considered as natural life-supporting chemical storehouse which provides highly nutritious food with biological packaging (eggshell) without any chance of adulteration [1]. Table eggs of chicken are a vital part of the diet and have been used as food by human beings since ancient times. Poultry eggs are regarded as inexpensive, convenient and low-calorie sources of high-quality protein with several other essential nutrients. Over the last four decades, poultry production in India has made impressive progress [2].

The Indian industry is presently focused on the production of eggs that have higher or enriched levels of certain nutrients. These eggs are capable of safeguarding the health of consumers. Designer eggs have high market demand because of the consumers' willingness to purchase designer eggs due to their nutritional qualities in addition to regular eggs [3]. Singh et al. [4] reviewed the possibilities of nutrification in eggs by nutritional manipulation of laying hens' ration which would be safe for human consumption without compromising the quality of products. Considering these facts, the current study was planned to produce vitamin D₃ enriched eggs by UVB exposure and to produce zinc-enriched eggs by supplementing laying

hens' diet with required levels of zinc. The dual enrichment with these two provides healthcare benefits over other food sources. Simultaneously we have to evaluate how the egg production parameters and feed consumption rate in layer chicken are affected by this enrichment

2. MATERIALS AND METHODS

The study was conducted using sixty-four numbers of 28-week-old crossbred (White Leghorn N strain X Desi) layer birds procured from All India Coordinated Research Projects (AICRP) on Poultry for Eggs, Mannuthy, KVASU. The birds were randomly distributed in a completely randomized experimental design and they were placed into four treatment groups, each with four replicates having four birds in each replicate (Table 1).

All birds were fed with a standard layer diet and raised under standard managemental conditions up to 28 weeks of age at AICRP on Poultry for Eggs Mannuthy, KVASU. The birds were immunized against diseases as per standard protocol followed on the farm. On completion of 28 weeks of age, the birds (average body weight of 1.28 ± 0.06 kg) were brought and housed in well-ventilated cages at $24.5 \pm 0.5^\circ\text{C}$ ambient temperature and at relative humidity ranging from 60-80 per cent [5]. A photoperiod of 16 hours per day was ensured throughout the period.

Table 1. Experimental layout

Treatment Groups	Treatment	No of birds
I	Control with standard layer diet (BIS 2007)	4X4
II	Zn enriched diet	4X4
III	UVB light exposed	4X4
IV	Zn enriched diet + UVB light exposed	4X4

Table 2. Percent ingredient composition of experimental diet

Sl. No.	Feed ingredients	Percentage
1	Yellow maize	56.50
2	De-oiled rice bran	6.50
3	Soya bean meal	27.50
4	Calcite powder	7.50
5	Dicalcium phosphate	1.50
6	Salt	0.50
Total		100.00
Feed Supplements (g/100 kg feed)		
1	L-Lysine	100
2	DL-Methionine	100
3	Vitamin premix	50
4	Toxin binder	100
5	Choline chloride	100
6	Trace mineral mixture	100
7	Liver tonic powder	25
Total		575

Birds of treatment groups III and IV were exposed to UVB (280-315 nm) light (M/s Philips India Ltd., Hyderabad) for 3h daily @ 8.00-8.30 a.m., 11.00 a.m.-12.00 p.m., 2.00-3.00 p.m. and 4.30-5.00 p.m. The UVB radiation dosage at a distance of 20 cm was 76 µW/cm² as claimed by the manufacturer. A 60cm long, 240 V, 50 Hz and 36 W UVB tube equipped with a heat protection reflector was placed in the lower front part of the cages to ensure optimal UVB light exposure on the featherless skin of feet and legs of birds, especially during the feeding time [6]. The zinc level in the control diet was formulated according to BIS 2007 standards (Table 2). Treatment

groups II and IV were provided with a diet incorporating inorganic zinc sulphate at the level of 75 mg/kg mash diet [7].

2.1 Egg Production

Daily egg production was recorded as follows and expressed in hen-housed and hen-day egg production per cent.

Hen Housed egg production = (Average daily number of eggs produced / Total number of birds housed at the beginning) x100

Hen day egg production = (Number of eggs produced on that day/ Total number of live hens on that day) x100

2.2 Feed Consumption

Feed intake was recorded in each replicate at the end of every 28-day period as the difference between the total feed offered and the cumulative feed left over. From the above data feed intake per bird/day was calculated.

3. RESULTS AND DISCUSSION

3.1 Hen-housed Egg Production

The results of the influence of UVB light treatment and zinc feeding on hen-housed egg production are presented in Table 3 (a) and (b).

Table 3 (a). Mean ±S.E. hen housed egg number in four treatment groups during 29-40 weeks of age

Treatment Groups	Period/age in weeks			Overall mean 29-40
	Period I 29-32	Period II 33-36	Period III 37-40	
I	21.13±0.16	21.88±0.16	19.63±0.13	20.88±0.12
II	20.75±0.10	21.75±0.10	19.69±0.12	20.73±0.09
III	21.00±0.10	22.13±0.07	19.56±0.06	20.90±0.06
IV	20.63±0.16	22.13±0.07	20.00±0.10	20.92±0.08
p-value	0.082	0.073	0.053	0.476

Table 3 (b). Mean \pm S.E. hen housed egg percent in four treatment groups during 29-40 weeks of age

Treatment Groups	Period/age in weeks			
	Period I 29-32	Period II 33-36	Period III 37-40	Overall mean 29-40
I	75.45 \pm 0.58	78.13 \pm 0.58	70.09 \pm 0.45	74.55 \pm 0.45
II	74.11 \pm 0.36	77.68 \pm 0.36	70.31 \pm 0.43	74.03 \pm 0.31
III	75.00 \pm 0.36	79.02 \pm 0.26	69.87 \pm 0.22	74.63 \pm 0.22
IV	73.66 \pm 0.58	79.02 \pm 0.26	71.43 \pm 0.36	74.70 \pm 0.27
p-value	0.082	0.073	0.053	0.476

The effect of influence of UVB light treatment and zinc feeding was not significant for mean hen-housed egg production. During the entire period, egg production did not vary between treatment groups. Egg production parameters showed no significant changes due to UVB treatment and zinc feeding. The overall mean percent hen housed egg production was also not significantly different among treatment groups.

3.2 Hen Day Egg Production

Data presented in Table 4 (a) and (b) illustrate the effect of UVB light treatment and zinc feeding on mean hen day egg production.

The effect of UVB treatment and zinc supplementation was not significant for the mean hen-day egg production during the experimental period. In the present study, the UVB light

treatment and zinc supplementation did not cause any significant change in both hen housed egg number as well as hen day egg number over the entire experimental period of 29 to 40 weeks of age, as shown in Table 3 (a) and 4 (a), respectively. Similarly, the mean per cent hen housed and hen day egg production also was not significantly affected by the treatments employed in the study [Table 3(b) and 4(b)].

Similar to the present findings, Supplee et al. [8] found no change with respect to egg production when zinc was supplemented in the feed for layers maintained on the floor or cages. Stahl et al. [9] recorded that 28 ppm of zinc in the control diet was appropriate for egg production and at no time layers consuming this basal diet have lower egg production than Zn supplemented group of birds. Hence 28 ppm of zinc which was naturally supplied was sufficient for egg production, hatchability and growth of progeny.

Table 4 (a). Period-wise mean \pm S.E. hen day egg number in four treatment groups during 29-40 weeks of age

Treatment Groups	Period/age in weeks			
	Period I 29-32	Period II 33-36	Period III 37-40	Overall mean 29-40
I	21.13 \pm 0.16	21.88 \pm 0.16	19.63 \pm 0.13	20.88 \pm 0.12
II	20.75 \pm 0.10	21.75 \pm 0.10	20.40 \pm 0.82	20.96 \pm 0.30
III	21.00 \pm 0.10	22.13 \pm 0.07	19.56 \pm 0.06	20.90 \pm 0.06
IV	20.63 \pm 0.16	22.13 \pm 0.07	20.00 \pm 0.10	20.92 \pm 0.08
p-value	0.082	0.073	0.490	0.989

Table 4 (b). Period-wise mean \pm S.E. hen day egg percent in four treatment groups during 29-40 weeks of age

Treatment groups	Period/age in weeks			
	Period I 29-32	Period II 33-36	Period III 37-40	Overall mean 29-40
I	75.45 \pm 0.58	78.13 \pm 0.58	70.09 \pm 0.45	74.55 \pm 0.45
II	74.11 \pm 0.36	77.68 \pm 0.36	72.86 \pm 2.93	74.84 \pm 1.06
III	75.00 \pm 0.36	79.02 \pm 0.26	69.87 \pm 0.22	74.63 \pm 0.22
IV	73.66 \pm 0.58	79.02 \pm 0.26	71.43 \pm 0.36	74.70 \pm 0.27
p-value	0.082	0.073	0.490	0.988

However, Durmus et al. [10] recorded a graded response on egg production, when the level of inclusion of zinc increased two times (1.21% more eggs), three times (1.64% more eggs) and 3.5 times (1.76% more eggs) than the control group though it was statistically non-significant. On the contrary, Kim and Patterson [11] remarked that hens fed with diet containing 3000 ppm Zn brought about a significant decrease in egg production, egg weight and shell thickness.

Extra zinc supplementation in the diet was practised as a mode for inducing moulting in birds, in order to boost the production in subsequent cycles. Zinc can also cause follicular atresia and interrupt the laying of eggs [12,13]. Shippee et al. [14] also found that the addition to the layer ration with 10,000 ppm (1 per cent) zinc, such as zinc oxide or zinc acetate for 14 days, caused a decline in hen day egg production from 60 to 0% in 6 days, which was marked with induction of moulting.

The trend of egg production recorded in this study fully agreed with reports of Plaimast et al. [15] and Bahakaim et al. [16] who supplemented both inorganic (zinc sulfate) and organic (amino acid chelate) zinc at levels of up to 600 ppm in the diet and it did not have a detrimental impact on the production of eggs and quality of production.

Contrary to the above reports so far discussed, Aghai et al. [17] concluded that dietary supplementation with Zn beyond 40 mg/kg of diet improved egg production and eggshell strength in laying Japanese quail.

In the present study as discussed, zinc supplementation and UVB treatment did not bring any significant changes in egg production. Nowadays, the usage of insect traps lighted with UV bulbs is more common in poultry houses. The hypothalamic photosexual responses to direct UV reception, are very weak because of the shorter wavelength and poor penetration power of UV rays. Hence it did not involve hypothalamic control of reproduction in a controlled environment of poultry houses [18].

In the present study, the UVB exposure for a long period of 3h/day did not make any change in egg production. These findings were in accordance with the reports of Kuhn et al. [19] and Hogsette et al. [20] who demonstrated that UVB light treatment could cause any significance in laying performance.

Contrary to the present findings, Barrot et al. [21], Carson and Junnila [22] and Wei et al. [23] recorded an increase in egg production, when hens were exposed to UV lamps which could be attributed to the favourable effects brought about by germicidal UV light of 253.7 nm wavelength. The beneficial effects of UVB exposure in older birds to maintain egg production were reported by England and Ruhunke [24].

Even though UV light did not play any role in photosexual responses related to egg production, it possessed the ability to synthesise vitamin D₃ which would lead to improved egg production in birds receiving vitamin D₃ deficient diet.

The current study was in accordance with reports of Carson and Beall [25] and Schutkowski et al. [26], as they found neither UVB light exposure nor vitamin D₃ supplementation brought any significant change in egg production. Since, daily supplementation of white incandescent light with 8h to 12h of UV radiation did not cause a shift in timing of the release of pre-ovulatory luteinising hormone which determined the time of oviposition [27].

3.3 Feed Consumption

Results of the effect of UVB light treatment and zinc feeding on mean feed intake (g)/ bird/ day between 29 and 40 weeks of age are presented in Table 5.

The effect of UVB light treatment and inclusion of inorganic zinc sulphate in the diet was highly significant ($p < 0.01$) for mean daily feed consumption during the entire three periods of the experiment. Throughout period I, significantly lower feed consumption was observed in a group of birds fed with inorganic zinc sulphate without UVB treatment (106.16 g) and in the UVB light-treated group without zinc supplementation (108.04 g) compared to control group I (110.00 g). However, the highest feed intake was observed in a group which received UVB light treatment along with zinc supplementation (115.80) during the period I. In periods II and III, the group with zinc supplementation alone (group II) displayed a similar pattern, showing lower feed intake relative to other groups, while the group with both UVB light therapy and zinc supplementation (group IV) showed the highest feed consumption. Feed intake was significantly different for all the groups during period II and period III. The overall feed consumption was

Table 5. Mean \pm S.E. daily feed consumption (g/bird) under UVB light treatment and zinc feeding

Treatment Groups	Period/age in weeks			
	Period I 29-32	Period II 33-36	Period III 37-40	Overall mean 29-40
I	110.00 ^b \pm 1.27	115.54 ^b \pm 0.83	104.29 ^b \pm 0.26	109.94 ^b \pm 0.72
II	106.16 ^a \pm 0.40	112.68 ^a \pm 0.26	102.68 ^a \pm 0.07	107.17 ^a \pm 0.24
III	108.04 ^{ab} \pm 0.16	114.82 ^b \pm 0.22	106.25 ^c \pm 0.07	109.70 ^b \pm 0.13
IV	115.80 ^c \pm 0.15	116.96 ^c \pm 0.13	106.79 ^d \pm 0.16	113.18 ^c \pm 0.14
p-value	0.00**	0.00**	0.00**	0.00**

Means bearing different superscript within a column differ significantly

** Highly significant ($p < 0.01$)

found to be significantly different ($p < 0.01$) and mean overall feed intake was lower in group II (107.17 g) while no significant differences were seen in control group I (109.94 g) and UVB light-treated group III (109.70) however, the highest mean feed intake was noticed in group IV (113.18). On the other hand, feed intake in groups with both UVB treatment and zinc supplementation was greater. Hens in group II did not consume feed as generally expected. The overall mean of feed intake was significantly different among treatments and the lowest value was recorded in layers fed diet with zinc supplementation alone.

The mean daily feed consumption during the entire three periods of the experiment and the overall feed consumption during the experimental period were significantly ($p < 0.01$) affected by UVB light treatment as well as inorganic zinc sulphate in diet, as summarized in Table 5.

During the present research study, significantly lower feed consumption was observed in the group of birds fed with inorganic zinc sulphate without UVB treatment (group II) compared to other groups. Such a finding was in full agreement with reports of Shippee et al. [14] and Tabatabaie et al. [28] who reported reduced feed intake in zinc fed group.

Contrary to the present study findings, Idowu et al. [29] and Naz et al. [30] reported that additional supplementation of Zn enhanced feed intake and improved growth in poultry birds. However, Sunder et al. [31] found that there was no major effect on body weight, feed consumption and feed conversion efficiency by supplementing Zn to the basal diet at graded levels.

The addition of a high level of zinc decreases palatability [32] and depresses appetite [33]

which could contribute to reduced feed intake and subsequent production performance. This hypothesis was well supported by the findings of Kim and Patterson [11] who recorded that supplementation of zinc (3,000 to 20,000 μ g Zn/g diet) could reduce hens' feed intake due to the anorectic effect at high levels.

In the current study feed intake was higher in the group of birds which received UVB light treatment along with zinc supplementation than the birds with zinc supplementation alone. These results were in line with previous studies of Kuhn et al. [19] who exposed birds to UVB light (76 mW/cm²) for varying periods ranging from 0 to 6h per day, where they found a significant increase ($p < 0.01$) in food intake. However, Schutkowski et al. [26] reported that UVB radiation and dietary vitamin D₃ did not influence food intake. Similarly, Lewis et al. [34] reported that feed intake, growth and feed conversion efficiency were similar for all groups, irrespective of UV supplementation (60-160 mW/m² intensity at bird-head height) when turkeys were given nutritionally complete diets. Therefore feed consumption and feed conversion efficiency were not significantly affected by UV supplementation in turkeys.

Contrary to the findings discussed so far, Lewis et al. [27] reported that the supplementation of continuous incandescent illumination with 12 h of UVA from a blacklight-blue lamp inexplicably reduced feed intake in laying hens during the period of exposure, even though the total daily feed intake was not significantly different from that of non-UVA supplemented controls.

In the present study, it was recorded that during period II (33 to 36 weeks) the egg production performance was at its peak and the feed consumption was increased in groups with UVB treatment along with zinc supplementation. This

may be due to the enhanced calcium-binding protein formation as well as calcium absorption from the duodenal segment which was channelized for egg laying as reported by Williams et al. [35] and Tian et al. [36]. Higher feed intake and egg production were recorded in period II as per the present study.

4. CONCLUSION

There were no significant differences were observed in egg production of birds due to UVB light exposure and zinc supplementation, since the present study, neither caused any toxicity nor imparted any negative effect on the production performance. However, a highly significant effect was observed on feed consumption between the treatment groups.

ACKNOWLEDGEMENT

The authors are thankful to Central Instruments laboratory, Kerala Veterinary and Animal Sciences University for providing the facilities needed for carrying out the research.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. McConnon A, Fletcher PL, Cade JE, Greenwood DC, Pearman AD. Differences in perceptions of functional foods: UK public vs. nutritionists. *Nutr. Bull.* 2004;29:11-18.
2. Landstrom E, Hursti UKK, Becker W, Magnusson M. Use of functional foods among Swedish consumers is related to health-consciousness and perceived effect. *Br. J. Nutr.* 2007;98:1058-1069.
3. Skrivan M, Skrivanova V, Marounek M. Effects of dietary zinc, iron, and copper in layer feed on distribution of these elements in eggs, liver, excreta, soil, and herbage. *Poult. Sci.* 2005;84: 1570-1575.
4. Singh,V, Pathak V, Akhilesh, KV. Modified or enriched eggs: A smart approach in egg industry. *Am. J. Food Technol.* 2012;7:266-277.
5. Megha PS, Ramnath V. Production of Vitamin D3 Enriched designer chicken eggs by direct Ultra Violet Blue (UVB) Light Exposure. *J. Anim. Res.* 2021;11(6):977-982.
6. Megha PS, Ramnath V. Influence of direct ultraviolet blue (UVB) light exposure on serum levels of vitamin D3, calcium and phosphorus in layer chickens. *J. Pharm. Innov.* 2023;12(1):16-19.
7. Megha PS, Ramnath V, Raji K, Babitha V, Binoj C. Production of zinc enriched designer eggs through dietary supplementation. *J. Vet. Anim. Sci.* 2021;52(1):77-80.
8. Supplee WC, Blamberg DL, Keene OD, Combs GF, Romoser GL. Observations on zinc supplementation of poultry rations. *Poult. Sci.* 1958;37(5):1245-1246
9. Stahl JL, Cook E, Sunde ML. Zinc supplementation: Its effect on egg production, feed conversion, fertility, and hatchability. *Poult. Sci.* 1986;65(11):2104-2109.
10. Durmus IS, Atasoglu C, Mizrak CENGIZHAN, Ertas S. Kaya. Effect of increasing zinc concentration in the diets of brown parent stock layers on various production and hatchability traits. *Archiv. Anim. Breed.* 2004;47(5):483-489.
11. Kim WK, Patterson PH. Effects of dietary zinc supplementation on hen performance, ammonia volatilization, and nitrogen retention in manure. *J. Environ. Sci. Health B.* 2005;40(4):675-686.
12. Scott JT, Creger CR. The use of dietary zinc as an effective molting agent in laying hens. *Poult. Sci.* 1976;55(5):2089.
13. Berry WD, Brake J. Comparison of parameters associated with moult induced by fasting, zinc and low dietary sodium in caged layers. *Poult. Sci.* 1985;64(11): 2027-2036.
14. Shippee RL, Stake PE, Koehn U, Lambert JL, Simmons Iii RW. High dietary zinc or magnesium as forced-resting agents for laying hens. *Poult. Sci.* 1979;58(4):949-954.
15. Plaimast H, Sirichakwal P, Puwastien P, Kijparkorn S. Effect of supplementary zinc from organic and inorganic sources on laying performance and zinc deposition in eggs. *Thai. J. Vet. Med.* 2008;38(3):47-53.
16. Bahakaim A, Abdel Magied H, Osman S, Omar A, Abdel Malak NY, Ramadan N. Effect of using different levels and sources of zinc in layer's diets on egg zinc enrichment. *Egypt. Poult. Sci. J.* 2014;34(1):39-56.
17. Aghai A, Khosravinia H, Mamuoiei M, Azarfar A, Shahriari A. Effects of dietary supplementation of zinc and α -tocopheryl

- acetate on performance and zinc concentrations in egg and tissues of Japanese quails. *Poult. Sci. J.* 2017;5(1):57-64.
18. Lewis PD, Gous RM. Responses of poultry to ultraviolet radiation. *World's poult. Sci. J.* 2009;65(3):499-510.
 19. Kuhn J, Schutkowski A, Kluge H, Hirche F, Stangl GI. Free-range farming: A natural alternative to produce vitamin D-enriched eggs. *Nutr.* 2014;30(4):481-484.
 20. Hogsette JA, Wilson HR, Semple Rowland SL. Effects on White Leghorn hens of constant exposure to ultraviolet light from insect traps. *Poult. Sci.* 1997;76(8):1134-1137.
 21. Barott HG, Schoenleber LG, Campbell LE. The effect of ultraviolet radiation on egg production of hens. *Poult. Sci.* 1951;30(3):409-416.
 22. Carson JR, Junnila WA. Ultraviolet irradiation of the turkey hen. *Poult. Sci.* 1953;32(5):871-873.
 23. Wei Y, Zheng W, Li B, Tong Q, Shi H, Li X. Effects of b-wave ultraviolet supplementation using light-emitting diodes on caged laying hens during the later phase of the laying cycle. *Anim.* 2020;10(1):15.
 24. England A, Ruhnke I. The influence of light of different wavelengths on laying hen production and egg quality. *World's Poult. Sci. J.* 2020;76(3):443-458.
 25. Carson JR, Beall G. Absence of response by breeder hens to ultraviolet energy. *Poult. Sci.* 1955;34(2):256-262.
 26. Schutkowski A, Kramer J, Kluge H, Hirche F, Krombholz A, Theumer T, Stangl GI. UVB exposure of farm animals: study on a food-based strategy to bridge the gap between current vitamin D intakes and dietary targets. *PLoS One.* 2013;8(7):e69418.
 27. Lewis PD, Perry GC, Sherwin CM, Moinard C. Effect of ultraviolet radiation on the performance of intact male turkeys. *Poult. Sci.* 2000b;79(6):850-855.
 28. Tabatabaie MM, Aliarabi H, Saki AA, Ahmadi A, Siyar SA. Effect of different sources and levels of zinc on egg quality and laying hen performance. *Pak. J. Biol. Sci.* 2007;10(19):3476-3478.
 29. Idowu OMO, Ajuwon RO, Oso AO, Akinloye OA. Effects of zinc supplementation on laying performance, serum chemistry and Zn residue in tibia bone, liver, excreta and eggshell of laying hens. *Int. J. Poult. Sci.* 2011;10(3):225-230.
 30. Naz S, Idris M, Khalique MA, Alhidary IA, Abdelrahman MM, Khan RU, Chand N, Farooq U, Ahmad S. The activity and use of zinc in poultry diets. *World's Poult. Sci. J.* 2016;72(1):159-167.
 31. Sunder GS, Panda AK, Gopinath NCS, Rao SR, Raju MVLN, Reddy MR, Kumar CV. Effects of higher levels of zinc supplementation on performance, mineral availability, and immune competence in broiler chickens. *J. Appl. Poult. Res.* 2008;17(1):79-86.
 32. Fox MRS. Zinc in human biology. Springer, London. 1989;365-370.
 33. Brink MF, Becker DE, Terrill SW, Jensen AH. Zinc toxicity in the weanling pig. *J. Anim. Sci.* 1959;18(2):836-842.
 34. Lewis PD, Perry GC, Morris TR. Ultraviolet radiation and laying pullets. *Br. Poult. Sci.* 2000a;41(2):131-135.
 35. Williams SN, Miles RD, Quart MD, Campbell DR. Short-term high level zinc feeding and tissue zinc concentration in mature laying hens. *Poult. Sci.* 1989;68(4):539-545.
 36. Tian XQ, Chen TC, Lu ZHIREN, Shao QING, Holick MF. Characterization of the translocation process of vitamin D₃ from the skin into the circulation. *Endocrinol.* 1994;135(2):655-661.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/114856>