

Evaluation of Oestrus Response, Pregnancy Rate and Oestradiol-17 β of Nigerian Goat Following Synchronisation with Prostaglandin F_{2 α}

C. E. Dung ^{a*}, M. U. Kawu ^b, H. J. Makun ^c, H. Buhari ^b, L. S. Yaqub ^b,
I. O. Suleiman ^d, J. J. Kalang ^e, F. E. Sa' Ayinzat ^f and J. A. Oko ^b

^a Department of Veterinary Physiology, Pharmacology and Biochemistry, University of Jos, Nigeria.

^b Department of Veterinary Physiology, Ahmadu Bello University, Zaria, Nigeria.

^c National Animal Production Research Institute, Shika, Ahmadu Bello University, Zaria, Nigeria.

^d Department of Animal Science, Bayero University, Kano, Nigeria.

^e Department of Veterinary Medicine, Surgery and Radiology, University of Jos, Jos, Nigeria.

^f Department of Theriogenology and Production, University of Jos, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/CJAST/2022/v41i383977

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here:

<https://www.sdiarticle5.com/review-history/90939>

Original Research Article

Received 05 June 2022
Accepted 09 August 2022
Published 29 October 2022

ABSTRACT

Reproductive inefficiency is one of the most important and cost limiting factor to profitable small ruminant production in the tropics. The aim of this study was to evaluate the influence of breed on oestrus response, reproductive and oestradiol 17 β profiles in Red Sokoto (RS), Sahelian (SH) and West African Dwarf (WAD) goats following synchronisation with prostaglandin F_{2 α} (PGF_{2 α}). Thirty (N = 30) apparently healthy; RS (n = 10), SH (n = 10) and WAD (n=10) goats, aged between 1.5 – 2.0 years, and body condition score of 3.0 – 4.0 were used for the study. Each doe was synchronised with 1 mL (0.26 mg) cloprostenol sodium (Synchromate[®]) given as a single intra-muscular injection on Day 0. Prior to synchronisation, a 5mL blood sample was collected from each doe on Day 0, (pre- synchronisation sample) and subsequently on Day 1, Day 2 and Day 3 post-synchronisation for determination of serum reproductive and oestradiol17 β profiles. The findings of the study were as follows: Oestrus response rates (%): RS; 100.0; SH; 100.0; WAD; 0.0; pregnancy rate (%): RS; 70.0; SH; 20.0; WAD; 0.0. The Sahelians that kidded (SHK) had significantly higher (P < 0.05)

*Corresponding author: E-mail: edward.dung@yahoo.com;

serum progesterone concentration than Sahelian that did not kid (SHN); Red Sokoto goats that kidded (RSK) had significantly higher ($P < 0.05$) serum concentrations of oestradiol-17 β than Red Sokoto that did not kid (RSN); oestradiol-17 β was higher in RSK than SHN ($P < 0.05$). It is, therefore, concluded that: Red Sokoto goats responded better to PGF_{2 α} -based oestrus synchronisation than Sahelian and WAD. Oestradiol-17 β activity was comparatively higher in RS than in SH and WAD between Day 1 and Day 3 post-synchronisation with PGF_{2 α} . WAD goats were completely non-responsive to oestrus synchronisation with PGF_{2 α} . It is therefore recommended that WAD goat will require more time (approximately 6 months) to adapt to the Northern Guinea Zone for their hypothalamo-pituitary- Ovarian axis to be activated sufficiently to cycle, ovulate and develop functional corpus luteum that will respond to PGF_{2 α} .

Keywords: Goat; oestrus response; oestrus synchronisation; prostaglandin F_{2 α} .

1. INTRODUCTION

“Goats are the most prolific of all domesticated ruminants under tropical and subtropical conditions and are able to breed throughout the year” [1]. “They provide meat, milk and skin, and other by-products such as manure to maintain soil fertility” [2]. Goats are hardy animals and are ubiquitous in Nigeria. They are year-round breeders with age and weight at first oestrus being 4-6 months and 10-18 kg, respectively [1]. “The Nigerian goat population is the largest in Africa and the 4th largest in the world after India, China and Pakistan” [3,4]. “The three recognized breed of goats in Nigeria are: Red Sokoto, Borno Sahel and West African Dwarf. The Red Sokoto goat is probably the most wide spread and well-known type in Nigeria” [5,6] and consist of about 60% of Nigerian goat population [7,8], (RIM, 1992). “Exogenous hormones are used to modify the physiological chain of events involved in the sexual cycle, while the non-hormonal methods of oestrus synchronisation involve the use of light control or exposure to a male” [9-12]. “In the doe, the window of opportunity is generally greater during the luteal phase, which is of longer duration and more responsive to manipulation” [13]. “It is essential that any oestrus synchronisation technique should not only establish synchrony, but also ensure reasonable levels of fertility in the synchronised cycle” [14]. “The advantages of oestrus synchronisation in goats include, amongst others, better oestrus detection, increased application of Artificial Insemination (AI), multiple ovulation and embryo transfer (MOET), shortening of kidding intervals, concentration of kid crop, improved management of pregnant does, induction of puberty in doelings and the more efficient use of labour and animal facilities” [14, 15]. “An easy-to-apply method of oestrus synchronisation in goats is by the use of prostaglandins to cause

luteolysis so as to induce the subsequent follicular phase of the oestrous cycle. In small ruminants, prostaglandin F_{2 α} is the primary luteolytic agent” [16]. “Dinoprost tromethamine marketed as Lutalyse® and Carboprost® are frequently used natural prostaglandins, while cloprostenol sodium, marketed as FenprostenoL®, Estrumate® and EstroPLAN®, is a synthetic prostaglandin” [17,14]. “Factors reported to affect oestrus response and subsequent fertility following administration of prostaglandin or its analogues include the dose level of the prostaglandin” [6,19], “the interval between administration of the prostaglandin” [6], “the responsiveness of the corpus luteum to the prostaglandin/ stage of the oestrous cycle” [6, 20], season [6]. “With the discovery of prostaglandin F_{2 α} as a luteolysing agent in farm animals as a synchronisation agent, it has been implemented to shorten the cycle and synchronize females for breeding. Both single” [21] and “double prostaglandin F_{2 α} injections, given at 10–11 day intervals in does have been successful” [22]. The limitation of this regimen is however that it is only effective in cycling does.

2. MATERIALS AND METHODS

2.1 Experimental Location

The study was carried out at the Small Ruminant Research Programme, National Animal Production Research Institute (NAPRI), Shika, Ahmadu Bello University, Zaria, Nigeria. Shika is located between latitudes 11 and 12° N and between longitudes 7 and 8° E at an altitude of 640 m in the Northern Guinea Savanna zone. The average annual rainfall in Shika is approximately 1100 mm, and mainly during the months of April to October. The maximum ambient temperature range in Shika is 27 – 35°C depending on season [23].

2.2 Experimental Animals

Thirty apparently healthy and nonpregnant RS (n=10), SH (n=10) and WAD (n=10) selected from the main flock were used for the study. The ages of the does range between 1.5 – 2.0 years, parity, 1-2; and body condition score (BCS), of 3.0 - 4.0 [24]. The does were identified by means of plastic ear tags. They were fed *Digitaria smuttsi* hay as basal diet and 15% crude protein concentrate ration. Water was provided ad libitum

2.3 Oestrus Synchronisation Protocol and Blood Sampling

Oestrus synchronisation was conducted in the late rainy season (August to September). The goats were synchronised using 1ml containing 0.26 mg cloprostenol sodium (Synchromate^(R) BREMER PHARMA, GMBH, Germany), The cloprostenol sodium was administered to each doe by a single deep intra-muscular injection (thigh). Treatment with cloprostenol sodium was carried out between 8.00am and 9.00am on day 0. The does were randomly allotted into 6 pens with 5 does each, and were allowed to stay together for one week with a proven buck in each pen for the purpose of oestrus detection and natural breeding. Thereafter, the does were separated from the buck and returned to the main flock. Five millilitres (5 ml) of blood sample was collected from each doe by jugular vein puncture on Days 0, 1, 2 and 3 of synchronisation. Day 0 sample was collected immediately prior to injection of cloprostenol sodium (Pre-synchronisation sample).

2.4 Oestrus Detection and Breeding

Beginning from day 1 post-synchronisation, the treated does were observed visually for behavioural manifestation of oestrus twice daily between 8.00 am and 10.00 am, and 4.00 pm and 6.00 pm, respectively. The does were exposed to proven sexually active bucks in the ratio of 1: 5 (male: female) to aid oestrus detection [25]. The does were considered to be in oestrus when they stood to be mounted by females (homosexual mount), or by males (heterosexual mounts). Other signs such as vigorous tail-wagging, reddened and swollen vulva, clear mucoid vaginal discharge, restlessness, frequent bleating and frequent adaptation of urination posture were also used to determine oestrus [26]. Mated does were allowed to run with other does in the group. Oestrus

activity occurring within 24 - 48 h (1-2days) was classified as synchronised.

2.5 Pregnancy Diagnosis

Pregnancy was diagnosed using real-time B-mode trans-abdominal ultrasound equipment (Aloka 500 V with a 3.5 MHz transducer, Corometrics medical systems, Wallingford, CT, USA). The procedure was carried out between days 30 and 35 after natural mating. Food and water were withheld over night for 12 hours before commencement of scanning in the morning (8.00 am and 10.00 am). Scanning was performed using the shaved hairless inguinal region of the does. The does were gently restrained by two persons on an examination table, while the hind legs of the does were folded up at the time of scanning for proper placement of the probe. An ultrasound coupling gel was applied each time to the probe to develop good contact and to remove hair between probe and the animal skin. A doe was confirmed pregnant by imaging apparent conceptus (anechoic, elongated structure) within the uterine fluid.

2.6 Determination of Oestrus Response Rate, Pregnancy rate and Serum Oestradiol -17 β Profile Profile

Oestrus response, and Pregnancy Rate, were determined as follows:

2.6.1 Oestrus response rate (%)

The number of does that showed standing oestrus and were subsequently mated, over the total number of does in each treatment group multiplied by 100 [27].

2.6.2 Pregnancy rate (%)

The total number of does that kidded divided by the total number of does mated multiplied by 100 (Sahare et al. 2009).

The Oestradiol-17 β assay was carried out using Microplate Enzyme Immuno-assay oestrogen test system (Accubind, ELISA Microwell; Monobind Inc, Lake Forest, CA 92630, USA: Product Code: 4925-300).

2.7 Data Analysis

Oestrus response rate, and pregnancy rate were expressed as percentages. While serum oestradiol -17 β concentration were expressed as mean (\pm SEM) and analysed using analysis of

variance (one- way and Two-way repeated measures) and Duncan post hoc-test. Graphpad Prism (version 5.0) statistical package was used for the data analysis. Values of $P < 0.05$ were considered significant.

3. RESULTS

The oestrous response rate, following synchronization with PGF_{2α} for Red Sokoto (RS), Sahelian (SH) and West African Dwarf (WAD) goats were 100.0%, 100.0%, and 0.0%, respectively as shown on Table 1. The pregnancy rate, following synchronization with PGF_{2α} for Red Sokoto (RS), Sahelian (SH) and West African Dwarf (WAD) goats were 70.0%, 20.0%, and 0.0%, respectively. The oestradiol - 17 β concentrations following synchronization with PGF_{2α} for Red Sokoto (RS), Sahelian (SH) and West African Dwarf (WAD) goats that kidded and those that did not kid is shown in Table 2.

4. DISCUSSION

The 100% synchronisation of oestrus achieved with prostaglandin in the present study is higher than the values of 64% and 84% following single

and double injections, respectively, with prostaglandin in Brown goats have also been reported in the same environment (Ogunbiyi et al. 1980). This value is in agreement with 90% estrus response recorded in Nadooshani goats in Iran (Bitaraf et al. 2007). The differences in these results may be due to breed, nutritional status, prevailing meteorological parameters, dose and potency of the agent used, stage of oestrous cycle or stage of maturity of the corpus luteum (Niswender et al. 2000). The pregnancy rates of 70 % and 20 % recorded for RS and SH goats, respectively, is lower than values reported in Brown goats following double injections of PGF_{2α} (Ogunbiyi et al. 1980). The higher pregnancy rates in Red Sokoto goats compared to the Sahelians suggests a better physiological adaptation of Red Sokoto goats as compared to Sahelian or West African dwarf goats to the Savannah agro-ecological zone [28,29]. Furthermore, goats that exhibited sexual receptivity had higher levels of oestradiol 17 β throughout the Days 0,1,2 and 3 and in the overall. This may be due to the fact that high oestradiol 17β levels are believed to cause surge in GnRH and consequently LH peak resulting in spontaneous ovulation towards the end of oestrus [14].

Table 1. Synchronization with prostaglandin F_{2α} in Red Sokoto, Sahelian and West African Dwarf goats

Fertility index	Breed		
	Red Sokoto (n=10)	Sahelian (n=10)	West African dwarf goats (n=10)
Oestrus response rate (%)	100.0 (10/10)	100.0 (10/10)	0.0 (0/10)
Pregnancy rate (%)	70.0 (7/10)	20.0 (2/10)	0.0 (0/10)

(n) = Number of animals that showed oestrus/ Became pregnant/Kidded

Table 2. Mean (±SEM) Serum Concentration of Oestradiol 17- β (pg/ml) in Red Sokoto, Sahelian, and West African Dwarf Goat Between Day 0 and Day 3 following Oestrous Synchronization with Prostaglandin F_{2α}

Breed	Oestradiol 17 β (ng/ml)				
	Day0	Day1	Day2	Day 3	Mean (±SEM)
RSK	357.00±111.2	300.00± 286.70	450±99.6	460.0 ±209.0	392 ±38.4 ^a
RSN	0.00 ± 0.00	120.0± 10.0	0. 0±0.0	150±32.0	67 .± 39.45 ^b
SHK	292.00±0.00	180.±5.0	502. 0 ± 0.0	417±0.0	348 ±70.62
SHN	138.00±323.00	180.±56	258.0 ± 85	99±91	168.0 ± 34.0 ^a
WADN	1380.0±5.40	477± 34	327.0 ± 225	612± 147	388.0± 101.8

a, b = $p < 0.05$

(n) = Number of animals/Samples analysed, RSK =Red Sokoto Goat that Kidded, RSN = Red Sokoto Goat that did not Kid, SHK = Sahelian Goat that Kidded, SHN = Sahelian Goat that did not Kidd, WADN = West African Goats that did not kidded

5. CONCLUSIONS

There was an increase in the oestrus response and fertility in Red Sokoto and Sahelian goats than West African Dwarf goats. The serum level of reproductive hormones (Oestradiol 17- β) were generally higher in Red Sokoto than in SHK and WAD following prostaglandin-based oestrus synchronization. The poor ovarian response in WAD may be due to poor adaptability of the breed to Northern Guinea Savannah zone.

ETHICAL APPROVAL

Ethical clearance for the experiment was obtained from the Ahmadu Bello University Committee on Animal Use and Care (ABUCAUC). No. ABUCAUC/2021/064.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Mamabolo MJ, Webb EC. Goat production survey – fundamental aspects to model goat production systems in Southern Africa: a case study-agricultural commission-WITFOR; 2005.
- Boyazoglu J, Hatziminaoglou I, Morand-Fehr P. The role of the goat in society: past, present and perspectives for the future. *Small Ruminant Research*. 2005; 60:13-23.
- Blench R. Traditional livestock breeds: geographical distribution and dynamics in relation to the ecology of West Africa. London. Overseas Development Institute. 1999;67.
- Food and Agriculture Organization of the United Nations. Production handbook. FAO. Rome. 1999;53:251.
- Osinowo OA. Small ruminant statistics. Nigeria Monograph. 1992;1-5.
- Kawu MU. Post-partum uterine and ovarian changes and the effects of oestrus synchronisation on rebreeding in the savannah Brown goats. M Sc. thesis submitted to the Post Graduate School, Ahmadu Bello University, Zaria. 2000;25-34.
- Molokwu ECI, Igonu MO. Reproductive performance and pattern in the Brown goat of the Nigerian Savannah zone, Proceedings of the 4th World Conference on Animal Production, Argentina. 1978; 578-590.
- Wilson RT. Small ruminant production and the small ruminant genetic resource in Tropical Africa. F.A.O. Rome, Italy. 1991;231.
- Fonseca JF, Torres CAA, Costa EP, Maffili VV, Carvalho GR, Alves NG. Progesterone profile and reproductive performance of Oestrus-induced Alpine goats given hCG five days after breeding. *Animal Reproduction*. 2005;60:75-80.
- Igbokwe IO, Grema HA, Ikpo AE, Mshelbwala FM, Igbokwe NA. Unilateral cryptorchidism in Nigerian Sahel bucks. *International Journal of Morphology*. 2009; 27(3):805-810.
- Maina A, Waruiru M, Mbuthia PG, Munyua W, Otieno R, Mutune M. Prevalence of gastrointestinal parasites in indigenous chickens ???; 2017.
- Zakari AY, Molokwu, ECI, Osori DIK. Effect of season on the oestrous cycle of cows (*Bos indicus*) indigenous to northern Nigeria. *The Veterinary Record*. 1981;109:213-215.
- Wildeus S. Current concepts in synchronization of oestrus: sheep and goats. *Journal of Animal Science*. 2000; 77:1-14.
- Rahman ANMA, Abdullah RB, Wan-Khadajah WE. Oestrus synchronisation and superovulation in goats: a review. *Journal of Biological Science*. 2008;8:1129-1137.
- Abecia JA, Forcada F, Gonzalez-Bulnes A. Hormonal control of reproduction in small ruminants. *Animal Reproduction Science*. 2012;130:173- 179.
- McCracken J, Glew ME, Scaramuzzi RJ. Corpus luteum regression induced by prostaglandin F_{2 α} . *J Clin Endocrin Metab*. 1979;30:544-546.
- Bello AA. Characterization of oestrus, endocrine events and fertility in Red Sokoto goat following comparative oestrus synchronisation with prostaglandin and progestagen (?). MSc. Thesis submitted to Ahmadu Bello University, Zaria; 2011.
- Omontese BO, Rekwot PI, Makun HJ, Ate IU, Rwuaan JS. Induction of oestrus in Sahel goats using fluorogestone acetate (FGA) sponges and equine chorionic gonadotrophin (eCG). *Sokoto J Vet Sci*. 2012;10(2):21-25.
- Tauheed M. The effect of dose and parity on oestrus response and fertility following

- single treatment with prostaglandin F2a in anoestral Red Sokoto goats during the mid-rainy season. DVM Project submitted to the Faculty of Veterinary Medicine. Zaria: Ahmadu Bello University; 2010.
20. Lassal A, Hernandez-Cerón J, Rodriguez-Maltos R, Gutierrez CG. The influence of the corpus luteum on ovarian follicular dynamics during estrous synchronisation in goats. *Anim Reprod Sci.* 2004;84: 369-375.
 21. Greyling JPC, Van Niekerk CH. Effect of pregnant mare serum gonadotrophin (PMSG) and route of administration after progestagen treatment on oestrus and LH secretion in the Boer goat. *Small Ruminant Research.* 1990;3:511–516.
 22. Khanum SA, Hussain M, Kausar R. Manipulation of estrus cycle in dwarf goat (*Capra hircus*) using estrumate under different managemental conditions. *Animal Reproduction Science.* 2006; 92:97–106.
 23. Taiwo BBA, Buvenandran V, Adu IF. Effects of body condition on the reproductive performance of Red Sokoto goats. *Nigerian Journal of Animal Production.* 2005;32:1-6.
 24. Spahr LI. Body condition scoring in meat goats. *Penn State College of Agricultural Science.* 2005;23:11-15.
 25. Abecia JA, Forcada F, Gonzalez-Bulnes A. Pharmaceutical control of reproduction in sheep and goats. *Vet Clin Food Anim Prac.* 2011;27:67-79.
 26. Fatet A, Pellicer-Rubio MT, Leboeuf B. Reproductive cycles of goats. *Animal Reproduction Science.* 2011;124: 211-219.
 27. Omontese BO, Rekwot PI, Rwuaan JS, Ate IU, Makun HJ. Comparative estrus induction in indigenous Sahel goats using two progestagens (CIDR and FGA) and eCG. *Livest Res Rural Dev.* 2013;25(4):66.
 28. Baldassarre H, Karatzas CN. Advanced assisted reproductive technologies (ART) in goats. *Animal Reproduction Science.* 2004;82–83:255–266.
 29. Fasanya OOA, Molokwu ECI, Eduvie LO, Dim NI. Dietary supplementation in savanna brown goats. II. Gestation and post-partum activity in primi-parous does. *Food Agric. Org. Animal Reproduction Science.* 1992;29:167-174.

© 2022 Dung et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/90939>