



Integrated Farming is an Effective Multiple Water use System to Enhance the Livelihood Security of Small and Marginal Farmers: A Review

S. Manivannan ^{a*} and V. Kasthuri Thilagam ^a

^a *ICAR-Indian Institute of Soil and Water Conservation, Research Centre, Udhamandalam – 643 004, Tamil Nadu, India.*

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2022/v34i930912

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/85109>

Review Article

Received 12 January 2022
Accepted 17 March 2022
Published 17 March 2022

ABSTRACT

The agriculture sector faces numerous challenges like increasing population, decreasing per capita land, fragmented land holdings, land degradation and climate change in 21st Century. The downtrend of per capita land availability poses a severe threat to the sustainability and profitability of agriculture. This issue is severe for small and marginal farmers as they face many problems sustaining single farm enterprises. The global assessment on water scarcity indicates that about 66 % of the global population live under severe water scarcity conditions. Increasing land and water productivity for enhancing livelihood security of rural poor is the challenge for various stakeholders of agriculture and allied sectors. Larger numbers of integrated farming system models have been developed and demonstrated across the globe. In most cases, integrated farming systems combine crops and livestock components. However, water component is rarely included in a few integrated farming system models and needs to be upscaled its research and demonstration on a larger scale. Hence, it is the right time to review the existing integrated farming system models and analyze the research gap on multiple water-use integrated farming systems to derive a future road map. This systematic review analyzed all available integrated farming systems' prospects and retrospects and their impacts on production and productivity. The reviews proved that crop and livestock based integrated farming systems are outperforming and enhancing farmers income, employment opportunities and fulfilling the family's nutritional requirements by optimizing the farm's resources.

*Corresponding author: E-mail: smanicar1997@gmail.com;

However, the role of harvested water and its effect on the integrated farming system is rarely studied. Hence, the research on multiple water use based integrated farming systems needs to be developed and demonstrated in all agro-ecological regions of different countries. The water productivity functions for multiple water use-based integrated farming systems need to be derived and tested to understand the role of harvested water in multiple water use scenarios.

Keywords: Integrated farming system; livestock based IFS; poultry based IFS; fish based IFS; multiple water use.

1. INTRODUCTION

The development of the agricultural sector is the primary concern for all the developing countries as agriculture is an essential sector in action for food production [1]. However, increasing population, fragmented land holdings, decreasing per capita land, land degradation and climate change are challenging agricultural development. Our country's growing population and shrinking land resources leave no scope for further land expansion for agricultural purposes and poses a severe threat to the sustainability and profitability of Agriculture [2]. Though small and marginal farmers constitute 85% of the Indian farming community, they share only 44% of the total agricultural land [3]. Nevertheless, farmers face challenges to generate adequate income and year-round employment with a single commodity [4]. Hence, integrating water resources, agriculture, horticulture, fishery, duckery and other livestock components at the farm level with suitable technologies is the best way to maintain land productivity and increase farmer income while protecting the environment. In the climate change scenario, water is becoming a scarce commodity worldwide. The water scarcity problem can be managed by runoff water harvesting at the farm level and put into multiple uses for agriculture and allied activities. This multiple water use based Integrated Farming system is a systematic approach to use low external inputs for crops and livestock production [5] consisting of various interrelated components [6]. This can be a sustainable alternative to commercial farming systems, particularly for small and marginal farmers, to reverse the resource degradation [7]. Integration of various farm enterprises with harvested water, recycling crop residue and farm-based by-products surely enhance farm income [8] and provide opportunities for maintaining and extending biodiversity [9].

Multiple water use based IFS provides an opportunity to increase the income per unit area by adopting diversification of crops and

livestocks with integration of water. There is much scope for water and nutrient recycling within the system to economize and sustain the entire system and minimizes the dependence on off-farm and fossil fuel-based chemical inputs in farming [10]. Traditionally IFS is effectively undertaken by farmers of Indonesia, China, Malaysia, Vietnam, Rwanda and Thailand [11]. However, these ancient systems have been gradually replaced by single commodity based intensive farming systems [12]. In recent days, the crops and livestock-based IFS are being adopted by farmers to increase net income using river or canal or well water. But, the awareness for integrating harvested rainwater with crops and livestock is being followed to a limited extent and needs to be upscaled in a wider range. This review aims to analyze the research gap on multiple water use integrated farming system to derive a future road map.

2. COMPREHENSIVE INTEGRATED FARMING SYSTEM

IFS aim to sustain the productivity, efficient recycling of farm wastes, better resource utilization, employment generation and risk reduction while maintaining environmental harmony [13,14]. Whereas multiple water use based IFS aims for a sustained increase in agricultural production and water use efficiency. Crop components, horticulture, dairy, livestock, and other subsidiary enterprises are all part of the agricultural system method [15,16,17]. Increasing organic inputs by way of farming system has a crucial role for increasing carbon content [18]. Intensive agricultural systems change the properties of organic matter and induce carbon losses [19] and lead to the low organic matter in soil.

Farming systems significantly affect soil organic matter's quantity and quality [20,21]. No single farm enterprise is likely to sustain the small and marginal farmers without integrated farming system [22]. IFS is a combined economic-environmental model [23] and entrepreneurial

orientation model [24] where by product / waste from one component becomes an input to another [25-28]. IFS was defined by various authors differently like it is an agricultural system that is integrated with livestock [29] an integrated farming system that introduces the changes in the farming system, system for waste recycling with increased input efficiency [30] no-cost farming with increased income [31] and a mixed agricultural system [32].

3. INTEGRATED FARMING SYSTEM MODELS

Various IFS models developed and demonstrated have been reviewed, and impacts on various aspects were analyzed. In any IFS model all the components are interdependent and the resources are recycled in the total farming system. This makes the farmer self-sufficient in inputs and balanced diet of the family. This interaction among the different components were explained by Behera and Mahapatra [33].

4. LIVESTOCK BASED INTEGRATED FARMING SYSTEM

The interaction of soil, water, plant, animal and environment enables the system to be more viable and profitable than the arable farming system. The method of integrating livestock with perennial crops must be comprehensive, considering all system components [34]. Livestock based Integrated farming systems offer optimizing resource utilization rather than maximizing individual elements in the system [9]. Live stock based IFS was described as a mixed animal crop system that often raises the animal component on agricultural waste products [35]. On the other hand, the animal is used to cultivate the soil, provide manure and fuel [36] and have better food security and higher income [37].

The advantages of livestock crop integration are in the following perspectives: (a) Agronomical (b) Economical (c) Ecological (d) Nutritional and (e) Social. Plants and livestock interact to create synergy [34].




Fig. 1. Interactions among different components of an IFS
 Source: Behera and Mahapatra (1998).

Kumar *et al.* [38] found that a combination of crop + duck + fish + goat had produced a good quantity of manure, poultry 2.3 t yr⁻¹, duck 1.6 t yr⁻¹, goatry 2.9 t yr⁻¹, cattle 14.0 t yr⁻¹ and plant residue 11.3 t yr⁻¹ whereas crop + dairy + sheep + rabbit + hen + quails, generated 750 man-days of employment [39]. Strategies should thus incorporate links with existing networks and institutional arrangements in designing crop-livestock integration beyond the farm level [40]. Integration of goats [41] and bovines [42] in IFS would rapidly increase the family income in case of reduced landholding pattern due to disintegration of the family in addition to high protein feeds to goats. Similarly, goat dung can be highly recycled into organic manure that can fertilize the soil and increase the growth of plants [43]. Cattle and pigs are also a cornerstone element for promoting local recycling of nutrients [44] and increasing the farm income [45] through abundant cattle manure [46].



Fig. 2. Advantages of Livestock based IFS

Integrating pig, goats or sheep with the plantation crops will reduce the greenhouse gas emissions [47] in addition to augmenting the quality of life of the farmers [48]. Live stock based IFS is a zero waste approach [49] that prevents wasting residuals [50] to minimize agricultural production costs [51]. Cattle with maize, sugarcane and oil palm IFS [52-54] enhances the soil quality by the decomposing capacity of crop residues and earthworms improvement as bioindicator of soil fertility [55-57]. in addition to increasing soil moisture.

5. FISH INTEGRATED BASED FARMING SYSTEMS

Fish based integrated farming systems plays an essential economic and ecological role,

particularly in Asia, by recycling waste and by-products from other production systems [58,59] that maximize productivity through optimized resource use [60] and reduced input costs [61,62]. Production of fish in ponds, with pigs, duck or chicken rearing in pens, beside or over the ponds constitutes continuous organic fertilization of the pond and increases the efficiency and rentability of livestock and fish culture [63]. Fish farming is an income-generating activity for a few farmers, like other household activities [64,65] which is lower yield than fish based IFS [66]. Integrated fish farming is a zero-waste, low-cost, and low-energy production method in which by-products from one business are recycled as inputs into another [67] resulting in increased family food consumption [68]. Manure wastes from pigs used as a source of nutrients for polyculture of carps, channel catfish and largemouth bass produced maximum fish yield [69]. Intensive fish culture with chicken droppings increased the fish yield by 21 per cent and decreased the feed conversion rate by 0.4 [70]. Integration of animal-fish farming can increase the returns from a limited land area and reduce risk by diversifying crops [71]. The gap between supply and demand can be achieved faster through integrated fish farming enhanced with organic matter, nitrogen, phosphorus, and micronutrients collected in pond [72,73,74].

Integration of aquaculture with agriculture has significant potential to contribute to sustainable intensification and nutrition security goals, reducing local environmental impacts associated with sediment disposal and increasing agricultural production [75,76]. Fish based IFS depending on the financial capacity of the farmer [77] could create horizontal integration of crop production by maximizing their workforce and cash flow that produce large fish at low cost, in line with the local market and consumers' expectations. The low stocking density allows for the use of natural pond productivity, ensures enough food for optimal fish growth and optimizes fish yield [78]. The leading adopter of rice-fish farming system in China has devoted 15 per cent of its rice-suitable farmland to this type of farming, producing 1.2 million tons of fish and other aquatic organisms annually [79]. Rice-fish farming has been advocated in Korea to improve resource efficiency and safeguard the environment [80] and grain yield was significantly higher (4.92 t ha⁻¹) with fish [81]. Rice-fish farming in rainfed lowland increases the organic carbon content of soil by 7 per cent and

exchangeable ammonium by 25 per cent, besides a 6 per cent increase in the available phosphorus [82]. However, rice-fish farming systems inevitably demand significant labour, resulting in high production costs in developed countries [83].

Integrated farming of fish, poultry and vegetables has experimented with a practical approach for sustainable production, income generation and employment opportunity for resource-poor rural households [84]. No supplementary feed was provided to the fish under integration throughout the culturing period. The product of Adama red onion (*Allium cepa*) grown in the plot (260m²) integrated with the poultry and fish components was estimated to be 10, 800kg ha⁻¹. The obtained yield was ranked to a better production level when compared to the product obtained using chemical fertilizers [85].

6. DUCK-BASED INTEGRATED FARMING SYSTEM

The IFS model of fish culture + duck farming + azolla + pulses, given three times more income than conventional farming and in a sustainable manner [86]. The benefit-cost ratio in IFS model is 2.28 compared to the traditional model. The total Global Warming Potential (GWP) of integrated rice-duck farming was 13.3 per cent less than the conventional rice cultivation [87]. The rice duck system can protect and enhance natural resources, ensure food security, and improve the economy [88]. Duck integrated farming is becoming more popular as a source of income for people in rural and urban areas and is one option for meeting community demands for nutritious food that is more disease resistant and develops quicker than native chicken become the foundation of life for most individuals who live in agro-ecosystem wetland areas [89]. Ducks were fed agricultural waste with high crude fibre [90]. In the integrated rice-duck farming (IRDF) system, rice paddies provide food (weeds and pests) for ducks, and ducks play a role in fertilizing rice plants [91].

7. POULTRY BASED INTEGRATED FARMING SYSTEM

IFS involving crop + poultry + dairy + piggery enterprises had a positive advantage in terms of economic returns [92,93]. Incorporation of poultry manure and FYM positively affected the plant growth parameters and yield contributing characters and thus resulted in the highest grain

and straw yield of rice [94]. The gross return and gross margin from poultry and vegetables exhibited an additional income that remarkably contributed to the households' increased income. Furthermore, unlike other livestock, poultry and chickens play an essential role in the smallholder farming systems in developing countries. Local poultry stocks often serve as a significant source of animal protein to the poor since they are accessible even to landless households [95,96]. Integrated farming of fish, poultry and vegetables has experimented with a practical approach for sustainable production, income generation and employment opportunity for resource-poor rural households [84]. The fish were feeding upon the planktons and other organisms harboured in the integrated pond with the aid of poultry waste; poultry waste is either eaten directly by fish or fertilizes pond water to support the plankton community used by fish as natural organic feed [96].

8. RICE AND CATTLE - FISH INTEGRATED FARMING SYSTEM

An integrated farming system of rice, cattle and fish is an improved system for paddy productivity that is combined with livestock. The selection of rice and cattle in farming is based on reciprocity relationships, where rice provides straw and bran for cattle feed. Conversely, cattle produce faeces as organic fertilizer. In rice, plants can improve soil structure, encourage absorption, better humidity, reduce power absorbency, and prevent surface soil crusting [97]. The rice, cattle and fish farming system helped to improve the land productivity, water and air quality and created harmony between the socio-cultural environment of the local community [98]. Intensification and efficiency gains in animal production result in less greenhouse gas emissions per unit of milk and more milk per unit of water [99]. This system produces grain and animal protein simultaneously, generating additional income sources and reducing the adverse effects of agriculture on the environment [100].

Impact of IFS on farm income: Water use productivity of IFS model (Crop + horticulture + dairy + poultry + fishery) was high at Rs. 991 ha cm compared to conventional farming (soyabean+wheat+ fallow), which was Rs. 406 ha cm. Higher net returns of 284 and 176 percent through the pig and duck-based multiple uses of pond water through diversified farming (crop, fruit, livestock and fishery) than the farmers' practice (without integration), respectively [101].

Integration of cattle and rice can increase the value of the BC ratio to 6 compare to BC ratio of 4 in traditional farming system [102]. The income of farmers with the integrated farming system was Rs. 9,086,867 for 1 ha land and two cattle or Rs. 4,543,433 for 1 ha land and one cattle with an BC ratio of 1.56 [103]. However, Multiple water use based IFS has the immense potential to emerge as an effective tool for obtaining maximum income and profitability from available resources [104].

9. MULTIPLE WATER USE BASED INTEGRATED FARMING SYSTEM

Harvest of surface runoff during the rainy season enhances the opportunity for diversification in space and time dimensions. Water productivity is enhanced by about one to two times due to multiple uses of pond water compared to farmers' practice of rainfed crops [101,105]. Meeting the food demand, a farming system approach should be used to reduce risks and uncertainties, increase food production, and improve living standards [106]. According to Islam and Mandal [107] adding 6 cm of additional irrigation to rainfed rice improved productivity by 59 per cent. It was projected that a pond the size of 5 per cent of the total cultivable land might hold enough water for supplemental irrigation. Multiple water use based IFS increase the efficiency of all the available resources of the farm and optimize the use of harvested rain water. It also enhances the water productivity

and water harvesting efficiency that resulting in increased system productivity. A water harvesting pond at the downstream side of the farm land will harvest the runoff water from the entire farm. According to the slope, the catchment area needs to be treated with soil conservation measures like bunding and bench terracing to check soil erosion. Grassed waterways also help direct the water flow towards the pond inlet. Multiple use of harvested rainwater includes crop cultivation, plantation, fishery, livestock, poultry, duckery and small scale industries. Crops and livestock can be selected based on the local conditions and need. Various studies found that the Multiple water use based IFS can improve the economic level of farmers, employment opportunities and fulfil the family's nutritional requirements by optimizing the farm's resources.

The following components of multiple water use system found to be effective.

- Farm pond – Water, Fishery, Duckery
- Crop components – Annual crops, Plantation crops, Agro forestry, Fodder crops
- Livestock components – Dairy animals, Sheep & Goat rearing, poultry, piggery, Sericulture, Rabbitry, Pigeon rearing
- By product based components - Vermiculture, Compost production, Apiary Mushroom cultivation [108].




Fig 3. Schematic representation of Multiple water use based IFS

Table 1. The Benefits of livestock components with crop reported by various researchers

S.No	Livestock Components	Direct benefits	Indirect benefits*	Researchers
1.	Duck + fish + goat	Manure production	Soil health and reduced external inputs	Kumar et al. [38] Mowa et al. [43]
2.	Dairy + sheep + rabbit + hen + quails	Employment Generation	Poverty alleviation	Govardhan et al. [39]
3.	Pig, goats or sheep	Greenhouse gas emissions	Reduced global warming potential	Rati et al. [47]
4.	Fish + chicken	Increased fish yield	Increased Economic status	Rappaport and Sarig, [70]
5.	Rice+ Fish	Increased rice yield	Crop productivity and income	Poonam et al. [81]
6.	Rice+ Fish	Increased organic carbon exchangeable ammonium, available phosphorus in soil	Soil fertility	Sinhababu et al. [82]
7.	Poultry + fish	Increased onion yield	Increased Economic status	Desaleng and Aklilu [85]
8.	Fish + duck + azolla	Increased income	Livelihood security	Mitra et al. [86]
9.	Rice+duck	Reduced global warming potential	Environmental sustainability	Xu et al. [87]
10.	Poultry + dairy + piggery	High economic returns	Livelihood security	Mukherjee, [92] Ravishankar et al. [93]
11.	Rice+ Poultry	Nutritional security	Improved family health	Weigend et al. [95] Endabu et al. [96]
12.	Rice + cattle + fish	Land productivity	Increased Economic status	Suwandi, [98] Gupta et al. [34]

**The indirect benefits are reviewed and synthesized by the authors from various sources*

10. CONCLUSION AND FUTURE PERSPECTIVE

Sustainability through an integrated farming system is environment friendly, socially accepted by farmers and economically viable. IFS helps to maintain agricultural income by lowering production costs, creating a lot of job opportunities, and showing the way to the resilience of the farm productivity. Overall, an integrated farming system achieves multiple objectives: making farmers self-sufficient by ensuring a balanced diet for family members, raising the standard of living by maximizing total net returns and creating more jobs, reducing risk and uncertainty, and maintaining environmental harmony. Adequate IFS models have been developed by various stakeholders involving crops, livestock, fish, poultry and ducks and found to be highly productive and profitable. It can be advocated for small and medium farmers for enhancing their livelihood security. Impacts of IFS on soil health, production, productivity, nutrient recycling, income, waste recycling and crop growth have been reported in sizable quantities. However, the studies on IFS with harvested rainwater is lacking and only limited model has been reported. Hence, the following areas needs to be addressed for understanding the role of harvested rainwater in integrated farming system.

- Hydrology and water balance studies on integrated farming systems and its effect on supplemental irrigation
- Water productivity functions of multiple water use system incorporating crops, livestock and fish components
- Nutrient balance studies on various livestock, fish and crops based IFS models
- Role of IFS models and its impact in climate resilience agriculture

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Nainggolan HL, Aritonang J, Sihombing M, Supriana T, Tafsin M. Structural Modelling of Rice Fields-Buffalo Livestock Based Integrated Agricultural Systems in The Context of Regional Development in Humbang Hasundutan Indonesia. *Journal on Food Agriculture and Society*. 2019;7(2):Article No 106.
2. Gill MS, Singh JP, Gangwar KS. Integrated farming System and agriculture sustainability. *Indian Journal of Agronomy*. 2009;54(2):128–139.
3. GOI. Agricultural statistics at a glance, Directorate of Economics and Statistics, Govt. of India, New Delhi; 2014.
4. Mahapatra IC. Farming systems research challenges and opportunities. *Eastern Indian Farming System Research & Extension Newsletter*. 1992;6:3–10.
5. Little DC, Edwards P. Integrated livestock-fish farming systems. *Food and Agriculture Organization of the United Nations, Rome*. 2003;189.
6. Siddeswaran K, Sangeetha SP, Shanmugam PM. Integrated farming system for the small irrigated upland farmers of Tamil Nadu. In: *Extended Summaries: 3rd International Agronomy Congress, New Delhi, India*. 2012;992-993.
7. Lightfoot C, Minnick DR. Farmer-first qualitative methods: Farmers diagrams for improving methods of experimental design in integrated farming systems. *Journal for Farming Systems Research and Extension*. 1990;2:11-13.
8. Behera UK, Jha KP, Mahapatra IC (2004) Integrated management of available resource of the small and marginal farmers for generation of income and employment in eastern India. *Crop Res*. 2004;27:83-89.
9. Soni RP, Katoch M, Ladohia R. Integrated farming systems - A review. *IOSR J. Agric. Vet Sci*. 2014;7(10):36–42.
10. Ganesan G, Chinnasamy KN, Bala Subramanian A, Manickasundram P. Studies on rice based farming system with duck cum fish culture in the deltaic region of Thanjavur district, Tamil Nadu. *Farming Systems Newsletter*. 1990;1(2):14.
11. Praphan N. Resilient of indigenous knowledge, fight to world crisis. *Isan alternative farming network, Ubonratchathani, Thailand*; 2001.
12. Ruaysoongnern S, Suphanchaimant N. Land-use patterns and agricultural production systems with emphasis on changes driven by economic forces and market integration. In: Kam, S.P., Hoanh, CT., Trebuil, G. and Hardy, B. (Eds.). *Natural resource management issues in the Korat basin of northeast Thailand: An*

- Overview. International Rice Research Institute. 2001;67-77.
13. Biswas C, Singh R. Integrated farming system: An intensive approach. *Intensive Agriculture*. 2003;41(7- 8):22-30.
 14. Mali Hansram, Kumar Amit, Katara Pawan. Integrated farming system for irrigated and rainfed conditions. In: *Proceedings of National Symposium on Agricultural Diversification for Sustainable Livelihood and Environmental Security*, November 18-20, Ludhiana, India. 2014;546.
 15. Jayanthi C, Mythily S. Crop-poultry-fish-mushroom integrated farming systems for lowlands of Tamil Nadu, *Journal of Farming Systems Research and Development*. 2002;8:93-95.
 16. Swaminathan MS. Enhancing our agricultural competitiveness. *Current Science*. 2003;85:886-895.
 17. Kumar S, Jain DK. Are linkages between crops and livestock important for the sustainability of the farming system?, *Asian Economic Review*. 2005;47(1):90-101.
 18. Janzen HH, Campbell CA, Brandt SA, Lafond GP, Townley-Smith L. Light fraction organic matter in soils from long-term crop rotations. *Soil Science Society of America Journal*. 1992;56(6):1799-1806.
 19. Ding G, Novak JM, Amarasiriwardena D, Hunt PG, Xing B. Organic Matter Characteristics as Affected by Tillage Management. *Soil Sci. Soc. Am. J. Soil*. 2002;66:421-429.
 20. Schjonning P, Munkholm L, Elmholt S, Olesen JE. Organic matter and soil tilt in arable farming: Management makes a difference within 5-6 years, *Agriculture Ecosystems and Environment*. 2007;122: 157-172.
 21. Theng BKG, Tate KR, Sollins P. *Constituents of organic matter in temperate and tropical soils*. Honolulu: H. University of Hawaii Press; 1989.
 22. Mahapatra IC. Farming system research – A key to sustainable agriculture. *Fertilizer News*. 1994;39(11): 13-25.
 23. Knowler D. A review of selected models with environmental influences in fisheries. *Journal of Bioeconomics*. 2002;4:163–181.
 24. Lans T, Seuneke P, Wageningen AH, Klerkx L. *Agricultural entrepreneurship Encyclopedia of Creativity, Innovation, and Entrepreneurship*, New York; 2013.
 25. Chan GL. What does integrated farming system do? *Integrated farming system*; 2006. Available:<http://www.scizerinm.org/chanarticle.html>.
 26. Gangwar B. Farming systems research for accelerating agricultural development in less developed countries – a review. *Agricultural Reviews*. 1993;14(3):149-159
 27. Sasikala V, Tiwari R, Saravanan M. A Review on Integrated Farming Systems. *Journal of International Academic Research for Multidisciplinary*. 2015;3(7): 319-328.
 28. Tisdall JM, Oades JM. Organic matter and water stable aggregates in soils. *Journal of Soil Science*. 1982;33: 141-163.
 29. Jitsanguan T. *Sustainable Agriculture Systems for Small-Scale Farmers in Thailand*; 2001. Retrieved:<https://www.researchgate.net/publication>.
 30. Thorat BN, Thombre BM, Dadge AV. Management of dairy cow and buffalo in Integrated Farming Systems Model in Marathwada Region of Maharashtra. *International Journal of Tropical Agriculture*. 2015;33(2):653-657.
 31. Jayanthi C, Vennila C, Nalini K, Chandrasekaran B. Sustainable Integrated Management of Crop with Allied Enterprises, Tamil Nadu Agricultural University, India Tech Monitor. 2009;21-27.
 32. Okigbo BN. Major farming systems of the lowland Savanna of SSA and the potential for improvement. In: *Proceedings of the IITA/ FAO workshop*, Ibadan, Nigeria; 2009.
 33. Behera UK, Mahapatra IC. Income and employment generation for small and marginal farmers through integrated farming systems. *Indian Farm*. 1998;48 (3):16–28.
 34. Gupta V, Rai PK, Risam KS. Integrated Crop-Livestock Farming Systems: A Strategy for Resource Conservation and Environmental Sustainability. *Indian Research Journal of Extension Education, Special*. 2012;2:49-54.
 35. Preston TR, Ambio. Future strategies for livestock production in tropical third world countries, *AMBIO*. 1990;19(8):390–393.

36. Edwards P. Sustainable food production through aquaculture. *Aquaculture Asia*. 1997;2(1):4-7.
37. Hanafi H. The Role of the cage of the communal system of beef cattle, integrated agricultural waste in supporting food sovereignty in Yogyakarta. *Agros*. 2016;18(2):126- 133.
38. Kumar O, Sannathimmappa HG, Basavarajappa DN, Vijay SD, Pasha Akmal, Rajani SR. Integrated Farming System- An Approach Towards Livelihood Security resource Conservation and Sustainable Production for Small and Marginal Farmers. *International Journal of Plant and Soil Science*. 2017;15(3):1-9.
39. Govardhan M, Latheefpasha Md, Sridevi S, Kumari CP. Integrated Farming Approaches for Doubling the Income of Small and Marginal Farmers. *International Journal of Current Microbiology and Applied Sciences*. 2018;7(3):3353-3362.
40. Asai M, Moraine M, Ryschawy J, De Wit J, Hoshide AK, Martin G. Critical factors for crop-livestock integration beyond the farm level: A cross- analysis of worldwide case studies. *Land use policy*. 2018;73:184–194.
41. Rai RB, Dhama K, Chakraborty S, Damodaran T, Singh B, Ali H, Rai S, Wani MY, Ram RA. An exploratory study on using rural poultry as a source of bio-control agent for pantaloon and some vegetables. *International Journal of Current Research*. 2013;5(9):2625-2627.
42. Arene CJ. Profit function analysis of small ruminant enterprises in Nsukka local government area of Enugu state, Nigeria. *Economic affairs*. 2002;47(4):209-214.
43. Mowa E, Akundabweni L, Chimwamurombe P, Oku E, Mupambwa HA. The influence of organic manure formulated from manure on growth and yield of tomato. *African Journal of Agriculture Research*. 2017;12(41):3061 – 3067.
44. Rhodes CJ. The imperative for regenerative agriculture. *Sci Prog*. 2017;100:80–129.
45. Shankar KA, Yogesh LN, Prashant SM, Sheik PP, Desai BK. Integrated Farming System: Profitable Farming to Small Farmers. *Int. Jr. of Current Microbiology*. 2017;6(10):2819- 2824.
46. Maulana M. Social costs of System of Rice Intensification (SRI) and conventional rice production system in Indonesia. Master of Science thesis in Environmental Sciences of Wageningen University and Research Centre, Netherlands; 2015.
47. Rati M, Seharawat PS, Anil R, Jasvinder K. Sequestration of green house gases for eco-friendly agriculture. *International Journal of Agriculture Sciences*. 2016;8(26):1536 – 1539.
48. Behera UK, Babu Amjath, Kaechele H, France J. Energy self-sufficient sustainable integrated farming systems for livelihood security under a changing climate scenario in an Indian context: a case study approach. *CAB Reviews*., 2015;10(19):1–11.
49. Wulandari WA. Integration of cows with corn in sub optimal land in Bengkulu Province. Report. Institute for Agricultural Technology Studies, Bengkulu; 2014.
50. Sunanto, Nasrullah. Study of zero waste agriculture model with approach to integration of corn crops-cattle in South Sulawesi. *Proceedings of Insinas*. 2012;223-228.
51. Erlina Y, Anggreini T. Study of management of integration of corn plants and cattle in Kalampange Village, Sabangan Sub-District. *Journal of Socio Agricultural Economics*. 2014;10(1):01-08.
52. Elly FH. Impact of Transaction Costs on Household Economic Behaviour of Farmers in Cattle-Crop Farming in North Sulawesi. Doctoral Dissertation submitted to Bogor Agricultural Institute, Bogor; 2008.
53. Erniyani K, Wahyuni S, Yustina MSWP. Struktur komunitas mesofauna tanah perombak bahan organik pada vegetasi kopi dan kakao. *Agric*. 2010;3(1): 1–8.
54. Niswati A, Yusnaini S, Utomo M, Dermiyati, Arif MAS, Haryani S, Kaneko K. Long-term organic mulching and no-tillage practice increase population and biomass of earthworm in sugarcane plantation. *Earth Environ. Sci. IOP Conf. Ser*. 2015;215.
55. Yan S, Singh AN, Fu S, Liao C, Wang S, Li Y, Cui Y, Hu L. A soil fauna index for assessing soil quality. *Soil Biology Biochemistry*. 2012;47:158–165.
56. Menta C. Soil fauna diversity – function, soil degradation, biological indices, soil restoration. In: *Biodiversity Conservation and Utilization in a Diverse World*, chapter. 2012;3:59-94.
57. Grubert D, Butenschoen O, Maraun M, Scheu S. Understanding earthworm –

- Collembola interactions and their importance for ecosystem processes needs consideration of species identity. *European Journal of Soil Biology*. 2016;77:60–67
58. Edwards P, Kaewpaitoon K, McCoy EW, Chantachaeng C. Pilot small-scale crop/livestock/fish integrated farm. Asian Institute of Technology, Bangkok, Thailand, AIT Research Report No. 1986;184.
 59. FAO. The State of World Fisheries and Aquaculture, FAO: Rome, Italy; 2012.
 60. Lightfoot C, Bimbao MAP, Dalsgaard JPT, Pullin RSV. Aquaculture and sustainability through integrated resources management. *SAGE Journal*. 1993;22(3): 143–150.
 61. New MB (1991) Turn of the millennium aquaculture. Navigating troubled waters or riding the west of the wave. *World Aquaculture*, 22 (3), 28–49.
 62. Biswas S, Goswami B, Sahu NC. Fish-duck and dyke vegetable cultivation practices in rural integrated farming system. *Indian Res. J. Ext. Edu*. 2013;13(1):72-76.
 63. Delmendo MN. A review of integrated livestock-fowl-fish farming systems, Integrated agriculture-aquaculture farming systems. *In: Proceedings, ICLARM-SEARCA conf., 1979, Manila Philippines*. 1980;59–71.
 64. Brummett RE, Gockowski J, Bakwowi J, Etaba AD. Aquaculture Economics & Management, Analysis of aquaculture investments in peri urban Yaounde', Cameroon. 2004;8(5–6):319–328.
 65. Brummett RE, Gockowski J, Pouomogne V, Aboo Medjo JM, Mvilongo Mbassi D, Soua N, Teteo F. Development of integrated aquaculture –agriculture systems for small-scale farmers in the forest margins of Cameroon. Final technical report, London, UK Department for International Development; 2005. (NRE9800 605/ 522/003), 34.
 66. Karim M, Little DC, Kabir S. Md., Verdegem, MJC, Telfer T, Wahab A. Md. Aquaculture, Enhancing benefits from polycultures including tilapia (*Oreochromis niloticus*) within integrated pond-dike systems: A participatory trial with households of varying socio-economic level in rural and periurban areas of Bangladesh. 2011;23(14):225– 235.
 67. Ayinla OA. Integrated fish farming: A veritable tool for poverty alleviation/Hunger eradication in the Niger Delta Region. In: Conference Proceedings of Fisheries Society of Nigeria, Owerri, Nigeria. 2003;40-41.
 68. Prein M, Ahmed M. Integration of aquaculture into smallholder farming systems for improved food security and household nutrition. *Food and nutrition Bulletin*. 2000;21:466-471.
 69. Buck DH, Baur RJ, Rose CR. Utilization of swine manure in a polyculture of Asian and North American fishes. *Trans. Amer. Fish. Soc*. 1978;107(1):216-222.
 70. Rappaport U, Sarig S, Bamidgeh. The result of manuring on intensive growth fish farming at the Ginosar Station ponds in 1977;30(2):27-36.
 71. Jhingran AG. Integrated fish-livestock-crop farming and its role in developing rural economy. *Bull. Cent. Inland Fish. Res. Inst., Barrackpore*. 1986;(48):4.
 72. Oláh J, Pekár F, Szabó PJ. Nitrogen cycling and retention in fish-cum-livestock ponds. *Appl. Ichthyol*. 1994;10; 341–348.
 73. Rahman MM, Yakupitiyage A, Ranamukhaarachchi SL. Agricultural use of fishpond sediment for environmental amelioration. *Inter. J. Sci. Tech*. 2004;9:1–10.
 74. Phu TQ, Tinh TK. Chemical compositions of sludge from intensive striped catfish (*Pangasianodon hypophthalmus*) culture pond. *J. Sci. Can Tho Univ*. 2012;22:290–299.
 75. Rahman M, Yakupitiyage A. Use of fishpond sediment for sustainable aquaculture—Agriculture farming. *Int. J. Sustain. Dev. Plan*. 2006;1:192–202.
 76. Pouomogne V, Pemsil DE. Recommendation domains for pond aquaculture. Country case study: Development and status of freshwater aquaculture in Cameroon. *World Fish Centre Studies and Reviews No*. 2008;1871:60.
 77. Grosse O, Oswald M. The role of the farmers' group in fish innovation in an extension project's frame. *Innovation and Sustainable Development in Agriculture and Food*, Montpellier, France; 2010.
 78. Glasser F, Oswald M. High stocking densities reduce *Oreochromis niloticus* yield: model building to aid the optimization of production. *Aquatic Living Resources*. 2001;14:319 –326.

79. Hu L, Ren W, Tang J, Li N, Zhang J, Chen X. The productivity of traditional rice–fish co-culture can be increased without increasing nitrogen loss to the environment. *Agriculture Ecosystem and Environment*. 2013;177:28–34.
80. Inland Aquaculture Research Centre. *Technology Development of Eco-Culture by Using Paddy Field*, National Institute of Fisheries Sciences: Busan; 2016. Korea.
81. Poonam A, Sanjoy S, Nayak PK, Shahid MD. Optimization of rice plant density for enhancing grain yield and growth of fish under rice –fish system in coastal lowlands. In proceedings of 11th National Symposium ISCAR 'Innovations in Coastal Agriculture-Current Status and Potential under Changing Environment, January 14-17, Bhubaneswar, India. 2016;109.
82. Sinhababu DP, Jha KP, Mathur KC, Ayyappan S. Integrated farming system: rice-fish for lowland ecologies. In: *Proceedings of the International Symposium on Rainfed Rice for Sustainable Food Security*, September 23-25, Orissa, India. 1998;203-222.
83. Paul J, Modi A, Patel J. Predicting green product consumption using theory of planned behaviour and reasoned action. *Consum. Serv*. 2016;29:123–134.
84. Hirpo LA. Evaluation of integrated poultry-fish-horticulture production in Arsi Zone, Ethiopia. *International Journal of Fisheries and Aquatic Studies*. 2017;5(2):562-565.
85. Desaleng L, Aklilu S. Research experiences in Onion production, Ethiopian Agricultural Research Organization. Research report, No. 2003;55. Ethiopia.
86. Mitra K, Khan M, Mandal S, Addy R. Potentiality of integration of different components under fish based farming system for increasing farmers income. *International Journal of Agriculture Sciences*. 2018;10(13):6547-6549.
87. Xu G, Liu X, Wang Q, Yu X, Hang Y. Integrated rice – duck farming mitigates the global warming potential in rice season. *Science of the Total Environment*. 2017;575:58 – 66.
88. Mukhlis, Melinda N, Nofialdi, Mahdi. The Integrated Farming System of crop and livestock: A Review of rice and cattle Integration Farming. *International Journal of Sciences: Basic and applied Research*. 2019;42(3):68-82.
89. Sari FS, Roesdiyanto, Ismoyowati. Influence of *Azolla microphylla* and *Lemna polyrrhiza* use, peking duck feed on different protein level on carcass weight and percentage and carcass parts. *Scientific Journal Ranch*. 2013;1(3):914-923.
90. Amaludin F, Suswoyo I, Roesdiyanto. Weights, and percentage of portions rejects ducks mojosari carcass based systems and location of maintenance. *Scientific Journal Ranch*. 2013;1(3):924-932.
91. Furuno T. *The Power of Duck Integrated Rice and Duck Farming*, Australia. Tagari Publications; 2001.
92. Mukherjee TK. Integrated crop-livestock-fish production systems for maximizing productivity and economic efficiency of small holders farms. *Royal Academy of Overseas Sciences*, Brussels; 1995.
93. Ravishankar N, Pramanik SC, Rai, Shakila Nawaz RB, Tapan KR, Biswas, Nabisat Bibi. Study on integrated farming system in hilly upland areas of Bay Islands. *Indian Journal of Agronomy*. 2007;52:7-10.
94. Goverdhan M, Kumari P, Reddy GK, Sridevi S, Alibaba MD, Chiranjeevi K, Kumar MS. Evaluation of Integrated Farming System Model for Resource Recycling and Livelihood Security of Small and Marginal Farmers of Telangana State, India. *Current Journal of Applied Science and Technology*. 2020;39(34):17-26.
95. Weigend S, Romanov MN, Rath D. Methodologies to identify evaluate and conserve poultry genetic resources. In: *XXII World's Poultry Congress: Book of Abstracts*. World's Poultry Science Association, WPSA — Turkish Branch, Istanbul, Turkey. 2004;84.
96. Endabu M, Tugie D, Negisho T. Fish growth performance in ponds integrated with poultry farm and fertilized with goat manure: a case in Ethiopian Rift Valley. *International Journal of Fishery Science and Aquaculture*. 2016;3(2):40-45.
97. Sunyoto P, Rachman B. Assessment on the Crop Livestock (CLS) System in the Paddy Field of Lebak District, Banten. In: *Proceeding of the National Conference on Animal Husbandry and Veterinary Technology*; 2005.
98. Suwandi. Sustainability of Integrated Wetland Paddy-Livestock At Sragen: A Rap CLS Approach. Dissertation submitted to Bogor Agricultural University

- Postgraduate School, Bogor, Indonesia; 2005.
99. Duncan AJ, Tarawali SA, Thorne PJ, Valbuena D, Descheemaeker K, Tut SHK. Tropical Grasslands-Forrajes Tropicales, Integrated Crop-Livestock Systems—a Key to Sustainable Intensification in Africa. 2013;1:202-206.
100. Kibria ASM, Haque MM. Potentials of integrated multi-trophic aquaculture (IMTA) in freshwater ponds in Bangladesh. Aquaculture Reports. 2018;11:8–16.
101. Das A, Choudhury BU, Ramkrushna GI, Tripathi AK, Singh RK, Ngachan SV, Patel DP, Layek J, Munda GC. Multiple Use of Pond Water for Enhancing Water Productivity and Livelihood of Small and Marginal Farmers. Indian Journal of Hill Farming. 2013;26(1):29-36.
102. Sariubang M. The farming integration system of bali cattle breeding with rice plants in rice fields. Agrisistem Journal. 2010;6:36-41.
103. Basuni R. The Integration Rice-Beef Cattle on Farming Systems in Rice Fields Case Study in Cianjur Regency, West Java, Bogor Agricultural University Postgraduate School, Bogor, Indonesia; 2012.
104. Nanda PK, Bandopadhyay UK. Recent advances in integrated livestock cum fish farming in India. In: Training manual of short course on advances in production of livestock management practices. IVRI, ERS, Belgachia, Kolkata; 2011.
105. Ghosh PK, Saha R, Das A, Tripathi AK, Samuel MP, Lama TD, Mandal S, Ngachan SV. Participatory Rain water Management in Hill Ecosystem – a success story. Technical Bulletin No. 67. FPARP- Phase I, ICAR Research Complex for NEH Region, Meghalaya; 2009.
106. Govindan R, Chinnusamy KN, Chandrashekar B, Budhar MN, Prince J. Poultry- fish culture in rice farming system in Cauvery Delta region of Tamil Nadu. Indian Journal of Agronomy. 1990;35:23-29.
107. Islam MDJ, Mondal MK. Agricultural Water Management, Water management strategy for increasing monsoon rice production in Bangladesh. 1992;22(4):335-343.
108. Lal M, Patidar J, Kumar S, Patidar P. Different integrated farming system model for irrigated condition of India on basis of economic assessment: A case study: A review, International Journal of Chemical Studies. 2018;6(4):166 - 175.

© 2022 Manivannan and Thilagam; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/85109>