



# **The Influence of Agitation Rates, pH and Calcium Carbonate on L-lysine Production by *Bacillus subtilis* using Agricultural Products as Carbon and Nitrogen Sources**

**J. Okpalla <sup>a\*</sup> and I. A. Ekwealor <sup>b</sup>**

<sup>a</sup> *Department of Microbiology, Chukwuemeka Odumegwu Ojukwu University, P.M.B. 02 Uli, Anambra State, Nigeria.*

<sup>b</sup> *Department of Applied Microbiology and Brewing, Nnamdi Azikiwe University, Awka, Anambra State, Nigeria.*

## **Authors' contributions**

*This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.*

## **Article Information**

DOI: 10.9734/MRJI/2022/v32i111351

## **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here:

<https://www.sdiarticle5.com/review-history/95449>

**Original Research Article**

**Received: 20/10/2022**

**Accepted: 28/12/2022**

**Published: 29/12/2022**

## **ABSTRACT**

L-Lysine is an essential amino acid that is required in the diet of humans and animals. It is utilized in human medicine, cosmetics and pharmaceutical industry. The influence of agitation rates, pH and calcium carbonate on L-lysine production by *Bacillus subtilis* using agricultural products as carbon and nitrogen sources was studied. The L-lysine-producing bacteria had already been isolated from Nigerian soil. They were purified and Identified as *B. subtilis* PR13 and *B. subtilis* PR9, using cultural, biochemical and molecular characteristics. Optimization of some parameters which

\*Corresponding author: E-mail: judyzuby@yahoo.com;

included agitation rates, pH values and CaCO<sub>3</sub> concentrations, on L-lysine production by the *Bacillus* species was carried out. The L-lysine was produced in 250 ml flasks containing fermentation media (FM1 and FM2). The findings revealed that, enhanced L-lysine yield of 2.10 and 1.33 mg/ml was observed at agitation rate of 180 rpm for *B. subtilis* PR13 and PR9 respectively. There was a positive correlation between agitation rates and L-lysine production by *B. subtilis* PR13 and PR9 ( $r = 0.96$  and  $0.83$  respectively). The pH of 7.5, stimulated optimum L-lysine yield of 2.27 mg/ml for PR13 and 1.38 mg/ml for PR9. There was a positive correlation between pH values and L-lysine production by *B. subtilis* PR13 and PR 9 ( $r = 0.63$  and  $0.50$  respectively). The supplementation of 40g/l of CaCO<sub>3</sub>, enhanced optimum L-lysine yield of 2.18 mg/ml for *B. subtilis* PR 13 and 1.30 mg/ml for *B. subtilis* PR9. There was a positive correlation between varying concentrations of calcium carbonate and L-lysine production by the *B. subtilis* PR13 ( $r = 0.35$ ), while negative correlation was observed for *B. subtilis* PR 9 ( $r = -0.10$ ). The results obtained in the study illustrated that the optimization of process parameters could increase the L-lysine yield from agricultural products by *B. subtilis* PR13 and *B. subtilis* PR9.

**Keywords:** *Bacillus* species; L-lysine; submerged fermentation; agitation rate; pH.

## 1. INTRODUCTION

“Amino acids are molecules used by all living things to make proteins and they contain both amino and carboxylic acid functional groups. Amino acid fermentation has reached a stage where it is playing a vital role for the supply of natural amino acids at industrial level. The discovery of glutamic acid producing bacterium, *Micrococcus glutamicus* (later renamed as *Corynebacterium glutamicum*) gave a new dimension to amino acid production. This break through laid the foundation for other researchers who lately reported many bacteria involved in amino acid fermentation” [1]. “Microbial fermentation provides 100% L-amino acids, whereas by chemical method, 50% D and 50% L- amino acids are obtained” [2]. “Amino acids are major industrial products derived by fermentation, covering a world market of more than 5 million tons per year” [3]. “The most extensive formed amino acid (roughly 900,000 tons each year) occurs as L-glutamic acid, trailed by L-lysine (420,000 tons each year) as well as DL-methionine (350,000 tons each year) while the rest of the amino acids trail behind” [4].

“L-Lysine is one of the 9 amino acids which are essential for human and animal nutrition. It is not synthesized biologically in the body. L-lysine is useful as medicament, chemical agent, food material and feed additive” [5]. “L-Lysine along with some other amino acids like aspartic acid is used extensively in the pharmaceutical industry in the formulation of diets with balanced compositions and in amino acid infusion” [6,7]. “As a fine chemical, it is utilized in human medicine, in cosmetics and as precursor for industrial chemicals. L- Lysine deficiency in man

has been linked to the development of various diseases and physiological conditions including anemia, nausea, dizziness, blood shot eyes, hypersensitivity to sound, impaired growth, hair loss, disorders of reproductive system, osteoporosis, cystinuria, and immunodeficiency” [8,9].

“L-Lysine production is not restricted to any particular group of microorganisms, though the high yielding strains are mostly the species of *Arthrobacter*, *Corynebacterium* and *Brevibacterium*” [10]. “Very good producer strains have also been developed from *Bacillus subtilis* and *Escherichia coli*” [11].

“Both chemical and biochemical methods are used for L- lysine production” [5,12]. “From the commercially manufactured L-lysine, 80% is manufactured by biochemical method and only 20% by chemical means” [13]. “Among biochemical methods, fermentation is the most economical and practicable means of producing lysine” [14]. “Its major commercial form is L-lysine-HCL (L-lysine monohydrochloride). Of course, each L-lysine commercial form requires different downstream processing to achieve the degree of purity (implying significant differences in investment costs, losses during down streaming, amount of waste volume, and user friendliness), so that L-lysine-HCl involves higher operating costs due to the larger number of equipment and energy consumption. Extensive research has been made in order to improve the fermentation process not only from the point of lowering production costs but also of increasing productivity” [15,16]. “Improvements have included for example, increased yield of desired metabolites, removal of unwanted co-

metabolites, improved utilization of inexpensive carbon and nitrogen sources, or alteration of the morphology to a form better suited for separation of the organisms from the product” [17].

“As Nigeria is a developing country, a huge amount of foreign exchange is spent in the importation of L-lysine for our industries. There is huge potential in production of L-lysine locally by microbiological methods using available agricultural products. In an earlier study, we had isolated three *Bacillus* species (which included *Bacillus subtilis* PR13, *Bacillus subtilis* PR9, and *Bacillus pumilus* SS16) from Nigerian soil, which produced various yields of L-lysine” [18]. In a further study, the *Bacillus* species were used for L-lysine production using carbohydrates as carbon and seed meals as nitrogen sources [19].

The present research work was aimed at determining the influence of agitation rates, pH and calcium carbonate on L-lysine production by *Bacillus subtilis* using agricultural products as carbon and nitrogen sources.

## 2. MATERIALS AND METHODS

### 2.1 Isolation of Bacteria

The two bacteria isolates used in this study were isolated from different locations in Awka town, Anambra state, Nigerian [18]. They were purified and Identified as *B. subtilis* PR13 and *B. subtilis* PR9, using cultural, biochemical and molecular characteristics. The bacteria cultures were maintained at 4°C until used and examined for the production of L-lysine.

### 2.2 Inoculum Preparation

Two loopfuls of *B. subtilis* PR13 and PR9 were inoculated in an Erlenmeyer flask containing 50 ml of seed medium which had already been sterilized at 121 °C for 15 min. The seed medium consisted of peptone, 10.0g; yeast extract, 10.0 g; NaCl, 5.0 g; water, 1litre; pH adjusted to 7.2. The inoculated flasks were incubated for 24 h on a rotary shaker at 120 rpm and 30 °C. Duplicate flasks were used.

### 2.3 Media for Fermentation

Two different fermentation media (FM1 and FM2) were used for L-lysine production by the two *Bacillus subtilis*. A cotton plugged Erlenmeyer flasks (100ml) containing 20ml of fermentation medium (FM1) comprising of KH<sub>2</sub>PO<sub>4</sub>, 0.5g;

K<sub>2</sub>HPO<sub>4</sub>, 0.5g; MgSO<sub>4</sub>.7H<sub>2</sub>O, 0.001g; MnSO<sub>4</sub>.H<sub>2</sub>O, 0.001g; FeSO<sub>4</sub>.7HO, 0.001g; CaCO<sub>3</sub>, 50g, the carbon source (glucose) was replaced by millet starch hydrolysate 60g; the nitrogen source (ammonium sulphate) was replaced by soyabean meal 40g ; water, 1 litre; pH adjusted to 7.2 was used for *Bacillus subtilis* PR13. Another cotton plugged Erlenmeyer flasks (100) containing 20ml of fermentation medium (FM2) comprising of KH<sub>2</sub>PO<sub>4</sub>, 0.5g; K<sub>2</sub>HPO<sub>4</sub>, 0.5g; MgSO<sub>4</sub>.7H<sub>2</sub>O, 0.001g; MnSO<sub>4</sub>.H<sub>2</sub>O, 0.001g; FeSO<sub>4</sub>.7HO, 0.001g; CaCO<sub>3</sub>, 50g, the carbon source (glucose) was replaced by sorghum starch hydrolysate 60g, the nitrogen source (ammonium sulphate) was replaced by deffated peanut meal (an agricultural product) 40g; water, 1 litre; pH adjusted to 7.2 was used for *Bacillus subtilis* PR9.

## 2.4 Optimization of Culture Conditions for L-lysine Production

### 2.4.1 Effect of agitation rates

The effect of the different agitation rates on growth and lysine production by *B. subtilis* PR13 and PR9 were studied. A (100ml) Erlenmeyer flasks containing the 20 ml of fermentation media (FM1 and FM2) as was previously described was used for L-lysine production. After sterilization, the media were cooled to room temperature and 1ml (1.8×10<sup>7</sup>) volume of the cultures of *Bacillus* species (24h) was inoculated into the fermentation media. Uninoculated flasks served as control. The flasks were placed on a rotary shaker at different agitation rates (120, 140, 160, 180 and 200 rpm) and incubated at 30°C for 72h. At the end of incubation, samples of the fermentation medium were aseptically dispensed into cuvettes using micropipettes. Thereafter, the cuvettes were placed in the spectrophotometer and the reading for bacteria growth was determined at 660nm. For the determination of L-lysine and residual sugar, the fermentation medium was subjected to centrifugation at 5,000 rpm for 15 min to obtain the cell free supernatant which is the crude L- lysine. The cell free supernatant was used for the determination of lysine and residual sugar. The experiments were conducted in triplicate

### 2.4.2 Effect of pH values

The effects varying pH values on growth and L-lysine production by *B. subtilis* PR13 and PR9 were studied. A (100ml) Erlenmeyer flasks containing 20 ml fermentation media (FM1 and

FM2) as was previously described was used for lysine production. The media were prepared at different pH values of 6.5, 6.8, 7.0, 7.2, 7.5 and 8.0 and sterilized at 121°C for 15 min.

After sterilization, the media were cooled to room temperature and 1ml ( $1.8 \times 10^7$  cfu /ml) volume of the cultures of *Bacillus* species (24h) was inoculated into the fermentation media. Uninoculated flasks served as control. The flasks were placed on a rotary shaker (at 160 rpm) and incubated at 30 °C for 72 h. Thereafter, residual sugar, bacterial growth and L-lysine production were determined from the broth culture as was previously described. The experiments were conducted in triplicate.

#### 2.4.3 Effect of calcium carbonate

The effects of different concentrations of calcium carbonate on growth and L-lysine production were determined. A (100ml) Erlenmeyer flasks containing the 20 ml of fermentation media (FM1 and FM2) as was previously described was used for L-lysine production. Various concentrations of calcium carbonate (20, 30, 40, 50 and 60g/l) were added to the fermentation media and sterilized at 121°C for 15min. After sterilization, the media were cooled to room temperature and 1ml ( $1.8 \times 10^7$  cfu/ml) volume of the cultures of *Bacillus* species (24h) was inoculated into the fermentation media. Uninoculated flasks served as control. The flasks were placed on a rotary shaker (at 160 rpm) and incubated at 30°C for 72h. Thereafter, residual sugar, bacterial growth and L-lysine production were determined from the broth culture as was previously described. The experiments were conducted in triplicate

#### 2.5 Quantitative Determination of Lysine

L-lysine in the broth culture was determined by acidic ninhydrin method of Chinard [20]. A 5ml volume of the culture broth of the isolate was centrifuged at 5000 xg for 20min, and the cell-free supernatant was collected and assayed for lysine production. 1ml of glacial acetic acid was added to 1ml of supernatant in a test tube. Thereafter, one ml of a reagent solution which contains an acid mixture, 0.4ml of 6M orthophosphoric acid, 0.6ml of glacial acetic acid and 25mg of ninhydrin, was also added to the supernatant in the test tube. The blank contains 1ml of glacial acetic acid, 1ml of the acid mixture without ninhydrin and 1ml supernatant. Both tubes were capped and the contents mixed properly for 10min before heating at 100°C in a

water bath for 1h. The test tubes were cooled rapidly under tap water and 2ml of glacial acetic acid was added to each test tube to give a final volume of 5ml. The optical density of the reacting mixture was read against the blank at 515nm in a spectrophotometer. Results obtained with the test samples were extrapolated from a standard lysine curve.

#### 2.6 Estimation of Reducing Sugar

The reducing sugar content was determined by dinitrosalicylic acid (DNS) method of Miller [21]. Reducing sugar was estimated by adding 1ml of DNS to 1ml of the supernatant. The mixture was heated in a water bath at 100 °C for 10min and allowed to cool. The volume of the mixture was thereafter increased to 12 ml with distilled water. After allowing the reaction mixture to stand for 15min at room temperature, the optical density was measured at 540 nm in a spectrophotometer against a blank prepared by substituting the supernatant with water. The reducing sugar content was subsequently determined by making reference to a standard curve of known glucose concentrations.

#### 2.7 Statistical Analysis

Data generated from this work were analyzed using correlation analysis with a software application SPSS version 14

### 3. RESULTS

The effect of agitation rates on growth and lysine production by *B. subtilis* PR13 and PR 9 are presented in Figs. 1-2. The results showed that maximum growth and lysine yields by *Bacillus subtilis* PR13 and *Bacillus subtilis* PR9, were observed at agitation rate of 180 rpm. The highest lysine accumulation of 2.10 and 1.33 mg/ml by *B. subtilis* PR13 and PR9, corresponded with a residual sugar of 0.60 and 0.53 mg/ml respectively. There was a positive correlation between agitation rates and lysine production by *B. subtilis* PR13 and PR9 ( $r = 0.96$  and  $0.83$  respectively).

The result of the effect of pH on growth and lysine production by *B. subtilis* PR13 and *B. subtilis* PR 9 are presented in Figs. 3-4. The results show that maximum growth and lysine yields by the *Bacillus* strains, were observed at a pH 7.5. The highest lysine accumulation of 2.27 and 1.38 mg/ml by *B. PR13* and *PR9*

respectively, corresponded with a residual sugar of 0.45 and 0.51 mg/ml respectively. There was a positive correlation between pH values and lysine production by *B. subtilis* PR13 and PR 9 ( $r = 0.63$  and  $0.50$  respectively).

The effect of calcium carbonate on growth and lysine production by *B. subtilis* PR13 and PR 9 are presented in Figs. 5-6. The results showed that maximum growth and lysine yields by all the *Bacillus* species, were observed when 40g/l of  $\text{CaCO}_3$  was added. The highest lysine accumulation of 2.18 and 1.30mg/ml by *B.subtilis* PR13 and PR9 respectively, corresponded with residual sugars of 0.33 and 0.50 mg/ml respectively. There was a positive correlation between varying concentrations of calcium carbonate and lysine production by the *B. subtilis* PR13 ( $r = 0.35$ ), while there was negative correlation for *B. subtilis* PR 9 ( $r = -0.10$ ).

#### 4. DISCUSSION

The study revealed that maximum L-lysine yield by *B. subtilis* PR13 and *B. subtilis* PR9 was observed at agitation rate of 180rpm, which later decreased. This was contrary to the findings of Shah et al. [22] who studied “the effect of agitation on lysine production by *Corynebacterium glutamicum*. They observed

that L- lysine production increased from 50rpm to the maximum at 200rpm”. Also, Wang et al. [23] worked “at 200 rpm for fermentation of L-lysine on a rotator shaker”. Rehman et al. [24] reported that “L-lysine production increased from 12.5 mg/mL to 15.1 mg/mL in shake flask cultures when agitation rate was increased from 150 to 300 rpm”. Siripoke et al. [25] reported “an increase in L- lysine production when the agitation rates were increased from 300 to 400 rpm, however L-lysine yield decreased when the agitation rate was increased to 450rpm”. They opined that the decrease in L-lysine production may be due to the increasing foam observed which has resulted in the reduction of the oxygen transfer rate to the cells. Lee et al. [26] suggested that “agitation rates above 200 rpm will lead to denaturation of enzymes with attendant low production of metabolites”. Darah and Ibrahim [27] pointed out that “excessive agitation could lead to cell lysis and increase cell permeability due to abrasion by shear forces”. Shafee et al. [28] pointed out that “mixing is especially important because oxygen is a very low solubility nutrient. The oxygen transfer capability of flask can limit the amount of biomass that can be grown in the flask. When premature oxygen limitations are imposed on growth, changes in physiological state occur which results in reduced inoculum effectiveness”.




Fig. 1. Effect of agitation rates on lysine production by *Bacillus subtilis* PR13




Fig. 2. Effect of agitation rates on lysine production by *Bacillus subtilis* PR9




Fig. 3. Effect of pH values on lysine production by *Bacillus subtilis* PR13




Fig. 4. Effect of pH values on lysine production by *Bacillus subtilis* PR9





Fig. 5. Effect of calcium carbonate on lysine production by *Bacillus subtilis* PR13



**Fig. 6. Effect of calcium carbonate on lysine production by *Bacillus subtilis* PR9**

Oxygen is essential for growth and yields of metabolites in aerobic organism. For the oxygen to be absorbed it must be dissolved in aqueous solution along with the nutrients by agitating with rotary shaker. Agitation is done to provide homogeneity. Agitation intensity influences the mixing and oxygen transfer rate in many bacterial fermentations thereby influencing growth and product formation. "Under aerobic conditions, sufficient oxaloacetate is available for aspartate conversion and then flows towards L- lysine formation. However, under anaerobic conditions, carbon metabolism shifts towards formation of ethanol, lactic acid and succinate" Wendisch et al. [29].

The result of the study showed that maximum L-lysine yield was observed at pH 7.5 for both *B. subtilis* PR13 and *B. subtilis* PR9. This is contrary with the report of Siripoke et al. [25] who reported maximum lysine production of 6.73g/l for *Bacillus* SWU41, at a pH 6.0, unlike most lysine producing strains which prefer pH 7.0–7.5. [30] reported maximum lysine yield of 69.5g/l at pH 7.0 using *Brevibacterium flavum*, above pH 7.5 the sugar utilization and L-lysine production

slightly decreased. Hua et al. [31] in a study of external pH on lysine production by *Corynebacterium glutamicum* observed that the optimum pH for lysine flux was found to be 7.0. Liu [32] found that L-lysine synthesis decreased at pH lower than 6.5, but found no difference between 6.5 and 8.0. An initial pH 7.5 was recommended by L-lysine production by *Corynebacterium glutamicum* Broer and Kramer [33]. Rehman et al. [24] observed that at pH value of 7.6, optimum production of L-lysine (12.5 mg/mL) in basal medium was in agreement with the results of Broer and Kramer [33] and Hua et al [31]. Broer et al. [34] found that the optimum pH for maximum velocity of transport by *Corynebacterium glutamicum* was 7.4 – 7.8. Kelle et al. [35] studied "the effect of pH on L-lysine transport by *Corynebacterium glutamicum* and found that pH value governed the transport activity and the specific L-lysine export rate. Membrane potential was found to be an important factor influenced by pH of fermentation medium towards lysine excretion by the cells".

The results from the study revealed that 40 g/l of CaCO<sub>3</sub> was optimum for lysine production by the



*Bacillus* species. This is in contrast to with the report of Shah et al. [22] that at 2% CaCO<sub>3</sub> *Corneybacterium glutamicum* produced maximum yield of L-lysine (16.58g/l). Ekwealor and Obeta [36], Rao et al. [37] and Siripoke et al. [25] incorporated different concentrations of CaCO<sub>3</sub> in the fermentation medium during the production of lysine by *Bacillus megaterium*, *Corneybacterium glutamicum* and *Bacillus* SWU 41 respectively. Young and Chipley [38] reported that the addition of CaCO<sub>3</sub> with yeast extract resulted in a relatively constant pH and increased depletion of glucose, but reduced lysine production. "Decrease in pH could be due to accumulation of acids such as pyruvic acid, lactic acid, gluconic acid as a result, bacterial growth ceased with concomitant decrease in the product yield. pH is one of the most important factors affecting microbial propagation" [1]. "As nutrients are consumed and converted into product during the fermentation process, the pH changes markedly in the absence of suitable control mechanism. In order to maintain optimal pH, reagents like Calcium carbonate must be added to the culture medium at the beginning of the fermentation" [1]. Calcium carbonate neutralized the pH of the broth, acting as internal neutralizing or buffering agent, it would be useful for shortening the fermentation time. Wang et al. [39] reported that though the pH of the fermenter was automatically controlled by ammonia water, but still small amount of CaCO<sub>3</sub> must be added. It eliminates the lag phase of cell growth, thereby shortening fermentation time. The free hydrogen ions are consumed resulting in decreased hydrogen ion activity. In other words calcium carbonate acts to neutralize or buffer the solution by consuming hydrogen ions.

## 5. CONCLUSION

The study revealed that some agricultural products could be used for L-lysine production. It also revealed that there was improved L-lysine production by *B. subtilis* PR13 and *B. subtilis* PR9 during the optimization study which included, agitation rates, pH and calcium carbonate. The agitation rate of 180 rpm, pH 7.5 and calcium carbonate 40g/l were optimum for L-lysine production. However, the highest L-lysine yield was observed with *Bacillus subtilis* PR13. The *Bacillus* species (*Bacillus subtilis* PR 13 and PR9) have shown potential for lysine production using readily available agricultural products. This development indicates that they can be used for large scale production of the L-lysine, to meet present-day needs in the Nigeria's industrial

sector. The microbiological process of L-lysine production, if well developed could lead to the availability of the product in Nigeria and this to some extent will reduce the importation of the product into the country. Further research is needed to study the effect of other parameters on L- lysine production.

## ACKNOWLEDGEMENT

The authors are very grateful to Marian Figge of the Netherlands culture collection of bacteria (NCCB) for her assistance in the purchase of the *E. coli* lysine auxotroph used for this work. Special appreciation goes to Prof. I.A. Ekwealor, for supervising the research. Also, I want to thank Dr. Paul Egwim for facilitating the procurement of s-2-amino-ethyl-cysteine (Sigma) used in the research from the United States.

## COMPETING INTERESTS

Authors have declared that no competing interest exist

## REFERENCES

1. Kinoshita S. Glutamic acid producing microorganisms. In: M.C. Flickinger and S.W. Drew(Eds), Encyclopaedia of Bioprocess Technology: Fermentation, Biocatalysis and Bioseparation, John Wiley and Sons, New York. 1999;1330-36.
2. Khan S H, Rasool G, Nadeem S. Bioconversion of cane molasses into amino acids. Pakistan J Agr. Sci. 2006; 43(3- 4):156- 161.
3. Eggeling L, Bott M. A giant market and a powerful metabolism: L-lysine provided by *Corynebacterium glutamicum*. Appl Microbiol Biotechnol. 2015;99(8):3387-3394.
4. Tryfona T, Bustard MT. Fermentative production of lysine by *Corynebacterium glutamicum*:transmembrane transport and metabolic flux analysis. Process Biochem. 2005;40(2):499–508.
5. Anastasiadis S. L-Lysine fermentation. Recent Pat Biotechnol. 2007;1:11-24.
6. Hirose Y, Okada H. Microbial production of amino acids. In:H.J. Pepller and D. Perlman (Eds.), Microbial Technology. 1979;1:211–40. Academic Press.
7. Costa-Ferreira M, Duarte JC. Amino acid accumulation by an analogue-sensitive

- mutant of *Corynebacterium glutamicum*. Biotechnol Lett. 1992;13:721-6.
8. Meister A. Amino acid antagonists. In Biochemistry of the amino acids. Academic Press, London. 1965;231.
  9. Chen C, Sander JE, Dale NM. The effect of dietary lysine deficiency on the immune response to Newcastle disease vaccination in chickens. Avian Diseases. 2003;47 (4):1346-51.
  10. Sen SK, Chatterjee SP. Influence of B-vitamins and trace elements on lysine production by *Micrococcus varians* 2fa, Acta Biotechnol. 1989;9:63-67.
  11. Leuchtenberger W. Amino acids – technical production and uses. In Rehm H.J., Reed, G., Phuler, A., Stadler, P., editors. Products of primary metabolism. Biotechnology. Wiley- VCH, Weinheim, German. 1996;6:492.
  12. Nelofer R, Syed Q, Nadeem M. Statistical optimization of process variable for L-Lysine production by *Corynebacterium glutamicum*. Turk. J. Biochem. 2008;33(2):50- 57.
  13. Malothu R, Dowlathabad MR, Balanarasaiah T. Bioprocess Technology Development and Convalescing Production of L-Lysine in *Corynebacterium glutamicum*. International Conference on Chemical Engineering and its Applications, Bangkok (Thailand); 2012.
  14. Ekwealor IA, Obeta JAN. Studies on lysine production by *Bacillus megaterium*. Afri J Biotechnol. 2005;4(7):633-638.
  15. Wittmann C, Becker J. The L-Lysine Story:From Methabolic Pathways to Industrial Production. Microbiol Monogr. 2007;5:40-62.
  16. Kelle R, Hermann T, Bathe B. L–Lysine production. In:Eggeling L, Bott M (eds) Handbook of *Corynebacterium glutamicum*. CRC press, Boca Raton, USA. 2005;465-488.
  17. Nielsen J. Metabolic engineering. Appl Microbiol Biotechnol. 2001;55:263-283.
  18. Okpalla J. and Ekwealor IA. Screening for lysine production by bacteria isolated from Nigerian soil. World Wide J Multidiscip Res Develop. 2019a;5(4):10-18.
  19. Okpalla J. and Ekwealor IA. Studies on lysine accumulation in broth culture of *Bacillus* species using carbohydrates as carbon sources and seed meals as nitrogen sources. Inter J Trend in Scientific Res Development. 2019b;3(2):760- 772.
  20. Chinard FP. Photometric estimation of proline and ornithine. J Biol Chem. 1952;199:91-95.
  21. Miller GL . Use of dinitrosalicylic acid reagent for detection of reducing sugar. *Annals of Chem*. 1959;31:427-431.
  22. Shah AH, Hameed A, Guj Madid K. Improved microbial production of lysine by developing a new Auxotrophic mutant of *Corneybacterium glutamicum*. Pak J Biol Sci. 2002;5(1):80-83.
  23. Wang JS, Yung CK, Chen WL and Liu YT. Optimization of culture conditions for L-lysine fermentation by *Brevibacterium* species Report of Taiwan Sugar Res. Instit. 1991;134:37-48.
  24. Rehman HU, Hameed A, Ahmed S.. Selection and characterization of a lysine yielding mutant of *Corynebacterium glutamicum* - A soil isolate from Pakistan. Pak Vet J. 2012;32(1):20-24.
  25. Siripoke S, Potivejkul K, Lerstaveesin P, Leungsakul S. Optimal conditions for lysine production by *Bacillus* SWU 41 in a Lab-Scale fermenter. 31<sup>st</sup> Congress on Science and Technology of Thailand at Suranaree University of Technology; 2005.
  26. Lee I, Lee K, Namgoong K, Lee YS. The use of ion exlusion chromatography to achieve a more efficient lysine recovery from fermentation broth. Enzyme Microbiol Technol. 2002;30:798-803.
  27. Darah I, Ibrahim CO. Effect of agitation on production of lignin-degrading enzymes by *Phanerochaete chrysosporium* grown in shake-flask cultures. Asian Pacific J. Mol Biol Biotechnol. 1996;4:174-182.
  28. Shafee N, Aris SN, Rahman RNZA, Basri M, Salleh AB. Optimization of environmental and nutritional conditions for the production of alkaline protease by a newly isolated bacterium *Bacillus cereus* strain 246. J Appl Sci Res. 2005;1:1-8.
  29. Wendisch VF, Bott M, Eikmann BJ. Metabolic engineering of *Escherichia coli* and *Corynebacterium glutamicum* for biotechnological production of organic acids and amino acids. Curr. Opinion in Microbiol. 2006;9:268-274.
  30. Javed A, Jamil A, Rezaei-Zarchi S. Optimization and hyper-expressed production of lysine through chemical mutagenesis of *Brevibacterium flavum* by N-nitroso-N-ethylurea. Afri J Microbiol Res. 2011;5(29):5230-5238.
  31. Hua Q, Yang C, Shimizu K. Metabolic control analysis for lysine synthesis using

- Corynebacterium glutamicum* and experimental verification. J Biosci Bioengine. 2000;90(2):184-192.
32. Liu YT. Studies on the fermentation production of L-lysine. Optimization of culture condition for L-lysine fermentation with cane molasses. Report Taiwan and Sugar Research Institute, 1986;114:45-66.
33. Broer S, Kramer R. Lysine excretion by *Corynebacterium glutamicum* 2. Energetics and mechanism of transport system. European Journal of Bacteriol. 1991;202: 137–143.
34. Broer S, Eggeling L, Kramer R. Strains of *Corynebacterium glutamicum* differing in lysine productivity possess different lysine excretion system. Appl Environ Microbiol. 1993;59:316-321.
35. Kelle R, Lanfer B, Brunzema C, Weuster BD, Kramer R, Wandrey C. Reaction engineering analysis of L-lysine transport by *Corynebacterium glutamicum*. Biotechnol Bioengine. 1996;51:40-50.
36. Ekwealor IA, Obeta JAN. Effect of vitamins and bivalent metals on lysine yield in *Bacillus megaterium*. Afri J Biotechnol.2007;6(11):1348-1351.
37. Rao BS, Muralidararao T, Swamy AN. Studies on continuous production kinetics of L-lysine by immobilized *Corynebacterium glutamicum* 13032. Middle-East J Sci Res. 2011;7(2):235-240.
38. Young TK, Chipley JR. Microbial production of lysine and threonine from whey permeate. Appl Environ Microbiol. 1983;45:610–615.
39. Wang JS, Yung CK, Chen WL, Liu YT. Optimization of culture conditions for L-lysine fermentation by *Brevibacterium* species. Report of Taiwan Sugar Res Institut. 1991;134:37-48.

© 2022 Okpalla and Ekwealor; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/95449>