



Intestinal Obstruction Caused by Small Bowel Volvulus

Mounir Bouali ^a, Kabira Falousse ^{a*},
Abderrahmane Lamnaouar ^a, Abdelilah El-Bakouri ^a,
Khalid El-Hattabi ^a, Fatima Zahra Bensardi ^a
and Abdelaziz Fadil ^a

^a *Department of Visceral Surgical Emergency, Faculty of Medicine and Pharmacy, University Hospital Center Ibn Rochd, Hassan II University, Casablanca, Morocco.*

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/96408>

Case Study

Received: 28/10/2022

Accepted: 30/12/2022

Published: 06/01/2023

ABSTRACT

Small bowel volvulus is a surgical emergency, the etiology may be primary or secondary, and the clinical presentation is that of an acute intestinal obstruction. Abdominal radiography and abdominal CT scan is the key of the diagnostic. We present the case of a small bowel volvulus admitted to the emergency department with an acute bowel obstruction, the diagnosis was confirmed by abdominopelvic CT scan and the surgical procedure consists in of ileocecal resection with ileocolostomy, with good postoperative results.

Keywords: Intestinal obstruction; small bowel volvulus; diagnosis; surgery.

1. INTRODUCTION

“Small bowel volvulus is a rare cause of intestinal obstruction in adult patients” [1]. “It is a twisting

of the small bowel around its mesenteric axis. It is a well-known condition in infants and children, but is rare in adults. Diagnosis is difficult because the symptoms resemble those of an acute

*Corresponding author: E-mail: falousse.kabira@gmail.com;

abdomen” [2]. “Prompt diagnosis and treatment are necessary to avoid intestinal necrosis” [3].

2. CASE PRESENTATION

The patient was 44 years old and was admitted to the emergency room for an occlusive syndrome that had been evolving for 7 days. On examination the patient was hemodynamically and respiratorily stable with a temperature of 37.7°C. Abdominal examination showed a distended and tympanic abdomen with generalized abdominal defense, the hernial orifices were free. A rectal examination showed an empty rectal ampulla without palpable mass. Abdominal radiography showed gaseous hydroaeric levels. Abdominal CT scan showed small bowel obstruction with moderate peritoneal effusion (Fig 1(A-B)).

After optimization of his general condition with a Naso-Gastric tube suction and intravenous fluids resuscitation, a decision was taken to proceed with an emergency laparotomy, under general anesthesia with endotracheal intubation. Preoperative prophylactic antibiotics were administered.

Intraoperative findings (Fig. 2,3) a peritoneal effusion of medium abundance with the presence of a small bowel volvulus on incomplete common mesentery with 2 counterclockwise spiral turns with small bowel necrosis from 1m extended from the ileocaecal junction to 2m40 from the duodenojejunal angle with perforation of distal ileal. The operation consisted of an ileocaecal resection taking 1m of small bowel and a double-barrel ileo-colostomy,



(A)



(B)

Fig. 1(A-B). CT scan of the abdomen showing the twisting of the ileum around its mesentery

The postoperative follow-up was simple with resumption of transit on the 2nd day and the patient was declared discharged on the 4th day with heparin therapy. A restoration of continuity was performed 4 weeks after, which took place without complications.

3. DISCUSSION

“Small bowel volvulus is torsion of the small bowel around its mesenteric axis, resulting in mechanical obstruction of the small bowel (1-4-5), Mesenteric torsion also results in occlusion of the mesenteric vessels with intestinal ischemia and ultimately necrosis. It is a rare surgical emergency. It represents 1% of the etiologies of intestinal obstructions” (1-4-6-7). “Depending on its etiology; small bowel volvulus can be divided into two categories: primary and secondary. Primary volvulus occurs in abdominal cavities in which there are no predisposing anatomical abnormalities. On the other hand, secondary small bowel volvulus occurs in the presence of predisposing lesions, congenital or acquired, such as malrotations, flanges and adhesions” [1].

“The clinical presentation of small bowel volvulus is non specific. [4-8], the diagnosis is that of an acute intestinal obstruction [4] marked by abdominal pain, vomiting and cessation of fluid and gas”. “Physical examination shows abdominal meteorism with tympany, and pelvic touches are painless. This may lead to a delay in

diagnosis. Mortality of small bowel volvulus ranges from 9% to 35%, but increases to 20% to 100% with intestinal necrosis” [2].

The unprepared abdomen is the first-line examination for acute intestinal obstruction, and can show hydro-aerosic levels of the small bowel [1-8].The barium enema can be useful in some cases, showing the Corskrew sign [1]. “Abdominal CT is the gold standard for the positive, topographical and severity diagnosis of small bowel volvulus [8]; it is rapid, non-invasive and has a sensitivity ranging from 89 to 100%”. The "swirl" sign appears to be pathognomonic for the majority of authors, (1-4-7) contrast injection allows visualization of the verticalization, or inversion, of the superior mesenteric vessels, with a vein lying above or to the left of the artery [8].The "beak" sign is also another sign on CT showing the tapering of three dilated intestinal segments, resembling a bird's bent beak [3].

Surgery is the mainstay of management of primary small bowel volvulus [3], and it must be prompt to avoid intestinal gangrene and thus reduce morbidity and mortality [4]. The management of primary volvulus without intestinal necrosis is based on either simple devolvulation with fixation, or to gallbladder resection in all cases of volvulus, whether necrotic or not. Treatment of secondary volvulus is based on cure of the underlying cause, which will guide subsequent management [3-9].



Fig. 2. Intraoperative photograph showing volvulus of ileum



Fig. 3. Distal iléal perforation

4. CONCLUSION

Primary SBV is an extremely rare situation [3]. The clinical presentation is not specific. The diagnosis can be made by CT scan and must be made early to avoid unnecessary bowel resections [4]. Treatment often involves urgent surgical exploration.

CONSENT

As per international standard or university standard, patient(s) written consent has been collected and preserved by the author (s).

ETHICAL APPROVAL

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Hayakawa M, Makino Y, Yajima D, Inokuchi G, Motomura A, Chiba F, et al. A fatal case of adult small bowel volvulus. *Leg Med (Tokyo)*. 2014;16(5):300-2.
2. Islam S, Hosein D, Dan D, Naraynsingh V. Volvulus of ileum: A rare cause of

- small bowel obstruction. *BMJ Case Rep*. 2016;2016:bcr2016216159.
3. Santín-Rivero J, Núñez-García E, Aguirre-García M, Hagerman-Ruiz-Galindo G, de la Vega-González F, Moctezuma-Velasco CR. Volvulo of the intestine delved. Case reports and bibliographic reviews. *Cirugía y Cirujanos*. 2015;83(6):522-6.
4. Bouassida M, Beji H, Chtourou MF, Ben Othmane N, Hamzaoui L, Touinsi H. Primary small bowel volvulus: A case report and literature review. *Ann Med Surg (Lond)*. 2022;80:104250.
5. Masson E. Le volvulus total du grêle sur malrotation chez l'adulte. About 11 cases [Internet]. To install EM-Consulte. Available at: <https://www.em-consulte.com/article/1079363/resume/le-volvulus-total-du-grele-sur-malrotation-chez-l-> Access on 5 February 2023
6. Kapadia MR. Volvulus of the small bowel and colon. *Clin Colon Rectal Surg*. 2017;30(1):40-5.
7. Small bowel volvulus after laparoscopic appendectomy. *Journal of Gastroenterology de México* [Internet]. Available: <http://www.revistagastroenterologiamexico.org/en-small-bowel-volvulus-after-laparoscopic-articulo-S2255534X21000840> Access on 5 February 2023

8. Alaoui A, et al. Imagery of volvulus of the grêle on the mesentery community incomplete in an adult: case report. *Pan Afr Med J.* 2020;37:287.
9. Garel C, Blouet M, Belloy F, Petit T, Pelage JP. Diagnosis of pediatric gastric, small-bowel and colonic volvulus. *Pediatr Radiol.* 2016;46(1):130-8.

© 2023 Bouali et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/96408>