



Prevalence of *Candida* Species in 'Thairu' a Traditional Fermented Milk of Kerala, India

Archana Chandran ^{a*}, C. K. Linsha ^a and A. K. Beena ^b

^a Department of Dairy Microbiology, College of Dairy Science and Technology, Pookode, Wayanad, India.

^b Department of Dairy Microbiology, Verghese Kurien Institute of Dairy and Food Technology, Mannuthy, Thrissur, India.

Authors' contributions

This work was carried out in collaboration among all authors. Authors AC and CKL performed the research. Author AC designed and prepared the original draft manuscript. Authors AKB supervised the study and provided the necessary guidance during the work and reviewed the manuscript. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/EJNFS/2022/v14i111261

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/92351>

Short Research Article

Received 29 July 2022
Accepted 02 October 2022
Published 11 October 2022

ABSTRACT

Aims: To determine the prevalence of *Candida* species in 'thairu', a traditional fermented milk prepared in the households of Kerala.

Study Design: 'Thairu' samples collected randomly from the households in Wayanad, Kozhikode, Malappuram, Palakkad and Kannur districts of Kerala. This was followed by enumeration and identification of yeast using API 20CAUX Kit. Determination of pH and acidity of samples.

Place and Duration of Study: Department of Dairy Microbiology, College of Dairy Science and Technology, Pookode, Wayanad, Kerala. September 2021- December 2021.

Methodology: A total of 30 household 'thairu' samples were collected from households of Wayanad, Kozhikode, Malappuram, Palakkad and Kannur districts of Kerala in sterile bottles. The samples were serially diluted in normal saline and pour plated on to Yeast Extract Glucose Chloramphenicol Agar for isolation and enumeration. The pH was measured using pH meter by directly inserting the probe into the homogenized sample. Titratable acidity in per cent lactic acid was measured using N/10 NaOH. The isolates were identified based on the sugar fermentation pattern using API 20C AUX kit (Biomerieux, France).

Results: The yeast count in the samples ranged from 5.0 to 6.7 log CFU/g with an average of

*Corresponding author: Email: archanac@kvasu.ac.in;

5.89±0.38 log CFU/g. Average acidity and pH of the samples were 1.92±0.34 % LA and 3.59±0.60 respectively. A total of 23 yeast isolates were identified based on morphology and carbohydrate fermentation pattern using API 20C AUX kit (Biomerieux, France). Exactly 86.95 per cent of isolates belong to *Candida* species.

Conclusion: Observations of the study revealed the high prevalence of *Candida* species in traditional fermented milk 'thairu'. Predominant *Candida* species were lactose fermenters, but the presence of a few species with spoilage and pathogenicity potential were also detected. The safety assessment of *Candida* species is essential before applying them as starter cultures for food fermentations.

Keywords: *Candida*; fermented milk; yeast; starter culture; traditional food; safety.

1. INTRODUCTION

'Thairu' is the traditional fermented milk product which serves as the key ingredient in the preparation of Kerala cuisines like 'pulissery', 'kalan' and 'pachadi'. The product is prepared in households by back slopping usually in cow milk. The microflora of this product varies from region to region within the state itself owing to the changes in culinary practices, incubation temperature and environmental conditions. Lactic acid bacteria, yeast and mold and enterococci play a significant role in fermentation of traditional milks [1]. The presence of lactic acid bacteria and enterococci in traditional fermented milks of Kerala have been previously reported [2,3]. The presence of yeast and molds are considered as indicators of poor hygiene and are thus regarded as undesirable in fermented milks like dahi and yoghurt. Yeasts can tolerate extreme conditions like low pH, low water activity and high salt concentrations, which helps them to proliferate in fermented foods and cause spoilage. But in traditional fermented foods like kefir and kumiss the role of yeast in development of technological and sensory attributes is well studied [4]. The ability of yeast to grow in milk is mainly dependent on the assimilation of protein, fat, lactose and citrate present in milk. The effect of reduction in pH of milk during fermentation is very less for yeast when compared to lactic acid bacteria [5].

Candida is a frequently isolated species from cocoa, cheese, meat, vegetables and spontaneously fermented milks. *Candida* is also used as co-culture along with *Saccharomyces* and *Lactobacillus* to produce fermented food products [6]. At the same time *Candida* genus, especially *Candida albicans* is widely known to cause infections and hence may represent a potential risk to public health [7]. The lack of good manufacturing practices like proper pasteurization of milk and use of defined starter

cultures can result in contamination of product with pathogens. The yeast communities in fermented milk products are generally linked to the production environment. The containers used for milk fermentation play an important role in the presence of yeasts in fermented milks and yoghurts [8]. The acidophilic character of yeast along with low sensitivity to the antagonistic activity of lactic acid bacteria can support the growth of yeast in fermented milks. Hence the study on the prevalence of *Candida* species in traditionally fermented milks are important from a technological as well as safety point of view. At the moment, there seems to be paucity of information on the prevalence of *Candida* species in 'thairu', a traditional fermented milk prepared in the households of Kerala hence the need to conduct more studies and bridge this information gap.

2. MATERIALS AND METHODS

2.1 Sample Collection

Thirty samples of freshly prepared 'Thairu' were collected aseptically in sterile bottle from different households in Kerala. The samples were stored at 4°C during transportation and analysed within 12 h of sampling.

2.2 Enumeration and Isolation of Yeast

The samples were serially diluted in normal saline and pour plated on to Yeast Extract Glucose Chloramphenicol Agar (YGCA) (ISO and IDF 2004). The plates were incubated at 25°C for 2-5 days and counted. Colonies with distinct colour, size and morphology were picked and stained using methylene blue. The colonies with yeast morphology were purified by streaking in the Malt Extract Agar (MEA). The purified isolates were stored at 4°C in MEA slants and in 70% glycerol stock at -20°C.

2.3 Determination of Acidity and pH of Samples

Ten grams of curd was mixed with 10 ml distilled water and titrated against N/10 NaOH using phenolphthalein indicator till the attainment of pale pink colour. pH was determined using pH meter by directly inserting the probe into the homogenized sample.

2.4 Identification of Yeast

The selected yeast isolates were identified based on the carbohydrate fermentation pattern using API20C AUX Test Kit (Biomerieux, France), according to manufacturer's instructions. The strips had 20 cupules containing dehydrated substrates which enable the performance of 19 assimilation tests. The ability of yeast to utilize the substrate as sole carbon source was evaluated. The reactions were compared with the control cupule and identified based on the numeric profile in apiweb™ identification software.

3. RESULTS AND DISCUSSION

The yeast count in the samples ranged from 5.0 to 6.7 log CFU/g with an average of 5.89±0.38 log CFU/g. Average acidity and pH of the samples were 1.92±0.34 % LA and 3.59±0.60 respectively. A total of 23 yeast isolates were identified based on morphology and carbohydrate fermentation pattern using API 20C AUX kit (Biomerieux, France). The isolates belonged to *Candida kefir* (8 isolates), *Candida spherica* (3 isolates), *Candida parapsilosis* (3 isolates), *Rhodotorula mucilogenosa* (3 isolates), *Candida rugosa* (1 isolate), *Candida colliculosa* (1 isolate), *Candida lusitaniae* (1 isolate), *Candida zeylanoides* (1 isolate), *Candida krusei* (1 isolate) and *Candida famata* (1 isolate) (Table 1).

Among these *Candida kefir*, *Candida spherica*, *Candida krusei* and *Candida famata* were able to ferment lactose. Majority of isolates were able to ferment galactose.

The traditional fermented milk 'thairu' was found to have comparatively higher acidity and low pH. The presence of yeast has been reported in an Italian fermented milk 'Gioddu' a product with pH in the range of 3.8-4.2. The strong acidity of that product was attributed to the presence of yeast along with lactic acid bacteria in the product [9]. The occurrence of yeast in traditional fermented

milks prepared under household conditions need to be thoroughly investigated as there is high microbial risk associated with some of these floras. In this study out of the 23 isolates 20 belong to *Candida* species showing a prevalence of 86.95 per cent. Among these the predominant species were *Candida kefir/Kluyveromyces marxianus* and *Candida spherica/Kluyveromyces lactis*. Both of these yeasts can assimilate lactose and metabolise citrate to ethanol, glycerol, lactic acid and propionic acid in milk [10]. *Candida krusei/Pichia kudriavzevii* is a potential probiotic yeast isolated from fermented food products [11]. These species are reported to be acid-resistant, thermotolerant and do not produce any extracellular toxins and can be used in the synthesis of single cell protein [12]. The presence of *C.lusitanae* and *C.colliculora* have been previously reported in Zimbabwean fermented milk [4]. *Candida rugosa* and *Rhodotorula mucilogenosa* are usually associated with milk fat causing spoilage in milk and milk products [13]. Both of these species were also detected in this study. Many of these yeasts are capable of causing spoilage in dairy products. In plankton form non *Candida-albicans* species like *C.krusei*, *C.lusitaniae*, *C.famata*, *C.parapsilosis* and *C.tropicalis* can produce biotypes and virulence factors similar to *C.albicans* showing the risk of their delivery through fermented foods. But the contribution of yeast in enhancement of flavour and other technological properties in traditional fermented milks like kefir, kumiss and fermented tea Kombucha have been well studied [14]. Thorough animal model studies have to be carried out to prove the invasiveness of these isolates before recommending them as starter cultures [15].

Table 1. Identification of yeast isolates by sugar fermentation pattern using API 20C AUX Kit

| Isolate No. | Identified as apiweb™ |
|---|------------------------|
| Y01 | <i>C. krusei</i> |
| Y06, Y10, Y13, Y14, Y23, Y24, Y29, Y30 | <i>C. kefir</i> |
| Y07, Y21, Y22 | <i>R. mucilogenosa</i> |
| Y08, Y11, Y20, | <i>C. spherica</i> |
| Y09 | <i>C. lusitaniae</i> |
| Y12 | <i>C. famata</i> |
| Y15 | <i>C. rugosa</i> |
| Y16, Y17, Y27 | <i>C. parapsilosis</i> |
| Y19 | <i>C. zeylanoides</i> |
| Y28 | <i>C. colliculosa</i> |

4. CONCLUSION

Traditional fermented milk have been evolved through various cultural and culinary practices and hence the microflora in these products are unique in contributing to the artisanal sensory attributes. The microflora of fermented milk can be affected by hygiene in the production and processing conditions. Environment plays a significant role in contributing yeast and nonstarter microorganisms to these products. At the same time several technologically relevant and probiotic yeast strains co-exist with lactic acid bacteria in fermented milks that can be potentially exploited as starter cultures for industrial applications. For this molecular characterization and safety assessment yeasts isolated from fermented milks need to be studied.

ACKNOWLEDGEMENTS

The authors thank the Kerala Veterinary and Animal Sciences University for the research facilities provided.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Mutukumira AN. Properties of amasi, a natural fermented milk produced by smallholder milk producers in Zimbabwe. *Milchwissenschaft*. 1995;50: 201-205.
- Chandran A, Beena AK, Baghya S, Rathish RL, Rahila MP. Probiotic characterisation of *Enterococcus faecalis* strain isolated from a household dahi sample of Wayanad district, Kerala. *J. Vet. Ani. Sci*. 2022;53(1):70-78.
- Krishna AM, James L, Beena AK, Rajakumar SN, Mercey KA. Changes in microbiological parameters, pH and titratable acidity during the refrigerated and room temperature storage of dahi prepared from the milk of Vechur cows: an indigenous cattle breed of Kerala. *J. Dairy Res*. 2021;88(4):436-9.
- Gadaga TH, Mutukumira AN, Narvhus JA. Enumeration and identification of yeasts isolated from Zimbabwean traditional fermented milk. *Int. Dairy J*. 2000; 10(7):459-66.
- Alvarez-Martin P, Florez AB, Hernández-Barranco A, Mayo B. Interaction between dairy yeasts and lactic acid bacteria strains during milk fermentation. *Food Control*. 2008;19(1):62-70.
- Zohre DE, Erten HÜ. The influence of *Kloeckera apiculata* and *Candida pulcherrima* yeasts on wine fermentation. *Process Biochemistry*. 2002;38(3):319-24.
- Rajkowska K, Kunicka-Styczynsk A. Typing and virulence factors of food-borne *Candida* spp. Isolates. *Int.J.Food Microbiol*. 2018;279:57-63
- Augustin M, Majesté PM, Hippolyte MT. Effect of manufacturing practices on the microbiological quality of fermented milk (Pendidam) of some localities of Ngaoundere (Cameroon). *Int J Curr Microbiol App Sci*. 2014;3(11):71-81.
- Maoloni A, Blaiotta G, Ferrocino I, Mangia NP, Osimani A, Milanović V, Cardinali F, Cesaro C, Garofalo C, Clementi F, Pasquini M. Microbiological characterization of Gioddu, an Italian fermented milk. *Int. J.Food Microbiol*. 2020;23:71-81
- Fleet G.H. Yeasts in dairy products— A review. *J. App. Bacteriol*. 1990;68:199-211
- Alkalbani NS, Osaili TM, Al-Nabulsi AA, Obaid RS, Olaimat AN, Liu SQ, Ayyash MM. *In vitro* characterization and identification of potential probiotic yeasts isolated from fermented dairy and non-dairy food products. *J. Fungi*. 2022; 8(5):544.
- Yadav JS, Bezawada J, Ajila CM, Yan S, Tyagi RD, Surampalli RY. Mixed culture of *Kluyveromyces marxianus* and *Candida krusei* for single-cell protein production and organic load removal from whey. *Bioresource Technol*. 2014;64:119-27.
- Jakobsen M, Narchus J. Yeasts and their possible beneficial and negative effects on the quality of dairy products. *Int. Dairy J*. 1996;6:755–768.
- Kruk M, Trzaskowska M, Ścibisz I, Pokorski P. Application of the “SCOBY” and kombucha tea for the production of fermented milk drinks. *Microorganisms*. 2021;9(1):123.
- de Melo Pereira GV, Maske BL, de Carvalho Neto DP, Karp SG, De Dea

Lindner J, Martin JGP, de Oliveira Hosken B, Soccol CR. What Is Candida Doing in My Food? A Review and Safety Alert on Its

Use as Starter Cultures in Fermented Foods. Microorganisms. 2022;10(9): 1855.

© 2022 Chandran et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/92351>