



## Screening of Some Commonly Used Medicinal Plants against Enteric Human Pathogen *Vibrio cholerae*

Sushmita Choudhury<sup>1\*</sup>, Latika Sharan<sup>2</sup> and M. P. Sinha<sup>1</sup>

<sup>1</sup>Department of Zoology, Ranchi University, Ranchi, Jharkhand-834008, India.

<sup>2</sup>Department of Botany, Ranchi Women's College, Ranchi, Jharkhand-834001, India.

### Authors' contributions

This work was carried out in collaboration between all authors. Author SC designed the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Authors LS and MPS managed the analyses of the study and the literature searches. All authors read and approved the final manuscript.

### Article Information

DOI: 10.9734/EJMP/2015/17973

#### Editor(s):

- (1) Thomas Efferth, Chair, Department of Pharmaceutical Biology, Institute of Pharmacy and Biochemistry, Johannes Gutenberg University, Germany.  
(2) Marcello Iriti, Professor of Plant Biology and Pathology, Department of Agricultural and Environmental Sciences, Milan State University, Italy.

#### Reviewers:

- (1) Atef Mahmoud Mahmoud Attia, Biochemistry department, National Research Centre, Egypt.  
(2) Anonymous, Texas Tech University Health sciences Center, USA.

Complete Peer review History: <http://sciencedomain.org/review-history/9803>

Original Research Article

Received 31<sup>st</sup> March 2015  
Accepted 4<sup>th</sup> May 2015  
Published 17<sup>th</sup> June 2015

### ABSTRACT

**Aim:** Present study aims to screen some of the commonly used medicinal plants (*Mangifera indica*, *Moringa oleifera*, *Pisidium guajava*, *Murraya koenigii* and *Ficus infectoria*) and determination of minimum inhibitory concentration (MIC) against enteric human pathogen *Vibrio cholera*.

**Place and Duration of Study:** Department of Zoology, Ranchi University, Ranchi, between January 2014-July 2014.

**Methodology:** Plant extraction, antibacterial analysis and phytochemical analysis were Done.

**Results:** The different methanolic leaf extracts differed in their vibriocidal properties. The MIC values of the plant extracts against test bacteria were in the range of 2-4 mg/mL by a disc diffusion method. *P. guajava* and *F. infectoria* showed promising vibriocidal property at 4 mg/mL concentration. Whereas the other three plants did not show any activity at the highest concentration. Broth dilution method showed 100% inhibition for all the extracts in the range 4-32 mg/mL.

\*Corresponding author: E-mail: [sushmita.choudhury7683@gmail.com](mailto:sushmita.choudhury7683@gmail.com);

**Conclusion:** All the extracts showed positive vibriocidal activity but *P. guajava* and *F. infectoria* showed promising activity and probably can be used as a safe medicine in diarrhoea.

**Keywords:** *Vibrio cholerae*; diarrhoea; MIC; ZOI.

## 1. INTRODUCTION

Medicinal plants have been used as traditional medicines for various human diseases for thousands of years. A major cause of morbidity and mortality throughout the world is diarrhoeal diseases which mainly effect the infants and children [1,2]. Resistance to antimicrobial agents, is now an increasing problem throughout the globe [3]. And this antibiotic resistance in bacteria of medical importance the interest is increasing our interest in plants as a source of agents for the treatment of microbial diseases [4]. The most significant bioactive compounds obtained from plants are alkaloids, flavonoids, tannins and phenolic substances [5]. It is a common belief that traditional medicine is cheaper and more effective than modern medicine and are being readily accepted for their primary health care needs, and most of this therapy involves the use of plant extracts and their active components [6]. Research on bioactive substances will definitely lead to the discovery of new compounds, which will result in the formulation of new and more potent antimicrobial drugs, which inturn will overcome the problem of resistance to currently available antibiotics. In India many plants are used traditionally as a medicine for the treatment of gastrointestinal disorders such as cholera, diarrhoea and dysentery [7-13]. Various studies have shown that the traditional use of such antidiarrhoeal medicinal plants by investigating the biological activity of extracts of such plants, which suppress gut motility, stimulate water adsorption or reduce electrolyte secretion, have delay intestinal transit, antispasmodic effects [14,15]. Out of the numerous phytochemicals (such as alkaloids, tannins, flavonoids and terpenes etc.) present in active extracts, tannins and flavonoids are thought to be responsible for antidiarrhoeal activity [16-18]. Some of the active ingredients are potentially toxic which should be evaluated for the safety of plant preparations. Few clinical trials have already been evaluated for the safety and tolerability of traditional and herbal medicine preparations which are used to treat diarrhoea and generally indicate that minimal side effects are observed [19].

*Vibrio cholerae* is the causative agent acute intestinal disease, cholera. This organism has a

short incubation period of less than a day, which extends up to five days. The enterotoxin produced by this rod shaped organism causes painless, watery diarrhoea leading to vomiting, severe dehydration, and even death if treatment is not prompt [20]. Cholera can spread as an endemic, epidemic, or pandemic. As a treatment antimicrobial agents are administered for 3-5 days; however it is already reported that a single-dose therapy with tetracycline, doxycycline, furazolidone, or ciprofloxacin has been found to be effective in reducing the duration and volume of diarrhoea [20].

Several plant and plant products are used in the treatment of cholera traditionally. But most of them have not been investigated pharmacologically so as to demonstrate their antibacterial properties, which could support their use as an anticholera or antidiarrheal remedies in traditional medicine. Therefore the objective of the present study was to evaluate the vibriocidal activities of the extracts of some plants viz. *Mangifera indica*, *Psidium guajava*, *Moringa oleifera*, *Murraya koenigii* and *Ficus infectoria* used in Indian traditional healthcare system, against enteric human pathogen- *V. cholerae*.

## 2. MATERIALS AND METHODS

### 2.1 Collection of Plant Material

The fresh and tender leaves were collected, dried in a shade under room temperature for six to seven days and then crushed into coarse powdery substance by using electric grinder. The coarse powdery substance was dried again and was then sieved to get fine powder using the fine plastic sieve, which was then stored in an air tight bottle in the laboratory until required.

### 2.2 Extract Preparation

50 g of the sieved powder was weighed accurately and subjected to extraction in a soxhlet apparatus at room temperature using ~350 mL methanol. The extract obtained was filtered, concentrated after dryness in rotary flash evaporator maintained at 45°C., percentage yield of each extract was calculated and the dried extract was stored in air tight containers at room temperature for further studies.

## 2.3 Phytochemical Analysis

Freshly prepared extracts of the powdered leaves were subjected to phytochemical analyses to find the presence of the following phyto constituents such as flavanoids, alkaloids, carbohydrates, glycosides, tannins, saponins, steroids, proteins, lipids, oils by standard methods [21,22].

## 2.4 Anti-bacterial Analysis

### 2.4.1 Test microorganisms

*Vibrio cholerae* (MTCC 3906) used during the present experiment was procured from Hi-media which is a potential causative pathogen for different diseases.

### 2.4.2 Concentrations screened

0.125, 0.25, 0.5, 1.0, 2.0 and 4.0 mg for agar disc diffusion method and for broth dilution method up to 64 mg/mL concentrations were used according to the sensitivity of samples.

### 2.4.3 Agar diffusion method

Media used consisted of Peptone-10 g, NaCl-10 g and Yeast extract 5 g, Agar 20 g in 1000 mL of distilled water. Initially, the stock cultures of bacteria were revived by inoculating in broth media and grown at 37°C for 18 h. The agar plates of the above media were prepared and wells were made in the plate. Each plate was inoculated with 18 h old cultures (100 µL, 104 cfu) and spread evenly on the plate. After 20 min, the wells were filled with different concentrations of samples. The control wells were filled with Gentamycin along with solvent. All the plates were incubated at 37°C for 24 h and the diameter of inhibition zones were noted.

### 2.4.4 Broth dilution method

Media used consisted of Peptone-10 g, NaCl-10 g and Yeast extract 5 g, in 1000 mL of distilled water. Initially, the stock cultures of bacteria were revived by inoculating in broth media and grown at 37°C for 18 h. The tubes containing above media were prepared, autoclaved and respective concentrations of the samples were added. Each tube was inoculated with 18 h old cultures (100 µL, 104 cfu). A control tube with inoculum and without any sample was prepared along with

a sterile media tube as blank. All the tubes were incubated at 37°C on a shaker with 140 rpm for 24 h and the growth was measured at 660 nm.

## 3. RESULTS AND DISCUSSION

It was revealed from the results that the leaf extracts showed different degree of inhibition against different microorganisms (Table 1, Fig.1). The diameter of zone of inhibition (ZOI) produced depends on several factors broadly classified as extrinsic and intrinsic parameters. The extrinsic parameters like pH of the medium, period and temperature of incubation, volume of the well, concentration of plant extracts and size of inoculum can be fixed and standardized during experiment, hence no error results due to extrinsic factors [23]. However, intrinsic factors such as nature of medicinal plant including its components, solubility and diffusing property are predetermined. Due to variable diffusibility, the antibacterial with very high potency may not demonstrate ZOI commensurate to its efficacy [23].

The medicinal importance of tannins, alkaloids, saponins, phenols, glycosides and flavonoids recorded in the present study is also common in various antibiotics used in treatment of common pathogenic strains, and these phytochemicals (Table 2) are naturally present in the plant extracts which makes the plant useful for treating different diseases and possess a potential of providing useful and safe drugs and drug leads for human use [24,25].


An ideal antidiarrhoeal agent should possess the ability to reverse (i) the increased luminal osmolarity of osmotic diarrhoea, (ii) the increased electrolyte secretion of secretory diarrhoea, (iii) the decreased electrolyte absorption and (iv) the deranged intestinal motility that causes decrease in transit time [26]. Previous studies have been reported which show that methanolic leaf extract of *M. oleifera* effective against castor-oil induced diarrhoea and significantly reduced the time of diarrhoea and total weight of the faeces and the result obtained establish the efficacy of *M. oleifera* plant extracts as antidiarrhoeal agents [15]. Also reports have shown that the leaf extracts of *Psidium guajava*, *Mangifera indica* possess antibacterial properties and are effective in combating various gastrointestinal disturbances [16].

**Table 1. Antimicrobial activity of *Mangifera indica* (M1), *Moringa oleifera* (M2), *Psidium guajava* (M3), *Murraya koenigii* (M4), *Ficus infectoria* (M5) tested against *V. cholerae* by disc diffusion method (Note: NF- MIC not found among the concentrations screened)**

Sample	Concentration						MIC mg
	0.125 mg	0.25 mg	0.5 mg	1.0 mg	2.0 mg	4.0 mg	
M1	0	0	0	0	0	0	NF
M2	0	0	0	0	0	0	NF
M3	0	0	0	0	0	2	4
M4	0	0	0	0	0	0	NF
M5	0	0	0	0	0	2	4
Methanol	0	0	0	0	0	0	NF
	25 µg	50 µg	100 µg	200 µg	400 µg	800 µg	MIC µg
Gentamycin	13	15	18	21	23	27	25

**Table 2. Phytochemical constituents of of *Mangifera indica* (M1), *Moringa oleifera* (M2), *Psidium guajava* (M3), *Murraya koenigii* (M4), *Ficus infectoria* (M5)**

Tests	Samples				
	M1	M2	M3	M4	M5
Carbohydrates	+	+	+	+	+
Glycosides	+	+	+	+	+
Polysaccharides	-	-	-	-	-
Proteins	+	+	+	+	+
Alkaloids	+	+	+	+	+
Steroids	+	+	+	+	-
Triterpenoids	-	+	-	+	-
Flavonoids	+	+	-	-	+
Tannins	-	+	-	-	-
Lipid	+	+	+	+	+
Oils	+	+	+	+	+
Saponins	-	-	-	-	-

**Fig. 1. Inhibition (%) of *V. cholerae* by methanolic extract of *Mangifera indica* (M1), *Moringa oleifera* (M2), *Psidium guajava* (M3), *Murraya koenigii* (M4), *Ficus infectoria* (M5) in broth medium**

Alkizim et al. [27] reported that *Mangifera indica* kernel extract inhibited jejunal contractility and with higher doses caused more pronounced effects. This was probably due to increasing

availability of agonistic molecules to bind to receptors, hence inhibiting the enteric nervous system (ENS), which in turn inhibit muscularis externa activity.

Out of all methanolic extracts from plants consistently provide more antimicrobial activity than those extracted in other more polar substances [28] and this property of possessing high antibacterial activity of methanol extracts is hypothesized to be due to the polarity of the solvent and to the capability to dissolve or diffuse into the medium used in the assays [28].

This study identified novel role of plant extracts and efficacies in controlling the growth of very challenging human pathogen- *V. cholerae*. For example, *Ficus infectoria* is being used for decades as a medicinal plant against various gastrointestinal disturbances, but to our knowledge, this is the first ever report of its efficacy in controlling growth of *V. cholerae*.

#### 4. CONCLUSION

From the above observations we can conclude that all the plant extracts show promising vibriocidal activities and can be used as a safe medicine in diarrhoea.

#### CONSENT

It is not applicable.

#### ETHICAL APPROVAL

It is not applicable.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Food safety as a public health issue for developing countries. In: Unnevehr LJ, ed. Food safety in food security and food trade. Brief 2 of 17 Washington DC: International Food Policy Research Institute. Fernando C, Ramon A; 2003.
2. Halley P. Effect of plants used in Mexico to treat gastrointestinal disorders on charcoal gum acacia induced hyperperistalsis in rats. *J. Ethnopharmacol.* 2010;28:49-51.
3. Albert MJ, Siddique AK, Islam MS, Faruque AS, Ansaruzzaman M, Faruque SM, et al. Large outbreak of clinical cholera due to *V. cholerae* non-O in Bangladesh. *Lancet.* 1993;341:704. [PubMed]
4. Monroe S, Polk R. Antimicrobial use and bacterial resistance. *Curr Opin Microbiol.* 2000;3:496–501. [PubMed]
5. Tiwari P, Kumar B, Kaur M, Kaur G and Kaur, H. Phytochemical screening and extraction: A review. *Int. Pharm. Sci.* 2011; 1:98-106.
6. Winston JC. Health-promoting properties of common herbs. *Am. J. Clin. Nutr.* 1999; 70:491-499.
7. Chopra RN, Nayer SL, Chopra IC. Glossary of Indian Medicinal Plants (3<sup>rd</sup> Edn). Council of Scientific and Industrial Research, New Delhi (India). 1956;7–246.
8. Kala CP, Farooquee NA, Dhar U. Prioritization of medicinal plants on the basis of available knowledge, existing practices and use value status in Uttaranchal, India. *Biodiversity and Conservation.* 2004;13:453-469.
9. Kala CK. Ethnomedicinal botany of the Apatani in the Eastern Himalayan region of India. *J Ethnobiol Ethnomed.* 2005;1:11. DOI:10.1186/1746-4269-1-11. Available:<http://www.ethnobiomed.com/content/1/1/11> Accessed 16 November, 2005.
10. Maikhuri RK, Nautiyal S, Rao KS, Semwal RL. Indigenous knowledge of medicinal plants and wild edibles among three tribal subcommunities of the central Himalayas, India. *Indigenous Knowledge and Development Monitor.* 2000;8:7-13.
11. Mukherjee PK, Saha K, Murugesan T, Mandal SC, Pal M, Saha BP. Screening of anti-diarrhoeal profile of some plant extracts of a specific region of West Bengal, India. *J Ethnopharmacol.* 1998; 60:85-89.
12. Muthu C, Ayyanar M. Raja N, Ignacimuthu S. Medicinal plants used by traditional healers in Kancheepuram District of Tamil Nadu, India. *J Ethnobiol Ethnomed.* 2006;2:43. DOI:10.1186/1746-4269-2-43. Available:<http://www.ethnobiomed.com/content/2/1/43> Accessed 7 October, 2005.
13. Nautiyal S, Rao KS, Maikhuri RK, Semwal RL, Saxena KG. Traditional knowledge related to medicinal and aromatic plants in tribal societies in a part of Himalaya. *J Medicinal Aromatic Plant Sciences.* 2000; 22/4a and 23/1A:441- 528.
14. Horn R, Perry A, Robinson S. A simple solution. *Time.* 2006;42-47.
15. Choudhury S, Sharan L and Sinha MP. Antidiarrhoeal potentiality of leaf extracts of *Moringa oleifera*. *British J. Appl. Sci. Tech.* 2013c;3(4):1086-1096.

16. Choudhury S, Sharan L, Sinha MP. Phytochemical and antimicrobial screening of *Psidium guajava* L. leaf extracts against clinically important gastrointestinal pathogens. J. Natural Prod. Plant Resour. 2012;2(4):524-529
17. Choudhury S, Sharan L and Sinha MP. Pharmacological efficacy of some medicinal plants used for treatment of gastrointestinal diseases. Proceedings of National Seminar on Ecology, Environment & Development, the Ecoscan. 2013a;3: 111-116.
18. Choudhury S, Sharan L and Sinha MP. Phytochemical and antimicrobial standardization of the methanolic leaf extracts on *Murraya koenigii* Linn. Archives des Sciences. 2013b;66(3):67-80.
19. Palombo EA. Phytochemicals from traditional medicinal plants used in the treatment of diarrhoea: Modes of action and effects on intestinal function. Phytoter. Res. 2006;20(9):717-724.
20. Sharma A, Patel VK, Chaturvedi AN. Vibriocidal activity of certain medicinal plants used in Indian folklore medicine by tribals of Mahakoshal region of central, India. Indian. J. Pharmacol. 2009;41(3): 129-133.
21. Sofowora A. Screening plants for bioactive agents. In: medicinal plants and traditional medicinal in Africa. 3<sup>rd</sup> Ed. Spectrum Books Ltd, Sunshine House, Ibadan, Nigeria. 1994;134.
22. Trease GE, Evans WC. Pharmacognosy. 15<sup>th</sup> Ed. Saunders Publishers, London. 2002;42
23. Prasai T, Lekhak B, Baral MP. Antibacterial property of medicinal plants against gram negative bacteria isolated from water. Proceedings of IV National Conference on Science and Technology. 2004;2:410-15.
24. Danish M, Singh P, Mishra G, Srivastava S, Jha KK, Khosa RL. *Cassia fistula* linn. (Amulthus)- An important medicinal plant: A review of its traditional uses. Phytochemistry and Pharmacological Properties J Nat Prod Plant Resour. 2011; 1(1):101-118.
25. Mensah JK, Ikhajiagb B. Phytochemical, nutritional and antibacterial properties of dried leaf powder of *Moringa oleifera* (Lam) from Edo Central Province, Nigeria e, NE Edema, J Emokhor. J Nat Prod Plant Resour. 2012;2(1):107-112.
26. Jinich H, Hersh T. Physicians' Guide to The Etiology of Diarrhoea. Medical Economics Company Inc.; Oradell, NJ; 1982.
27. Alkizim, FO, Matheka DM, Murithi, AW. Dose-dependent myocardial toxicity of *Mangifera indica* during diarrhoea treatment. Afr. J. Pharmacol. Therapeu. 2012;1(2):67-70.
28. Cowan MM. Plant products as antimicrobial agents. Journal of Clinical Microbiology. 1999;12:564-582.

© 2015 Choudhury et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<http://sciencedomain.org/review-history/9803>