



Registry of Ischemic Heart Disease Patients in the Region of Mid Delta, Egypt Data Base and Quality Initiative Project

Abdelrahman Adel Taha Shone^{a++*},
Mohammed Ahmed Elbarbary^{b#},
Suzan Bayoumy Elhefnawy^{bt}
and Mona Adel Elsaïdy^{bt}

^a Tanta Health Insurance Hospital, Gharbia, Egypt.

^b Tanta University, Gharbia, Egypt.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/CA/2023/v12i4367

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/106659>

Original Research Article

Received: 14/07/2023

Accepted: 20/09/2023

Published: 26/09/2023

ABSTRACT

Background: Cardiovascular diseases are significant contributors to global increased mortality rates. coronary artery disease (CAD) is the most common cause of heart disease globally.

Aims: Detection of the characteristics [clinical, electrocardiographic, echocardiographic and coronary angiographic] of ischemic heart disease patients in Med Delta region in Egypt and also their management plan.

⁺⁺ Cardiology Resident;

[#] Lecturer of Cardiology;

[†] Professor of Cardiology;

*Corresponding author: E-mail: Abdelrahmanshone@gmail.com;

Methodology: The registry included 1000 patients of ischemic heart disease (IDH) at Med Delta Region. All patients underwent full medical history, full cardiac examination, detection of ECG changes, cardiac enzymes, other labs, Echocardiography and coronary angiography

Results: Out of 1000 patients studied 72.1% had chronic coronary syndromes and 27.9% had acute coronary syndrome. Seven hundreds and sixty-seven (76.7%) were known to have hypertension and 725 (72.5%) patients were smokers. left anterior descending artery (LAD) was diseased in 54.9 % of the patients. Single vessel disease was affected in 61.7%, 21.5% had double vessel disease and about 16.8 % had multi vessel disease.

Conclusion: Chronic coronary syndrome was the most common presentation in the studied patients with hypertension being the most prevalent risk factor and LAD was the mostly affected vessel.

Keywords: Registry; MID-DELTA; IHD.

1. INTRODUCTION

Cardiovascular illnesses are typically divided into three groups: electrical, circulatory, and structural [1]. The most prevalent type of cardiovascular disorders is coronary artery disease (CAD), a circulatory-related condition that affects the coronary circulation [2]. Cardiovascular diseases are significant contributors to global increased mortality rates [3]. According to the World Health Organization (WHO), 32% of all deaths worldwide in 2019 were attributable to CVD, killing an estimated 17.9 million individuals (WHO, 2021) [4]. CAD is caused by atherosclerosis, which occurs within the walls of the coronary arteries as a result of cholesterol deposits (plaque) in the arteries. The atheromatous plaque grows gradually, leading to the narrowing of the arterial lumen. Atherosclerosis progresses slowly, causing the intima to thicken gradually [5]. There are two types of CAD risk factors: modifiable and non-modifiable. The modifiable factors are obesity, hypertension, diabetes, dyslipidemia, smoking, and physical inactivity that can all be modified and managed [6]. CAD can be diagnosed in patients using invasive and non-invasive techniques. Invasive coronary angiography is the gold standard in the diagnosis of CAD patients. Technology advancements have made it possible for us to non-invasively see coronary calcification [7].

2. METHODOLOGY

2.1 Patient Population

This study was carried out at Tanta Health Insurance Hospital, Tanta Cardiovascular department, Tanta university and also at different centers at the Mid-Delta region including Benha

University Hospitals Mansoura University Hospitals Menofia University Hospitals and Al Mahalla cardiology center. The duration of the study was one year, starting from 1st of March 2021 to 28th of February 2022. this study was a registry study and was conducted on (1000) patients with Ischemic heart diseases.

2.2 Inclusion Criteria

All patients presented with ischemic heart disease.

2.3 Exclusion Criteria

Patients refused coronary angiography.
Patients who were unfit for coronary angiography

2.4 Methods

All patients were subjected to the following:

- I. Full history taking with particular emphasis to age, sex, medical history, and family history with careful evaluation of complaint and common presenting symptoms (onset & duration), and focused on traditional risk factors such as hypertension, diabetes mellitus, and smoking.
- II. Laboratory investigations as, complete blood count, Na, K, serum creatinine, liver enzymes, lipid profile and cardiac enzymes.
- III. Well standardized 12-leads resting electrocardiogram (ECG) was done for all patients with special emphasis on heart rate, cardiac rhythm, QRS duration, conduction abnormalities, ST segment deviation, T-wave changes and ischemic changes were considered when there was primary ST-segment deviation, T-wave

changes and (or) pathological Q-wave in contiguous leads.

- IV. Echocardiography was done for assessment of Ejection fraction (EF) and resting wall motion abnormalities and any other abnormal findings
- V. Coronary angiography was done for all patients.

2.5 Statistical Analysis

Statistical analysis was performed using was calculated by SPSS version 25. The qualitative parameters were described by number of frequency and percentage while the quantitative variables were described by mean, standard deviation and range. Continuous variables were summarized using means or medians based on the normality; normally distributed variables were summarized using the mean and standard

deviation (SD), while the non-normally distributed variables were summarized using the median. Categorical data were summarized as the frequency and percentage.

3. RESULTS AND DISCUSSION

This registry was carried out on 1000 patients all from Mid delta region. The main bulk of our patients were recruited from Tanta Health Insurance Hospital. The rest of the centers are shown in Table 1.

3.1 Distribution of the Studied Cases according to Demographic data

Men constituted the main bulk of our patients as 710/1000 of all patients were males (71%) versus females who were 290 (29%) (Table 2).

Table 1. Distribution of the studied cases according to the number of cases in hospitals

Hospital name	Number of cases	%
Tanta Health Insurance hospital	558	55.8%
Benha University Hospitals	108	10.8 %
Tanta university hospitals	101	10.1%
Mansoura University Hospitals	92	9.2%
Al Mahalla cardiology center	72	7.2%
Menofia University Hospitals	69	6.9%

Table 2. Distribution of the studied cases according to demographic data

	No.	%
Age		
Min-max	30-78	
mean± sd	57.98±10.23	
Median	54	
Gender		
Male	710	71
Female	290	29
Occupation		
Working	390	39
Retired	450	45
Disabled	160	16
Marital status		
Single	140	14
Married	560	56
Widow	240	24
Divorced	60	6
Residency		
Rural	245	24.5
Urban	755	75.5
Educational level		
Illiterate	70	7
High school	357	35.7
University graduate	573	57.3

3.2 Distribution of Clinical Risk Factors and Associated Comorbidities for the Included Populations

Seven hundreds and sixty-seven (76.7%) of our enrolled patients were known to have hypertension which represented the most prevalent risk factor in our study, followed by smoking as the second most common risk factor with 725 (72.5%) patients were smokers (Table 3).

3.3 Distribution of the Studied Cases according to Medical History

According to medical history of ischemic heart disease, we found 217 (21.7%) of patients had previous ACS, 225 (22.5%) had previous CABG and 528 (52.8%) had prior PCI (Table4).

3.4 Distribution of the Studied Cases according to Patient's Presentation

The majority of our patients, seven hundreds and twenty-one (72.1%) had chronic coronary syndrome. About 190(19%) of patients presented with STEMI and 89(8.9%) had NSTEMI. The most common presenting symptom was chest pain with percentage of 79% (Table5).

3.5 Distribution of the Studied Cases according to ECG Findings Upon Admission and Changes During Patients' Hospital Stay

All enrolled patients had 12 leads ECG only 186(18.6) of all patients had normal ECG. Ischemic changes presented in most patients in 814(81.4%) patients (Table 6).

Table 3. Distribution of clinical risk factors and associated comorbidities for the included populations

Medical history and risk factors	No.	%
Hypertension	767	76.7%
Dyslipidaemia	518	51.8%
CKD	300	30%
Diabetes	660	66%
PAD	175	17.5%
Family history for IHD	475	47.5%
BMI		
18.5-24.9	365	36.5%
25-29.9	200	20%
30 or higher	435	43.5%
Smoking	725	72.5%

CKD(chronic kidney disease), PAD(Peripheral arterial disease, BMI (body mass index)

Table 4. Distribution of the studied cases according to medical history

	NO.	%
Previous ACS	217	21.7%
Prior CABG	225	22.5%
Prior PCI	528	52.8%

Table 5. Distribution of the studied cases according to patient's presentation

Presentation of IHD	No.	%
Acute coronary syndrome		
STEMI	190	19%
NSTEMI	89	8.9%
Chronic coronary syndrome with unstable angina	721	72.1%
Main presenting symptoms		
Chest pain	790	79%
Breathlessness	290	29%
Other symptoms as fatigue and sweating.	389	38.9%

STEMI (st-segment elevation myocardial infarction),NSTEMI(non- st-segment elevation myocardial infarction)

Table 6. Distribution of the studied cases according to ECG findings upon admission and changes during patients' hospital stay

Ecg findings	NO.	%
Normal ECG	186	18.6%
Ischemic changes	814	81.4%
Sinus rhythm	724	72.4%
AF	276	27.6%
LBBB	326	32.6%

AF (arterial fibrillation), LBBB(left bundle branch block)




Fig. 1. Presentation of IHD during the registry

STEMI (ST- elevation myocardial infarction) NSTEMI (NON-ST elevation myocardial infarction)




Fig. 2. Pre-tests which were done during the registry

3.6 Distribution of the Studied Cases according to Different Echocardiographic Parameters

Also all population had complete transthoracic echocardiography assessment. Three-hundreds and forty-one (34.1%) of the studied population had preserved ejection fraction while 322 (32.2%) were not preserved the rest of the population 337 (37.3%) had mid-range. From all these population 740 (74%) patients showed abnormal wall motion hypokinesia at rest. (Table7).

3.7 None Invasive Tests Before Coronary Angiography in Patients Subjected to Elective Coronary Angiography

From the chronic patients enrolled with us we found patients with previous non-invasive coronary testing and they were as following, stress ECG in 240 (24%) of all patients, MPI in 270 (27%) of total population enrolled during our

study, and 47 (4.7) underwent CT-Coronary angiography (Table8).

3.8 Distribution of the Studied Cases According to Different Parameters of Coronary Vessels Affected according to Results of Coronary Angiography

According to the results of coronary angiography that was done to all population about 617 (61.7) of the patients had single vessel disease, while 215 (21.5%) had double vessel disease and 168 (16.8%) had multi vessels disease (Table9).

3.9 Distribution of the Studied Cases according to Decision of Management After Coronary Angiography

Regarding the decision that was taken for the patients' management, 602 (60.2%) of patient underwent PCI while, 298 of all patients referred for surgical consultation and only 100 (1%) of the patients were advised to stay on medical treatment (Table10).

Table 7. Distribution of the studied cases according to different echocardiographic parameters

	No.	%
Wall motion hypokinesia	740	74%
EF		
Preserved EF (>50%)	341	34.1%
Mid-range EF(40-50%)	337	33.7%
Impaired EF(<40%)	322	32.2%

EF (Ejection Fraction)




Fig. 3. Distribution of number of vessel affected in the study population

Table 8. None invasive tests before coronary angiography in patients subjected to elective coronary angiography

Test	Stable coronary syndrome	
	No.	%
Stress exercise ECG	240	24%
CT Coronary angiography	47	4.7%
Myocardial Perfusion Imaging	270	27%
Total	557	55.7%

ECG (Electrocardiogram), CT (Computed Tomography)




Fig. 4. Distribution of sites of lesions

3.10 Distribution of the Studied Cases according to the Site of Lesion in Coronary Arteries

Regarding the site of lesions, we noticed that 63 (6.3%) of patients had left main coronary artery disease, 549 of the studied population had left anterior descending coronary artery lesions that accounted the most frequent affected vessel. (Table11).

4. DISCUSSION

The current registry reported general characteristics, demographics, clinical picture and management in 1000 patients with IHD in the Egyptian mid-delta region. In the following discussion, we also would try to compare our overall results with the available national, regional and international data.

Regarding the socio-demographic data, we reported that 45% of the studied populations were retired, 16% were disabled and 39% were regularly working before the admission. about

56% of our patients were married ,14% were single, 7% were divorced and 24% were widows. Patients from urban areas were more commonly found as compared to those from rural areas. We also noted that the previous history of CABG accounted for 22.5%, prior PCI 52.8% and prior ACS 21.7%.

The study done by Doll et al., who reported that 65.8% of their studied patients were married ,7.9% were widow ,6.9% were single 17.3% were divorced or separated and 2% did not answer, about the educational status, 1.5% had elementary or middle school, 8.4% had some high school, 36.9% college graduated. regarding the employment status, the retired patients were 43.8% while not working patients were 14.3%. prior to PCI was detected in 33%, prior MI in 31.5% and Prior CABG in 22.2% [8].

In contrary, the study performed by Maddox et al., who noted that prior PCI was detected in 29.5%, prior CABG in 1.4% and prior ACS in 18.9%. [9].

This contrast may be explained by different number of studied population, different period and geographical distribution of population.

As regarding age, the mean age in our study population was 54.

Parallel to this study, the study done by Kinsara., et al who revealed that the mean age was 56.7 ± 10.4 . [10]. As regard sex, there were 73% males and 27% females, agreeing with our study, the study which was done by Guo., et al which revealed that males approximately were 77% while females were 23%. Also a similar study done by Chiha., et al that detected males were 67.7% while females were 32.3%. [11,12]. Explaining that IHD is more common in males because men's coping with stressful events may be less adaptive physiologically, behaviorally and emotionally. also, estrogens have a protective effect on the cardiovascular system, such a situation makes women who suffer from coronary heart disease show the effects about 10 years later than do men. Therefore, postmenopausal women are considered to be at high risk of cardiovascular disease. Further, a reduction in ovarian function and a reduction in the level of estrogens during menopause induce accelerated aging of the vascular system. The effects include endothelial dysfunction and stiffening of the walls of the large arteries via an unspecified mechanism. Hence, there is a higher incidence of cardiovascular disease in women who have experienced premature menopause, independent of other risk factors such as hypertension, obesity and smoking [13,14]. Regarding the risk factors, hypertension presented in 76.7%, smoking presented in 72.5% of all patients, obesity in 43.5%, diabetes in 66%, dyslipidemia in 51.8% and family history presented in 47.5% of the study population. So, the most frequent risk factor was hypertension followed by smoking, diabetes and dyslipidemia. Concordant to our study, the study of Ntalla., et al who revealed that the most common risk factor was hypertension followed by smoking and dyslipidemia [15]. An Egyptian study which was done by Ellian., et al at Menofia university revealed the following, 64% of patients were obese (64%). Fifty-four percent had hypertension (54%). Thirty-four percent had diabetes (34%). Smoking was documented in (64%) of patients. Positive family history of premature CHD was present in (43%) of the participants [16]. Discordant to this study, the cohort study of Zeitouni., et al who detected that Past or current smoking was the most frequent cardiovascular

factor (60.8%), followed by hypertension (52.8%) and family history of CAD (39.8%). [17]. This discrepancy may be due to a complex interplay between genetic and environmental factors. Several risk factors such as increasing age in our study that may be associated with increased incidence of hypertension. Also family history, obesity, high sodium diets (greater than 3g/day), physical inactivity, have strong and independent correlations with the development of hypertension. The non-invasive tests before coronary angiography in stable coronary syndrome patients involved were stress exercise ECG in 24% of patients, CT angiography in 4.7% and MPI in 27% with total noninvasive pretesting representing 55.7%. In agreement with our study, the study by Bing., et al which detected that the total pre-tests which were done to 43% of all study populations [18]. Disagree with our study the study which was done by Reeh et al.

Table 9. Distribution of the studied cases according to different parameters of coronary vessels affected according to results of coronary angiography

	No.	%
Single vessel disease	617	61.7%
Two vessel disease	215	21.5%
Multi vessel disease	168	16.8%

Table 10. Distribution of the studied cases according to decision of management after coronary angiography

	No.	%
PCI	602	60.2%
Surgical refer	298	29.8%
Medical treatment	100	1%

PCI (percutaneous intervention)

Table 11. Distribution of the studied cases according to the site of lesion in coronary arteries

	No.	%
LM	63	6.3%
LAD	549	54.9 %
LCX	334	33.4 %
RAM	90	9%
RCA	313	31.3 %

LM(left main coronary artery) LAD (left anterior descending artery) LCX(left circumflex coronary artery) RAM(ramus coronary artery) RCA (right coronary artery)

which detected that the MPI was done in 62%, stress exercise ECG was in 35% while patients underwent CT angiography were 83%. [19]. The discrepancy in pre-tests investigations may be due to different hospitals, economic control and different hypothesis of its importance. As regard the symptoms at time of admission, the most presented symptom was chest pain with percentage of 79% followed by dyspnea with percentage of 29% while other symptoms like numbness and fatigue presented by 38.9%. Similarly, in the study performed by Robson et al., all patients presented by chest pain, 19.8% presented by dyspnea. [20]. Another study done by Mieres et al., demonstrated that chest pain was in 90.5%, dyspnea in 9.5% while combined symptoms were in 51.5%. [21]. Regarding ECG changes as (ST-segment deviation >1 mm, inverted T wave, hyper acute T wave in two or more contiguous leads, and new-onset left bundle branch block [LBBB]) (Ischemic evidence) About 81.4% of our studied population had ischemic changes. 32.6% had left bundle branch block and 27.6% of the patients had atrial fibrillation. Parallel study by Simkhada et al., who revealed that total (83.12%) patients had significant ischemic changes [22]. In contrast to our study, the study performed by Moisi et al., who detected that left bundle branch block and QS waves were mostly present [23]. We noticed about 18.6 % of population had completely normal ECG results may consider a good evidence that normal ECG never excludes coronary artery disease as long as symptoms are present. The different ECG results might be due to different study populations as regard number, risk factors and presentations, also the type of studies differ. According to the number of vessels affected, we found that the majority of patients accounted for 61.7% of patients had single vessel disease, 21.5% of them had double vessel disease while only 16.8% of all study population had multi-vessel disease. these results were detected after coronary angiography applied to all patients. Also according to the results of coronary angiography, detected sites of lesions of coronaries were as the following, the most frequent affected vessel was left anterior descending artery with a percentage of 54.9% followed by left circumflex by about 33.4% then right coronary artery 31.3%. the least affected artery was left main coronary artery. Similar results of the study done by Motloch et al., who detected that the percentage of LAD was 52.3% while RCA was 36.9%. this study also reported that about 27.4% had One-vessel affected, 25.2% had two vessels while 47.3% had MVD

[24]. The Egyptian study performed by El-Dosouky et al., who reported that 32.6% of the patients had single vessel disease, 40.3% had double vessel disease and they also reported LAD was the most presented 69.2% followed by RCA 55.7% and that the least presented one was Left main by percentage of 9.6% also has results similar to our study. [25]. As regard the strategy of treatment, our patients were divided into three groups according to the decision of management, the first who underwent PCI, the second who were referred to CABG, the third group who had non-significant lesion and were prescribed medical treatment. In our study, the patients who underwent PCI were 63.2%, the patients who referred to surgical consultation were 26.8% and who received medical treatment were 1%. In the study done by Doll et al., who detected that the patients who underwent PCI were 56.4%, medical treatment in 20.2% and CABG in 23.4%. [8]. about this section is given below. After reading these instructions, please delete this paragraph

5. CONCLUSION

Patients with IHD were commonly males of 4th decade with HTN being the most commonly found risk factor. Acute coronary syndrome recorded about 279 patients (27.9%) while chronic coronary presented in 721 patients (72.1%). The main presented symptom was chest pain. ECG ischemic changes presented in 81.4% of our enrolled patients, wall motion hypokinesia presented in 74% of total population and impaired ejection fraction presented in 32.2% , also coronary angiography was done to all patients at that study with a result of single vessel disease was the most common and the most prominent site of lesion was LAD followed by RCA while left main was the least recorded lesion.

CONSENT AND ETHICAL APPROVAL

An informed consent was obtained from all participants in this research. 2. Any unexpected risks appear during the course of their search will be cleared to participants and the ethical committee on time. 3. Adequate provision to maintain privacy of participants and confidently of the data will be maintained as all patients will be numbered and these numbers will be used instead of names and also all data will be for this research only. 4. Results of our research was only for academic interest to keep the confidentiality of the data.

ACKNOWLEDGEMENTS

Great thanks to my family and my supervisors especially Prof. Dr Mona Adel Elsaidy for her support and efforts.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Kingma DP, Ba JL, Adam. A method for stochastic optimization. 3rd Int Conf Learn Represent ICLR 2015 - Conf Track Proc. 2015;(3):1–15.
- Sood S, Kumar M, Pachori RB. et al. Application of empirical mode decomposition-based features for analysis of normal and CAD heart rate signals. *J Mech Med Biol.* 2016;(1):1-16.
- Roth GA, Johnson C, Abajobir A. et al. Global, Regional, and National Burden of Cardiovascular Diseases for 10 Causes, 1990 to 2015. *J Am Coll Cardiol.* 2017;(1):1–25.
- World Health Organization .Cardiovascular diseases fact sheets; 2021. Available:[https://www.who.int/en/newsroom/factsheets/detail/cardiovascular-diseases-\(cvds\)](https://www.who.int/en/newsroom/factsheets/detail/cardiovascular-diseases-(cvds)).
- Nakahara T, Dweck MR, Narula N. et al. Coronary artery calcification: from mechanism to molecular imaging. *JACC:Cardiovascular Imaging.* 2017;(5):582–593.
- Hajar R. Risk factors for coronary artery disease: Historical perspectives. *HeartViews : the official journal of the Gulf Heart Association.* 2017;(3):109–114.
- Agatston AS, Janowitz WR, Hildner F. et al. Quantification of coronary artery calcium using ultrafast computed tomography. *Journal of the American College of Cardiology.*1990;(4):827–832.
- Doll JA, Jones WS, Likhnygina Y. et al. Prepared study: a study of shared decision-making for coronary artery disease. *Circ Cardiovasc Qual Outcomes.* 2019;12(2):005244.
- Maddox TM, Ho PM, Roe M. et al. Utilization of secondary prevention therapies in patients with nonobstructive coronary artery disease identified during cardiac catheterization: insights from the National Cardiovascular Data Registry Cath-PCI Registry. *Circ Cardiovasc Qual Outcomes.* 2010;3(6):632–41.
- Kinsara AJ, Najm HK, Al Anazi M. et al. Resting heart rate in patients with ischemic heart disease in Saudi Arabia and Egypt. *J Saudi Hear Assoc.* 2011;23(4):225–32.
- Guo Y, Yin F, Fan C. et al. Gender difference in clinical outcomes of the patients with coronary artery disease after percutaneous coronary intervention: a systematic review and meta-analysis. *Medicine (Baltimore).* 2018;97(30).
- Chiha J, Mitchell P, Gopinath B. et al. Gender differences in the prevalence of coronary artery tortuosity and its association with coronary artery disease. *IJC Hear Vasc.* 2017;14:23–7.
- Carr MC. The emergence of the metabolic syndrome with menopause. *The Journal of Clinical Endocrinology & Metabolism.* 2003;88(6):2404-11.
- Mudali S, Dobs AS, Ding J. et al.,. Endogenous postmenopausal hormones and serum lipids: the atherosclerosis risk in communities study. *The Journal of Clinical Endocrinology & Metabolism.* 2005;90(2): 1202-9.
- Ntalla I, Kanoni S, Zeng L. et al. Genetic risk score for coronary disease identifies predispositions to cardiovascular and noncardiovascular diseases. *J Am Coll Cardiol.* 2019;73(23):2932–42.
- Ellaien A, Reda A, Elkersh A. The Pattern of Risk Factor Profile and Dutch Lipid Clinic Network Criteria of Familial Hypercholesterolemia in Egyptian Patients with Premature Coronary Heart Disease. 2021;1:331:146.
- Zeitouni M, Clare RM, Chiswell K. et al. Risk factor burden and long-term prognosis of patients with premature coronary artery disease. *J Am Heart Assoc.* 2020;9(24):017712.
- Bing R, Singh T, Dweck MR, Mills NL. et al. Validation of European Society of Cardiology pre-test probabilities for obstructive coronary artery disease in suspected stable angina. *Eur Hear Journal-Quality Care Clin Outcomes.* 2020;6(4):293–300.
- Jacob Reeh, et al. Prediction of obstructive coronary artery disease and prognosis in patients with suspected stable angina. *European Heart Journal.* 2019; 7;40 (18):1426-35.
- Reeh J, Therming CB, Heitmann M. et al. Prediction of obstructive coronary

- artery disease and prognosis in patients with suspected stable angina. *European Heart Journal*. 2019;40(18):1426-35.
20. Robson J, Ayerbe L, Mathur R. et al. Clinical value of chest pain presentation and prodromes on the assessment of cardiovascular disease: A cohort study. *BMJ Open*. 2015;5(4):e007251.
 21. Mieres JH, Heller G V, Hendel RC. et al. Signs and symptoms of suspected myocardial ischemia in women: Results from the What is the Optimal Method for Ischemia Evaluation in Women? Trial. *J Women's Heal*. 2011;20(9):1261–8.
 22. Simkhada R, Khan B, KC SS. et al. Angiographic Significance of ST Depression in Anterior Leads in Acute Inferior ST Elevation Myocardial Infarction. *Nepal Hear J*. 2021;18(2):7–10.
 23. Moisi MI, Rus M, Bungau S. et al. Acute coronary syndromes in chronic kidney disease: clinical and therapeutic characteristics. *Medicina (B Aires)*. 2020; 56(3):118.
 24. Motloch LJ, Reda S, Larbig R. et al. Characteristics of coronary artery disease among patients with atrial fibrillation compared to patients with sinus rhythm. *Hell J Cardiol*. 2017;58(3):204–12.
 25. El-Dosouky II& Abomandour HG. Fragmented QRS complex in acute coronary syndrome, does it have significance in the emergency room? A study from Egypt. *J Indian Coll Cardiol*. 2019;9(3):131.

© 2023 Shone et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/106659>