



Prevalence and Risk Factors of In-Hospital Mortality of ST Elevation Myocardial Infarction (STEMI) Patients in Bekasi Regency General Hospital in 2021

Khanza Rizqullah Syauqi ^{a++*}, Mailani Karina Akhmad ^{a#} and Omar Mokhtar Siregar ^{a#}

^a Bekasi Regency General Hospital, Bekasi Regency, Indonesia.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/CA/2023/v12i4369

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here:

<https://www.sdiarticle5.com/review-history/106600>

Original Research Article

Received: 20/07/2023

Accepted: 23/09/2023

Published: 02/10/2023

ABSTRACT

Background: Acute Coronary Syndrome (ACS) is a major cardiovascular problem and has been one of the leading death in the world. ST-Elevation Myocardial Infarction (STEMI) is an indicator of total coronary artery occlusion which requires revascularization to restore the blood flow and myocardial reperfusion using fibrinolytic agents or primary Percutaneous Coronary Intervention (PCI). Early treatment showed better outcomes and reduced in-hospital mortality in ACS patients.

⁺⁺ General Practitioner;

[#] Cardiologist;

^{*}Corresponding author: E-mail: khanzasyauqi02@gmail.com;

Aims: To determine the prevalence of Acute Coronary Syndrome (ACS) patients and the risk factors of in-hospital mortality of ST-Elevation Myocardial Infarction (STEMI) patients at Bekasi Regency General Hospital in 2021.

Place and Duration of Study: Cardiology Department and Emergency Room Unit in Bekasi Regency General Hospital, December 2021 to March 2022

Methods: This study is a descriptive cross-sectional study using secondary data (medical records) which included all the patients over 35 years old that admitted through the ER from January to December 2021 with complaints including chest pain or shortness of breath which were later hospitalized. The final sample used in this study is 117 subjects.

Results: There were 117 cases of ACS in 2021. They were 14% STEMI, 15% Non-ST Elevation Myocardial Infarction (NSTEMI), and 71% Unstable Angina Pectoris (UAP). Most patients were male (54%), and aged <65 years old (85%). The in-hospital mortality was 19% STEMI, 17% NSTEMI, and 12% UAP. Among 16 STEMI patients, 81% were male and 94% were aged <65 years old. There were 69% who came to the emergency room <12 hours and only 50% underwent fibrinolytic.

Conclusion: The risk factors of in-hospital mortality of STEMI patients were 40% late onset, 23% anterior STEMI, 20% hypertension, 60% diabetes mellitus, 50% chronic kidney disease, 33% tachycardia, and 100% Killip III.

Keywords: Acute coronary syndrome; St-Elevation myocardial infarction; in-hospital mortality; prevalence; risk factors.

ABBREVIATIONS

Abbreviation	Definition
ACS	: Acute coronary syndrome
WHO	: World Health Organization
STEMI	: ST-Elevation Myocardial Infarction
PCI	: Percutaneous Coronary Intervention
NSTEMI	: Non-ST Elevation Myocardial Infarction
UAP	: Unstable Angina Pectoris
ECG	: Electrocardiogram
CKMB	: Creatine kinase-MB
LVH	: Left Ventricular Hypertrophy
LBBB	: Left Bundle Branch Block
ER	: Emergency Room
JAC	: Jakarta Acute Coronary Syndrome

million deaths, while ACS caused 7.4 million deaths. In 2019 in Indonesia, ischemic heart disease was the second medical case that mostly caused mortality [3].

Coronary heart disease is a disease caused by plaque accumulation in the coronary arteries that supply oxygen to the heart muscles [4]. Acute coronary syndrome is also caused by increased oxygen demand, reduced blood oxygen transport, and is often caused by reduced coronary flow as a result of narrowing or obstruction of the arteries due to atherosclerosis [5]. Acute coronary syndrome consists of Unstable Angina Pectoris (UAP), non-ST elevation myocardial infarction (NSTEMI), and ST-elevation myocardial infarction (STEMI) [6,4]. Angina pectoris is a clinical syndrome caused by an imbalance between demand and supply of coronary artery flow [6]. Non-ST Elevation Myocardial Infarction (NSTEMI) is an acute coronary syndrome with clinical symptoms of chest pain typical of infarction, electrocardiogram (ECG) without ST-segment elevation, and elevated cardiac enzymes (CKMB, troponin T). NSTEMI can cause various complications, such as acute pulmonary edema, cardiac arrest, and even death [7].

Acute ST Elevation Myocardial Infarction (STEMI) is a clinical syndrome defined by myocardial ischemia symptoms related to persistent ST elevation and release of myocardial necrosis biomarkers [8]. Acute myocardial infarction is diagnosed based on the

1. INTRODUCTION

Based on the data from WHO in 2019, around 17.9 million deaths are caused by Cardiovascular Diseases each year. The majority of CVD deaths are caused by heart attacks and strokes, and more than half of these deaths are premature deaths among people under 71 years old [1]. Around 7.5 million of cardiovascular-related deaths are due to Ischemic Heart Disease, which among the 1.8 million deaths per year are mostly caused by Acute Coronary Syndrome (ACS) and sudden death [2]. Acute Coronary Syndrome (ACS) is a major cardiovascular problem. According to World Health Organization (WHO) in 2015, cardiovascular disease caused 17.5

typical clinical symptoms of chest pain, increased cardiac biomarkers, and changes in electrocardiogram patterns [9]. Increased cardiac-specific markers found in acute myocardial infarction are MB isoenzyme of creatine kinase (CKMB), troponin T, and troponin I. CKMB isoenzymes are plentiful in cardiac muscle cells. Increased levels of CKMB indicate the extent and severity of infarction in the heart muscle [9]. On the electrocardiogram, an assessment of ST elevation is performed at the J point and is found in two opposite leads. The ST-segment elevation threshold value in the diagnosis of STEMI for males and females in most leads is 0.1 mV [10]. ST-segment elevation without Left Ventricular Hypertrophy (LVH) or Left Bundle Branch Block (LBBB) is a new ST segment elevation at the J point $>2\text{mm}$ (0.2 mV) in males or $>1.5\text{ mm}$ (1.5 mV) in females in leads V2-V3 and/or $>1\text{ mm}$ (0.1 mV) in another precordial lead or limb lead, in at least 2 contiguous leads [8,10].

Typical chest pain in STEMI located in the substernal, with >20 minutes duration, accompanied by cold sweat, chest pain that is described as feeling like crushed and can spread to the left arm, back, jaw, or epigastrium. Risk factors for STEMI include diabetes, cholesterol, high blood pressure, chronic kidney disease, and heredity [6]. STEMI is an indicator of total coronary artery occlusion. This condition requires revascularization to restore blood flow and myocardial reperfusion as soon as possible, either medically using fibrinolytic agents or mechanically through primary percutaneous coronary intervention [10]. Several factors must be considered when choosing reperfusion therapy. Initiation of reperfusion management does not need to wait for the results of increased cardiac biomarkers [8,10].

Over three million cases of STEMI developed each year globally and over four million individuals displayed the pathology of STEMI [11]. ACS, especially STEMI is a major health problem in Indonesia. There are 608,043 ACS cases estimated each year with an estimated number of 296,725 cases of STEMI each year in Indonesia [12]. One of the interventions used to treat STEMI patients is Percutaneous Coronary Intervention (PCI).

Percutaneous coronary intervention (PCI) is a non-surgical intervention procedure using a catheter to dilate or open narrowed coronary

arteries due to atherosclerosis or thrombosis with ballooning or deploying a stent. For STEMI patients who come to a hospital with PCI facilities, primary PCI must be performed within 90 minutes with the onset of complaints <12 hours. For patients who come to a hospital without PCI facilities, fibrinolytic therapy should be initiated at <12 hours of onset and if primary PCI cannot be performed within 120 minutes of first medical contact [8,6]. Early reperfusion therapy can reduce in-hospital mortality rates and show better outcomes in patients with acute myocardial infarction. However, STEMI still causes various complications, including myocardial dysfunction, heart failure, cardiogenic shock, arrhythmias, mechanical complications, and pericarditis [13].

This study attempts to determine the prevalence of Acute Coronary Syndrome patients who were admitted through the emergency room in Bekasi Regency General Hospital and determine the risk factors of in-hospital mortality among STEMI patients. With the insight on the prevalence and risk factors of Acute Coronary Syndrome, health systems are expected to understand the burden of the disease in order to improve the service and care for the patients, especially in taking appropriate care in a timely manner for STEMI patients.

2. MATERIALS AND METHODS

This study is a descriptive cross-sectional study using secondary data. The study subjects included all the patients over the age of 35 that admitted through the ER in Bekasi Regency General Hospital from January to December 2021 with chest pain or shortness of breath and later hospitalized. The research was conducted from December 28th, 2021 to March 28th, 2022. This study used secondary data from the medical records of Bekasi Regency General Hospital.

The initial study was carried out by collecting register data for all patients with acute coronary syndrome who were admitted through the ER from January to December 2021. From the data register, there were 149 patients with acute coronary syndrome in the past year. Consequently, a search for medical record status was conducted to collect the necessary data. Out of the 149 statuses, 12 statuses could not be found and 20 statuses were not included in the inclusion criteria. Therefore, the final sample in this study is 117 statuses (Fig. 1).




Fig. 1. Sampling process

3. RESULTS

3.1 Acute Coronary Syndrome Characteristics

This study shows that the number of acute coronary syndrome cases admitted through the ER in the medical record data of Bekasi Regency General Hospital from January 1 to December 31, 2021, was 149 cases. Later, out of these 149 cases, 12 cases had no medical records, and 20 cases were not included in the inclusion criteria. Thus, the final sample of this study was 117 cases. Out of the 117 cases, there were 16 cases of STEMI (14%), 18 cases of NSTEMI

(15%), and 83 cases of UAP (71%). The distribution of Acute Coronary Syndrome (ACS) patients in Bekasi General Hospital in 2021 can be seen in Fig. 2.

This study found that more than half of acute coronary syndrome patients are male patients (54%), while the rest are female patients (46%). Age is a part of the risk factor for acute coronary syndrome. This study classified the patients' age into the under 65 years old age group and the over 65 years old age group. Out of 117 patients, 99 patients are under 65 years old (85%), and 18 patients are over 65 years old (15%).




Fig. 2. Distribution of Acute Coronary Syndrome (ACS) patients in Bekasi regency general hospital in 2021

Patients' comorbidities were dominated by hypertension (85%) with 99 patients, 46 patients suffering from diabetes mellitus (39%), and 36 patients suffering from chronic kidney disease (31%). Out of 117 acute coronary syndrome cases in Bekasi Regency General Hospital during the period of 1 January to December 31, 2021, 16 of these cases died during in-hospital treatment at the hospital. The dead cases were 3 STEMI cases (19%), 3 NSTEMI cases (17%), and 10 UAP cases (12%). Among the 16 acute coronary syndrome patients who died, 7 of them were male (11%) and 9 of them were female (17%). Based on the age group, the majority of acute coronary syndrome patients that were died from the under 65 years old age group (15%), while there was only 1 patient over 65 years old that died (6%). Comorbidities among the patients that died included hypertension; 11 patients (11%), diabetes mellitus; 9 patients (20%), and chronic kidney disease; 9 patients (25%) (Table 1).

3.2 STEMI Patients Clinical Characteristics

Clinical and demographic data of STEMI patients in Bekasi Regency General Hospital in 2021 is summarized in Table 2. Out of 16 STEMI patients, the majority of them are under 65 years old, which were 15 patients (94%) and 13 male patients (81%). Out of these 16 STEMI patients, three of them died (20%) during in-hospital treatment and three of them were under 65 years old. The majority of the patients that died were male, which were 2 patients (15%).

STEMI patients who came to the ER of Bekasi Regency General Hospital complained about chest pain with onset of attack <12 hours, which were 11 patients (69%). Only five patients (31%) come to the ER with the onset of attack >12

hours. Out of 11 patients with initial onset <12 hours, only 8 of them received fibrinolytic therapy, 2 patients did not receive it due to fibrinolytic contraindications, while the other one rejected the fibrinolytic therapy. Most of the acute myocardial infarctions were located in the anterior part of the heart, with 13 patients (81%), while three patients (19%) were in the non-anterior part of the heart. All three patients who died had a myocardial infarction in the anterior part of the heart.

Risk factors of STEMI patients are hypertension (94%), diabetes mellitus (31%), and chronic kidney disease (13%). Symptoms and clinical signs, of the patients, when they arrived were low systolic blood pressure (31%), high pulse rate (19%), and no arrhythmia (0%). Killip classifications when the patient arrives are Killip I (13%), Killip II (44%), Killip (13%), and Killip IV (31%) (Table 2).

4. DISCUSSION

4.1 Acute Coronary Syndrome Characteristics

Acute coronary syndrome is classified into three types, acute ST elevation myocardial infarction with (STEMI), acute non-ST elevation myocardial infarction (NSTEMI), and unstable angina pectoris (UAP). Unstable angina pectoris and NSTEMI can be distinguished from cardiac biomarkers. If a significant increase is found, the diagnosis is NSTEMI and if there is no significant increase, the diagnosis is UAP [10]. The result in (Fig. 2) is in line with the data register from the Jakarta Acute Coronary Syndrome (JAC) in 2008-2009 where there were more UAP patients than STEMI and NSTEMI patients [14].

Table 1. Acute coronary syndrome patient characteristic

Variable	Amount (n=117)	Death (n=16)
STEMI	16 (14%)	3 (19%)
NSTEMI	18 (15%)	3 (17%)
UAP	83 (71%)	10 (12%)
Sex		
Male	63 (54%)	7 (11%)
Female	54 (46%)	9 (17%)
Age		
<65 years old	99 (85%)	15 (15%)
>65 years old	18 (15%)	1 (6%)
Hypertension	99 (85%)	11 (11%)
Diabetes Mellitus	46 (39%)	9 (20%)
Chronic Kidney Disease	36 (31%)	9 (25%)

Table 2. STEMI patient characteristic

Variable	Amount (n=16)	Death (n=3)
Age		
<65 years old	15 (94%)	3 (20%)
>65 years old	1 (6%)	0 (0%)
Sex		
Male	13 (81%)	2 (15%)
Female	3 (19%)	1 (33%)
Onset		
<12 hours	11 (69%)	1 (9%)
>12 hours	5 (31%)	2 (40%)
Reperfusion		
Fibrinolytic	8 (50%)	1 (13%)
Non-Fibrinolytic	8 (50%)	2 (25%)
STEMI location		
Anterior	13 (81%)	3 (23%)
Non-Anterior	3 (19%)	0 (0%)
Hypertension		
Hypertension	15 (94%)	3 (20%)
Diabetes Mellitus		
Diabetes Mellitus	5 (31%)	3 (60%)
Chronic Kidney Disease		
Chronic Kidney Disease	2 (13%)	1 (50%)
Hypotension		
Hypotension	5 (31%)	0 (0%)
Tachycardia		
Tachycardia	3 (19%)	1 (33%)
Killip		
I	2 (13%)	0 (0%)
II	7 (44%)	0 (0%)
III	2 (13%)	2 (100%)
IV	5 (31%)	1 (20%)
Arrhythmia		
Arrhythmia	0 (0%)	0 (0%)

The result is also in line with the study finding in 2014 which stated that the 41-60 age group is the most affected age group [15]. Another study stated that old age is the main predictor of in-hospital mortality in ACS patients [16]. The risk of developing acute coronary syndrome is increasing in men who are over 45 years old and women who are over 55 years old [15]. The prevalence of myocardial infarction is higher in individuals over the age of 60 compared to individuals under the age of 60 [11]. A previous study on characteristics of in-hospital mortality in ACS patients found an increased risk of comorbidities among the elderly population [17].

Patients' comorbidities can also be a risk factor for acute coronary syndrome. This is explained by the result in Fig. 2 which was in line with a study finding in 2018 that found that hypertension is the most common risk factor for acute coronary syndrome patients [18]. Increased systemic blood pressure in hypertension causes resistance to the blood pump from the left ventricle and causes ventricular hypertrophy. Ventricular hypertrophy causes an increase in myocardial oxygen demand thus increasing the

workload of the heart. Ultimately, this can trigger angina and myocardial infarction. The incidence of acute coronary syndrome in patients with hypertension is five times greater than in normotensive patients. Previous studies found that hypertension is a major risk factor for acute coronary syndrome [18,17]. However, this result is not in line with a study by Pramudyo, which found that hypertension is not related to the mortality risk in ACS patients [16].

The second most common risk factor is diabetes mellitus. Patients with diabetes mellitus have a 200% greater risk of experiencing acute coronary syndrome than patients without diabetes mellitus. Another study states that male patients with diabetes mellitus have a 50% greater risk of experiencing acute coronary syndrome, while female patients with diabetes mellitus have twice the risk compared to patients without diabetes mellitus [18].

According to the 2018 Handbook of Management of Acute Coronary Syndrome—*Buku Pedoman Tata Laksana Sindrom Koroner Akut*, the prevalence of NSTEMI and UAP is greater than

STEMI. NSTEMI patients' initial mortality is lower than STEMI patients, however, after 6 months, the mortality of both is comparable. In the long term, NSTEMI patients' mortality will be greater subsequently [10]. Gender, age, and comorbid risk factors for acute coronary syndrome patients affect mortality. The result in Table 1 is in line with a study in 2017 which found that the causes of cardiogenic shock after acute coronary syndrome can be exacerbated by old age, diabetes mellitus, previous myocardial infarction, chronic occlusion of the ascending left coronary artery, and decreased ejection fraction which led to death [19].

4.2 STEMI Patients' Clinical Characteristics

Based on the study on ACS patient clinical profiles, almost half of the ACS patients have one or more comorbidities [20]. A research in 2017 found that the number of mortalities in acute coronary syndrome patients that were not treated properly is 40% and the majority of the deaths occurred in the first two hours since the initial onset (Table 2). The mortality and morbidity that occur in STEMI patients increase with age. Besides, STEMI has a high mortality number, oftentimes due to complications. The predictors of STEMI patient mortality include Killip III and IV, arrhythmia, anterior STEMI, non-reperfusion, chronic kidney disease, tachycardia, STEMI onset >12 hours, and diabetes mellitus [19].

The cause of death in myocardial infarction is multifactorial and depends on the elapsed time duration since the initial attack. Sudden death is commonly caused by ischemia precipitating ventricular arrhythmias during the acute phase of myocardial infarction. Other complications that commonly occur are ventricular tachycardia/ventricular fibrillation (13%), left ventricular dysfunction (11.9%), cardiogenic shock (10.7%), atrial tachycardia/atrial fibrillation (7.1%), and right ventricular infarction (7.1%). Mortality in patients who did not receive reperfusion therapy was 13.2%. The purpose of reperfusion is not only for restoration of arterial patency but also for restoration of coronary flow which has been shown to be associated with reduced heart wall motion abnormalities and improved clinical outcomes [19].

Patients with a previous acute myocardial infarction had the same hospital outcomes as patients without a previous acute myocardial

infarction but had a higher recurrence rate at 6 months. Refractory ischemia or recurrent ischemia is symptomatic ischemia within 10 minutes with ST-segment deviation or definitive T-wave inversion, and/or hypotension, pulmonary edema, or a heart murmur representing myocardial ischemia (despite the use of nitrates and beta-blockers or calcium channel blockers), occurs in up to 20% of ischemic patients. Refractory ischemia has mortality within 30 days two times greater than patients with ST-segment elevation and even greater mortality risk than patients without ST-segment elevation [19].

With the prevalence of the disease presented, the healthcare system could use it to estimate the burden of monetary cost, life expectancy, life quality, morbidity, etc. [21]. By referring to the result of this study, proper care and treatment could be conducted by considering the comorbidity, age, onset, and other variables of acute coronary syndrome patients and STEMI patients in particular.

4.3 Study Limitations

This study did have some limitations such as the size of the population which is limited to cases in Bekasi Regency General Hospital from January to December 2021. Hence the result might not be applicable to a broader population with longer time intervals since the result might differ with broader and more varied data. Besides, the selection bias also might occur due to the sample not including other ACS cases outside of cases in Bekasi Regency General Hospital.

5. CONCLUSION

Based on the study and discussions, it can be concluded that the number of acute coronary syndrome patients admitted through the ER of Bekasi Regency General Hospital in 2021 is 149 cases. There are 12 cases without medical records and 20 cases excluded from the inclusion criteria, thus, the final sample in this study is 117 cases. Acute coronary syndrome patients are classified into 16 STEMI patients (14%), 18 NSTEMI patients (15%), and 83 UAP patients (71%). Most patients of acute coronary syndrome were male, with a total of 63 patients (54%) while the others were female, with a total of 54 patients (46%). Most patients with acute coronary syndrome were under 65 years old, with a total of 99 patients (85%), and the rest were under the age of 65, with a total of 18 patients (15%).

The risk factors for acute coronary syndrome patients were 99 hypertension patients (85%), 46 diabetes mellitus patients (39%), and 36 chronic kidney disease patients (31%). The number of acute coronary syndrome patients who died during in-hospital treatment was 16 patients, including three STEMI patients (19%), three NSTEMI patients (17%), and ten UAP patients (12%). There were 7 male patients (11%) and nine female patients (17%). The number of patients under the age of 65 is 15 patients (15%), while the other patient over the age of 65 is only one patient (6%). The risk factors included 11 hypertension patients (11%), nine diabetes mellitus patients (20%), and nine chronic kidney disease patients (25%). Out of 117 cases, the number of STEMI patients is 16, and three of them have died during in-hospital treatment.

The majority of STEMI patients are under the age of 65 years, with a total of 15 patients (94%), while there was only one patient over the age of 65 (6%). Most of them were male patients, with a total of 13 patients (81%) and the other were female patients with a total of three patients (19%). STEMI patients that come through the ER with clinical symptoms of chest pain with <12 hours onset are 11 patients (69%), while patients with >12 hours onset are five patients (31%). STEMI patients that received fibrinolytic therapy are eight patients (50%) and only patients with initial onset <12 hours received the therapy. There were two patients who did not receive the fibrinolytic due to contraindications and there was one patient that rejected the fibrinolytic therapy.

Most of the acute myocardial infarctions were located in the anterior part of the heart with a total of 13 patients (81%), while the rest were located in the non-anterior part of the heart with a total of three patients (19%). STEMI risk factors were 15 hypertension patients (94%), five diabetes mellitus patients (31%), and two chronic kidney disease patients (13%). Symptoms and clinical signs when STEMI patients arrived included 5 low systolic blood pressure (31%), high pulse rate (19%), and no arrhythmia (0%). Killip classifications when STEMI patients arrived were two Killip I patients (13%), seven Killip II patients (44%), two Killip III patients (13%), and five Killip IV patients (31%).

The result of this study could be used as a reference to estimate the acute coronary syndrome burden by the healthcare system in order to take proper care of the patients in a timely manner by considering the risks and the

prevalence of the disease. Additionally, further studies could consider to use analytical research methods to study the correlation between the risk factors with acute coronary syndrome condition and use patient data that received PCI procedure. In addition, this study could be used as a basis for further research with a wider period of acute coronary syndrome patient registry.

CONSENT

As per international standard or university standard, patient(s) written consent has been collected and preserved by the author(s).

ETHICAL APPROVAL

The study is using the medical data for research purposes was approved by Bekasi Regency General Hospital by following the applicable ethical regulations and respecting the privacy concern. An additional permission from the National and Political Unity Agency has been obtained to conduct the study in the government institution (government-owned hospital).

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. World Health Organization. Cardiovascular Diseases (CVDs); 2021. Available:[https://www.who.int/news-room/fact-sheets/detail/cardiovascular-diseases-\(cvds\)](https://www.who.int/news-room/fact-sheets/detail/cardiovascular-diseases-(cvds))
2. Bueno H. Epidemiology of acute coronary syndromes. In *The ESC Textbook of Cardiovascular Medicine* (3rd edition). The European Society of Cardiology Series/. 2018;1214–1218. Available:<https://doi.org/https://doi.org/10.1093/med/9780198784906.003.0305>
3. World Health Organization. (2020). Global Health Estimates 2020: Deaths by Cause, Age, Sex, by Country and by Region; 2000-2019. World Health Organization. Available:<https://www.who.int/data/gho/data/themes/mortality-and-global-health-estimates/ghe-leading-causes-of-death>
4. Maulidia R. QRS complex fragmentation as a predictor of mortality and rehospitalisation in the first 6 months in acute coronary syndrome (STEMI, NSTEMI) in Haji Adam Malik General

- Hospital, Medan. Universitas Sumatera Utara; 2021.
5. Wibowo A, Andina M. Prevalence of hypertension with Acute Coronary Syndrome (ACS) in Hajj Hospital, Medan. *Jurnal Kedokteran Anatomica*. 2018;1(3).
 6. Indonesian Cardiovascular Specialist Association. Clinical Practice Guideline (CPGs) and Clinical Pathway (CP) in cardiovascular disease (1st ed.) (1st ed.). Indonesian Cardiovascular Specialist Association; 2016.
 7. Elfi EF. Acute coronary syndrome with complications of acute pulmonary edema and cardiac arrest. *Jurnal Kesehatan Andalas*. 2015;4(2):613–617. Available:<https://doi.org/10.25077/jka.v4i2.309>
 8. Gayatri NI, Firmansyah SSSH, Rudiktyo E. Predictors of in-hospital mortality in acute ST-Elevation Myocardial Infarction (STEMI) patients in Dr. Dradjat Prawiranegara Regional General Hospital, Serang, Indonesia. *Cermin Dunia Kedokteran Journal*. 2016;43(3): 171–174.
 9. Kurniawan LB, Bahrun U, ER D, Arif M. The relationship between Creatine Kinase-MB levels and mortality in patients with myocardial infarction during hospitalization in Dr. Wahidin Sudirohusodo Hospital, Makassar. *Buletin Penelitian Kesehatan*. 2015;43(2):125–130. Available:<https://doi.org/10.22435/bpk.v43i2.4146.125-130>
 10. Indonesian Cardiovascular Specialist Association. Guidelines for acute coronary syndrome management (4th ed.) (Keempat). Indonesian Cardiovascular Specialist Association; 2018.
 11. Salari N, Morddarvanjoghi F, Abdolmaleki A, Rasoulpoor S, Khaleghi AA, Hezarkhani LA, Shohaimi S, Mohammadi M. The global prevalence of myocardial infarction: a systematic review and meta-analysis. *BMC Cardiovascular Disorders*. 2023;23(206). Available:<https://doi.org/https://doi.org/10.1186/s12872-023-03231-w>
 12. Fahri I. STEMI in Indonesia - Social-Economics & demographics challenges. *Transcatheter Cardiovascular Therapeutics Asia Pacific*; 2022. Available:https://www.summitmd.com/html/technical/coronary/inside_topics_view.php?cmode=slide&subset=&part=e&year=&no_case=8625
 13. Wilar GI, Panda AL, Rampengan SH. 6 Months observation of Major Cardiovascular Events in patients with Acute ST-Elevation Myocardial Infarction (STEMI) in Prof. Dr. R. D. Kandou General Hospital, Manado during January-December 2017 period. *Jurnal Medik Dan Rehabilitasi (JMR)*. 2019;1(3):1–8.
 14. Dharma S, Juzar DA, Firdaus I, Soerianata S, Wardeh AJ, Jukema JW. Acute myocardial infarction system of care in the third world. *Netherlands Heart Journal*. 2012;20(6):254–259. Available:<https://doi.org/10.1007/s12471-012-0259-9>
 15. Torry SRV, Panda L, Ongkowitzaya J. Description of risk factors of patients with Acute Coronary Syndrome. *Jurnal E-CliniC (ECI)*. 2014;2(1):1–8.
 16. Pramudyo M, Yahya AF, Martanto E, Tiksnadi BB, Karwiky G, Rafidhinar R, Putri GNI. Predictors of in-hospital mortality in patients with acute coronary syndrome in Hasan Sadikin Hospital, Bandung, Indonesia: A retrospective cohort study. *Acta Medica Indonesiana*. 2022;54(3):379–388.
 17. Tessy DB, Pramudyo M, Cool CJ. Characteristics of in-hospital mortality among patients with acute coronary syndrome: A Single-Center Study in West Java, Indonesia. *Althea Medical Journal*. 2021;8(2):99–103. Available:<https://doi.org/https://doi.org/10.15850/amj.v8n2.2281>
 18. Suling FRW, Patricia MI, Suling TE. Prevalence and risk factors of acute coronary syndrome at Universitas Kristen Indonesia General Hospital. *Majalah Kedokteran UKI*. 2018;34(3): 110–114.
 19. Sutarmini SANY, Gunadhi IGNP. ST-Segment elevation as a predictor of Major Adverse Cardiovascular Events (MACE) in Acute Coronary Syndrome (ACS) patients during hospitalization in Sanglah General Hospital, Denpasar. Udayana University; 2017.
 20. Adam AA, Tiluata LJ, Yunita L, Putra MP, Wilujeng N, Homalessy LV, Hutabarat DCF. Clinical Profile of Acute Coronary Syndrome Patients in Kupang: a Result from 1-Year iSTEMI Registry. *Indonesian Journal of Cardiology*. 2021;42(4):109–118. Available:<https://doi.org/https://doi.org/10.30701/ijc.1194>

21. Noordzij M, Dekker FW, Zoccali C, Jager KJ. Measures of disease frequency: Prevalence and incidence. *Nephron Clinical Practice*. 2010;115(1):c17–c20. Available: <https://doi.org/https://doi.org/10.1159/000286345>

© 2023 Syauqi et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/106600>