



Risk Factors Associated with Diabetic Retinopathy among Patients Aged 50-75 Years Attending Diabetic Clinic at Mbagathi Hospital Nairobi County, Kenya

**Tom Michira Nyakaba ^{a*}, Dominic Mogere ^a, Lucina Koyio ^b
and Peterson Kariuki ^a**

^a School of Public Health, Mount Kenya University, Kenya.

^b Department of Health Services, Nairobi City County Government, Kenya.

Authors' contributions

This work was carried out in collaboration among all authors. Authors TMN, DM and LK participated in the entire research process. Author TMN participated in the data analysis process. Authors DM, PK and TMN played a key role in drafting the manuscript. All authors read, accepted and approved the final manuscript.

Article Information

DOI: 10.9734/IJTDH/2023/v44i151459

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/104697>

Original Research Article

Received: 15/06/2023

Accepted: 17/08/2023

Published: 28/08/2023

ABSTRACT

Background: According to research, nearly 60% of persons with type 1 diabetes are likely to experience diabetes retinopathy after 20 years after the initial diagnosis of diabetes type 1. Nearly 40 % of all persons with unrestrained type 2 diabetes are likely to experience diabetes retinopathy during their lifetime.

Objective: The study aimed to determine the risk factors associated with Diabetes Retinopathy among patients aged between 50 - 75 years seeking care at Mbagathi Hospital Nairobi County, Kenya.

*Corresponding author: Email: nyakabatom532@gmail.com;

Method: This study used an analytical cross-sectional study design. A systematic random sampling design was used to recruit study partakers. The sample size for this study was 151 study respondents. Both Bivariate and binary logistic regression techniques were also utilized to evaluate the degree of association between the independent and the dependent variable. Statistical significance was set at $p < 0.05$.

Results: The prevalence of diabetes retinopathy (non-proliferative diabetes retinopathy) in this study was 31.5% indicating this is a real public health concern that needs an urgent multisectoral approach. From this study, The presence of laboratory services (OR=10,95%CI=3.56-30.99), support group (OR=5.2,95%CI=1.81-14.85), provision of health care message (OR=11.6,95%CI=3.46-38.59), normal BMI (OR=3.6,95%CI=1.98-65.36) reduced the odds of diabetes retinopathy. Drinking alcohol (OR=22,95%CI=0.003-0.771), smoking (OR=33.95%, CI=0.004-0.262), uncontrolled blood sugars (OR=4,95%CI=19.89-65.36) increased the odds of diabetes retinopathy. Low education level (OR=5.9,95%CI=0.03-0.79), earning less than 6000 Ksh per month (OR=9,95%CI=0.04-0.29) smoking (OR=33.3,95%CI=0.004-0.262), uncontrolled blood sugars (OR=4,95%CI=19.89-65.36) increased the odds of diabetes retinopathy.

Conclusion: The prevalence of diabetic retinopathy was high, earning less than 6000 Ksh per month, drinking alcohol, smoking, Low education level, and having uncontrolled blood sugars increased the odds of diabetes retinopathy. The presence of laboratory services, support group, provision of health care messages, and normal BMI reduced the odds of diabetes retinopathy

Keywords: Diabetic macular edema; diabetes mellitus; diabetic retinopathy; eye complications.

ABBREVIATIONS

DM : Diabetes Mellitus, DME;
DR : Diabetic Retinopathy;
NACOSTI : National Commission for Science
Technology and Innovation;
SPSS : Statistical Package for Social
Sciences;
WHO : World Health Organization.

1. INTRODUCTION

The term Diabetes Mellitus (DM) stands for a group of metabolic disorders that are categorized by elevated levels of glucose often known as hyperglycemia in the absence of treatment [1]. Diabetes Mellitus is characterized by inadequate insulin production, poor insulin action, or both this often leads to disruption of fat, protein, and carbohydrate metabolism [2]. The most common long-term effect of diabetes includes diabetes neuropathy, retinopathy, and diabetic nephropathy among other complications [2]. The likelihood of vision loss in an individual with diabetes is up to 25 times more in non-diabetic people. People with poorly controlled diabetes are more likely to develop systemic complications, including diabetes retinopathy. Long-term diabetes management is crucial to prevent or delay the advancement of eye complications [3].

There are worrying trends of increased cases of diabetes, recent statistics from WHO there are

435 million cases of diabetes globally with a higher number of the cases being recorded in developing nations [4]. The international diabetes federation approximates that if no measures or interventions are put in place to curb the problem, it is estimated that by the year 2045, the number of diabetes cases will have increased by 48% in 2045. In recent statistics, the Age-adjusted magnitude in adults with diabetes surged from 4.7% in 1980 to 10.5% by 2019 [5]. Worldwide, the number of diabetes cases has got to an epidemic level which is worrying. Low and middle-income nations are highly affected by these emerging trends. In a report by the international diabetes federation of 2017, it's estimated there are more than 425 million in the world who have diabetes Mellitus [6]. In Africa, its approximated that 15.9 million adults are currently surviving diabetes which is a regional magnitude of 3.1% [7]. Despite diabetes being a public health concern, Africa has the most undiagnosed diabetes Mellitus cases and according to projected statistics, Africa will encounter the highest future rise in the load of DM of about 156% by 2045 [8].

The International Diabetes Federation approximates the magnitude of diabetes Mellitus in Kenya in 2015 to be 2-5%, and over 50% of cases are undiagnosed. Kenya is experiencing a rise in diabetes as a result of demographic, nutritional, and social changes such as urbanization [9]. According to the international diabetes federation, in the year 2015, the

prevalence of diabetes in Kenya was at 5% where 50% of these cases were still undiagnosed [10]. Kenya is steadily experiencing a spike in diabetes cases which has been linked to social changes such as urbanization, change in nutrition habits, and change in social demographic factors [11]. As the prevalence rises, patients face an even higher threat from long-term eye complications.

The likelihood of vision loss in a person with diabetes is 25 times higher than in non-diabetic persons. People with poorly controlled diabetes are more likely to develop systemic complications, including retinopathy [12]. Long-term diabetes management is important in delaying the progression of eye complications. Improved lipid profile and hypertension control can also slow the progression of retinopathy, especially when initiated soon after the diagnosis of diabetes [13]. Prevention measures for diabetes include dietary modification, the use of medical interventions, regular screening for Diabetes Retinopathy, and adopting a healthy lifestyle [14]. Vision loss from DR can be averted or reduced through adequate control of blood sugar, prompt diagnosis, and treatment of retinopathy. Periodic retinal examinations even when there are no signs and symptoms are key to preventing diabetes-related vision loss.

2. METHODOLOGY

2.1 Research Design

This study used an analytical cross-sectional study design. It involved figuring out how the independent variable (risk factors) and dependent variable (Diabetes retinopathy) were related to one another.

2.2 Study Area

Primary data was collected from the Mbagathi county hospital DM clinic. Mbagathi county hospital is a level five hospital located in Nairobi County, Lang'ata Constituency it serves an urban population mostly from the slum area. The health facility has over 400 patients seeking diabetes care every month [15].

2.3 Study Population

All diabetic patients who were either type 1 or type 2 aged between 50-75 years on follow-up at Mbagathi diabetic clinic presented during the study period. They were considered since they

were at high risk of developing diabetes retinopathy as compared to other age groups.

2.4 Sample Size Determination

The sample size was determined using the formula designed by Fisher as recommended by Mugenda and Mugenda (1999). As a result, 137 clients made up the final sample size for this study. Additionally, 10% of the non-respondent rate resulted in 151 study participants overall. The health facility has about 400 a point of diabetic patients every month. To determine the sampling interval the following formula was used.

$$\text{Sampling interval} = 400/151 = 3$$

To determine the initial starting point, the first participant was randomly selected.

2.5 Sampling Technique

Mbagathi county hospital was purposively selected, A systematic random sampling design was used to recruit study partakers in this study where all diabetic clients aged 50 to 75 years were carefully selected to participate in the study.

2.6 Data Collection Method and Instruments

A semi-structured questionnaire form was used to capture the data information required in this study. Section A of the questionnaire captured data on patients' factors associated with Diabetes Retinopathy, section B captured data on Diabetes Retinopathy screening, section C captured data on lifestyle factors and lastly, section D captured data on health facility factors associated with Diabetes Retinopathy.

2.7 Validity and Reliability

Validity refers to the correctness of the research instruments which promotes the collection of accurate data and information. An expert was requested to appraise the data-gathering tools, and supervisors and a diabetologist was requested to review the data-gathering tools before embarking on the data-collection process. The degree to which data collection is reliable provides the same finding after being subjected to the study participants than ones. Carefully selected study participants were used to carry out the reliability of the study tools. Cronbach alpha reliability test was carried out using SPSS version 26, literature accepts an internal consistency of above 0.7. According to the

findings of the reliability test, a score of 0.94 was obtained, which confirmed that the tools were reliable for data collection.

2.8 Data Processing and Analysis

Using Microsoft Excel, the collected data was cross-checked and verified for accuracy and completeness to prevent double entry before being entered into the Statistical Package for Social Sciences (SPSS) version 26 for analysis. Data was analyzed using descriptive statistics and presented as tables and percentages. The χ^2 test for independence was utilized in the bivariate analysis. A p-value of <0.05 was used as a cutting threshold for significant variables. Significant variables in the bivariate analysis were imported into the binary logistic regression. The variables that were significantly associated with DR were identified using binary logistic regression.

3. RESULTS

3.1 Prevalence of Diabetes Retinopathy

As indicated in Fig. 1, both eye clinical examination and fluorescein angiography were used to diagnose diabetes retinopathy (non-proliferative diabetes retinopathy). These procedures were conducted by competent and licensed clinical officers and medical doctors, from this study (31.5%) of the study partakers were diagnosed with diabetes retinopathy, while (68.5%) of the study participants didn't have diabetes retinopathy.




Fig. 1. Prevalence of Diabetes Retinopathy

3.2 Descriptive Statistics on Social Demographic Factors

As indicated in Table 1, more than half (53%) of the study respondents were males. Regarding age, close to half (47%) of the study participants

were aged between age 50-55 while only a few (16.1%) of the study respondents were aged 66-75 years. Concerning marital status, the majority (61.7%) of the study respondents were married. Regarding the level of education, close to half (49.7%) of the study participants in this study had attained a secondary level of education. Only a few (12.8%) of the respondents in this study had attained the tertiary level of education. In this study, the majority (82.6%) of the study participants were Christians as compared to Muslims who accounted for (17.4%). Finally, concerning the level of income, the majority (61.1%) of the study respondents reported earning a monthly income of above 6000 Ksh.

3.3 Association between Social Demographic Factors and Diabetes Retinopathy

In the social demographic factors; gender type ($\chi^2=6.256, df=1, p=0.012$), marital status ($\chi^2=10.36, df=4, p=0.035$), an education level ($\chi^2=35.707, df=3, p=0.000$), and income earned per month ($\chi^2=32.244, df=1, p=<0.000$), were statistically associated with diabetes retinopathy hence these variables were imported to binary logistic regression for further analysis. Type of religion ($\chi^2=1.690, df=1, p=0.194$) and age ($\chi^2=1.091, df=2, p=0.579$) were found not to be statistically associated with diabetes retinopathy. As indicated in Table 2, compared to diabetic patients with tertiary education, those with primary education had a 5.9 higher risk of developing diabetes retinopathy. From this study, study participants who earned less than 6000 Ksh per month had a 9-fold higher risk of developing diabetic retinopathy than their matched peers.

3.4 Association between Health System Factors and Diabetes Retinopathy

In the health system factor associated with diabetic retinopathy, availability of diabetes mellitus medication ($\chi^2=9.796, df=1, p=0.002$), availability of diabetes mellitus laboratory testing services ($\chi^2=54.022, df=1, p=<0.000$), presence of diabetes mellitus support groups ($\chi^2=21.102, df=1, p=<0.000$), and the provision of diabetes mellitus healthcare messages ($\chi^2=46.309, df=1, p=<0.000$) were found to be associated with diabetes retinopathy, as a result, these variables were imported in the binary logistic regression for further analysis. Adherence to diabetes mellitus

guidelines ($X^2=0.682, df=1, p=0.409$) was not associated with diabetes retinopathy. As indicated in Table 3, study participants who reported that laboratory test services (blood sugar monitoring and blood pressure monitoring) were available were 10 times less likely to be diagnosed with diabetes retinopathy as compared to their fellow counterparts. In

addition, participants in diabetes mellitus support groups had a 5.2-fold lower risk of developing diabetic retinopathy than those who didn't. Study respondents who reported receiving a healthcare message related to diabetes Mellitus were 11.6 less likely to experience diabetic retinopathy than their fellow counterparts.

Table 1. Social demographic characteristics of the study respondents

Independent Variables	Categories	Frequency	Valid Percentage%
Gender	Male	79	53
	Female	70	47
Age	50-55	70	47
	56-65	55	36.9
	66-75	24	16.1
Marital Status	Single	29	19.5
	Married	92	61.7
	Divorced	9	6
	Separated	11	7.4
	Widowed	8	5.4
Educational Level	Never been to school	23	15.4
	Primary level	33	22.1
	Secondary level	74	49.7
	Tertiary level	19	12.8
Religion	Christian	123	82.6
	Religion	26	17.4
Income level	6000ksh and below	58	38.9
	6000 ks and above	91	61.1

Table 2. Multivariate analysis of social demographic factors

Step 1a	B	S.E	Wald	Df	Sig	Exp(B)	95% C.I for EXP(B)		
							Lower	Upper	
Marital status			6.634	4	0.157				
	Single	2.12	1.08	3.888	1	0.049	8.331	1.01	68.53
	Married	2.36	1.00	5.59	1	0.018	10.644	1.45	75.60
	Divorced	2.31	1.33	3.04	1	0.08	10.09	0.75	135.68
	Separated	1.35	1.27	1.19	1	0.28	3.99	0.33	47.90
	Widowed					ref			
Education level			18.55	3	0.000				
	Never been to school	-0.03	0.829	0.001	1	0.975	0.975	0.192	4.994
	Primary	-1.75	0.769	5.17	1	0.02	0.17	0.03	0.79
	Secondary	0.87	0.74	1.39	1	0.23	2.38	0.56	10.06
	Tertiary					ref			
Gender			1.69	1	0.19	0.53	0.21	1.37	
	Income below 6k	-2.19	0.486	20.32	1	0.000	0.112	0.043	0.290
	Income above 6k					ref			
Constant			0.027	1	0.87	1.21			

3.5 Association between Lifestyle Factors and Diabetes Retinopathy

In the lifestyle factors associated with diabetic retinopathy, smoking ($X^2=28.575, df=1, p<0.000$), drinking alcohol ($X^2=45.928, df=1, p^*<0.000$), body mass index ($X^2=98.485, df=3, p^*<0.000$), physical activities (both aerobic and anaerobic physical activities) ($X^2=32.693, df=1, p<0.000$) were found to be statistically significant with diabetes retinopathy hence they were imported for logistic regression analysis. Controlled blood pressure ($X^2=0.358, df=1, p=0.55$) and period lived with diabetes mellitus ($X^2=2.161, df=1, p=0.142$) were found not to be associated with diabetes mellitus. As indicated in Table 4, When compared to their peers, study participants who

drank alcohol were 22 times more likely to develop diabetes retinopathy. Study participants who were smokers were 33.3 times more likely to be diagnosed with diabetes retinopathy as compared to their fellow counterparts. When compared to obese study participants, those with a normal body weight had a 3.6 times lower risk of developing diabetes retinopathy. In addition, study participants with uncontrolled blood sugars (persistently high blood sugar levels despite the presence of insulin and oral glycemic control agents) were 17 more likely than those with controlled blood sugars (presence of normal blood sugars after administration of insulin or oral glycemic control agents) to develop diabetes retinopathy.

Table 3. Multivariate analysis of health system factors

Step 1a	Variables	B	S.E	Wald	Df	Sig	Exp(B)	95% C.I for EXP(B)	
								Lower	Upper
	Availability of medicines	0.40	0.57	0.499	1	0.48	1.49	0.49	4.53
							ref		
	Lab test available	2.35	0.55	18.12	1	0.00	10.49	3.56	30.99
							ref		
	Support group	1.65	0.54	9.43	1	0.002	5.19	1.81	14.85
							ref		
	Receiving a message on DR	2.45	0.62	15.82	1	0.00	11.56	3.46	38.59
							ref		
	Constant	-2.23	0.52	18.85	1	0.00	0.10		

Table 4. Multivariate analysis on life style factors

Step 1a	Variables	B	S.E	Wald	Df	Sig	Exp(B)	95% C.I for EXP(B)	
								Lower	Upper
	Drinking alcohol	-3.09	1.44	4.58	1	0.03	0.046	0.003	0.771
							ref		
	smoking	-3.45	1.08	10.27	1	0.00	0.03	0.004	0.262
							ref		
	BMI			19.74	3	0.00			
	<18.5	5.66	1.73	10.69	1	0.00	2.86	9.651	85.085
	18.-24.9	5.88	1.48	15.86	1	0.00	3.65	19.887	65.356
	25-29.9	1.73	0.94	3.34	1	0.07	5.61	0.883	35.665
	>30						ref		
	Blood sugar under control(4.5-7.2 mmol/L)	2.85	0.97	8.59	1	0.00	17.33	2.571	116.794
							ref		
	Physical exercise	1.27	0.896	2.02	1	0.16	3.57	0.617	20.680
							ref		
	Constant	-2.43	0.95	6.55	1	0.01	0.09		

4. DISCUSSION

In this study, both eye clinical examination and fluorescein angiography were used to diagnose diabetes retinopathy, from this study (31.5%) of the study partakers were diagnosed with diabetes retinopathy. These findings were close to a study done in Australia which documented a prevalence of 34.6% among study participants [16]. Compared to diabetic patients with tertiary education, those with primary education had a 5.9 higher risk of developing diabetes retinopathy. The need for better health-seeking behavior becomes more and more understood as one's education level rises. These findings were in agreement with another study done in China where being educated reduced the odds of having diabetes retinopathy [17].

Study participants who were earning less than 6000 Ksh per month had a 9-fold higher risk of developing diabetes retinopathy than their matched peers. The probable reason for this is that better income promotes the affordability of various health services related to diabetes management. These findings were in agreement with another study done in Ethiopia [18]. Study participants who reported that laboratory test services were available were 10 times less likely to be diagnosed with diabetes retinopathy as compared to their fellow counterparts. This means the adequate provision of laboratory test services enables earlier detection of parameters that are likely to contribute to diabetes retinopathy, some of the essential services provided include blood sugar monitoring and blood pressure monitoring which are risk factors for diabetes retinopathy. This study's results were in harmony with another study done in Amhara northwest of Ethiopia [19].

In addition, participants in diabetes mellitus support groups had a 5.2-fold lower risk of developing diabetes retinopathy than those who didn't. The presence of support groups ensures patients who have diabetes receive maximum care and treatment which reduces the odds of developing diabetes-related complications. Some of the essential services provided by this support group include support to purchase diabetes medication, provision of regular health education, and promotion services related to diabetes management. These findings were in harmony with another study done in India [20]. Study respondents who reported receiving a healthcare message related to diabetes Mellitus were 11.6 less likely to experience diabetes retinopathy

than their fellow counterparts. The provision of healthcare messages serves protective factor since it reminds diabetes patients to observe preventive measures as well as the need to comply with the prescribed regimen. These findings were in harmony with another study done in Germany [21].

When compared to their peers, study participants who drank alcohol were 22 times more likely to develop diabetes retinopathy. Drinking of alcohol among diabetes patients has been observed to be a risk factor for diabetes mellitus and its associated complication which include DR. These findings were in harmony with a study in Sweden [22]. Study participants who were smokers were 33.3 times more likely to be diagnosed with diabetes retinopathy as compared to their fellow counterparts. Smoking has been linked to an increased risk of developing diabetes and its complications such as DR. These findings agreed with another study carried out in Russia [23]. When compared to obese study participants, those with a normal body weight had a 3.6 times lower risk of developing diabetes retinopathy. Being overweight and obese have been researched and have been recognized to be risk factors for diabetes complications such as diabetes retinopathy, These findings were in agreement with a study done in China [24]. Lastly, Study participants with uncontrolled blood sugars were 17 more likely than those with controlled blood sugars (4.5-7.2mmol/L) to develop diabetes retinopathy. In this study, uncontrolled blood sugar was considered a persistently high blood sugar despite the provision of insulin injections or oral agents for control purposes. Diabetes retinopathy is more likely to occur as a result of damaged nerves and blood vessels in the eye caused by elevated levels of sugar in the blood. These findings were in agreement with another study done in Cameroon [25].

5. CONCLUSION

The present investigation's prevalence of diabetes retinopathy was 31.5%, which was high indicating this is a public health concern requiring urgent interventions. In the social demographic factors, earning less than 6000 Ksh per month, and having a primary level of education increased the odds of having diabetes retinopathy. In the health system factors associated with diabetes retinopathy, the provision of laboratory testing services, the presence of a diabetes mellitus support group,

and the provision of diabetes mellitus health messages reduced the odds of diabetes retinopathy. In the lifestyle factors, drinking alcohol, smoking, and the incidence of uncontrolled blood sugars augmented the odds of diabetes retinopathy while having a normal body mass index reduced the odds of having diabetes retinopathy. There is a need for joint action by concerned stakeholders to reduce the risk of diabetes retinopathy among diabetic patients.

ETHICAL APPROVAL AND CONSENT

Ethical approval was sought from the MKU institutional and ethical review committee (ERC). An introduction letter was sought from MKU graduate school. NACOSTI granted the license to conduct the study. Additional approval was requested from the appropriate county offices (County Department of Health, County health committee, and HR Department). Consent forms were provided to the study participants before embarking on data collection. Participation in this study was purely voluntary. Study respondents' confidentiality and anonymity were highly upheld. Those who were found to be positive for DR were referred to an eye expert for appropriate treatment.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. WHO. "World Health Statistics 2016: Monitoring health for the SDGs, sustainable development goals; 2016.
2. Lima VC, Cavalieri GC, Lima MC, Nazario NO, Lima GC. "Risk factors for diabetic retinopathy: A case-control study." *Int. J. Retin. Vitr.* 2016;2(1):1–7.
DOI:10.1186/S40942-016-0047-6/TABLES/2
3. Tan F, et al. "Associated risk factors in the early stage of diabetic retinopathy." *Eye Vis.* 2019;6(1):1–10.
DOI:10.1186/S40662-019-0148-Z/FIGURES/3
4. Kajiwara A, et al. "Gender differences in the incidence and progression of diabetic retinopathy among Japanese patients with type 2 diabetes mellitus: a clinic-based retrospective longitudinal study." *Diabetes Res Clin Pr.* 2014;103(3):7-10.
DOI: 10.1016/j.diabres.2013.12.043
5. Ejigu T, Tsegaw A. "Prevalence of diabetic retinopathy and risk factors among diabetic patients at university of gondar tertiary eye care and training center, North-West Ethiopia." *Middle East Afr. J. Ophthalmol.* 2021;28(2):71–80.
DOI: 10.4103/MEAJO.MEAJO_24_21
6. Penman A, et al. "Risk factors for proliferative diabetic retinopathy in african americans with type 2 diabetes." *Ophthalmic Epidemiol.* 2016;23,(2):88.
DOI: 10.3109/09286586.2015.1119287
7. Carpineto P, et al. "Neuroretinal alterations in the early stages of diabetic retinopathy in patients with type 2 diabetes mellitus." *Eye.* 2016;30(5)673–679.
DOI: 10.1038/eye.2016.13
8. Burgess PI, MacCormick IJC, Harding SP, Bastawrous A, Beare NAV, Garner P. "Epidemiology of diabetic retinopathy and maculopathy in Africa: A systematic review." *Diabet. Med.* 2013;30(4):399.
DOI: 10.1111/J.1464-5491.2012.03756.X
9. Mathenge W, et al. "Prevalence and correlates of diabetic retinopathy in a population-based survey of older people in Nakuru, Kenya." *Ophthalmic Epidemiol.* 2014;21(3):169–177.
DOI: 10.3109/09286586.2014.903982.
10. Nyawira G, Samuel W, Maricianah O, Elizabeth AB, Kenneth N. "Determinants of antiretroviral therapy adherence among older adolescents living with HIV in Kenya during the transition to adult care; An observational study." *J. AIDS HIV Res.* 2020;12(2):24.
DOI: 10.5897/JAHR2020.0513.
11. Magliah SF, Bardisi W, Attah MAI, Khorsheed MM. "The prevalence and risk factors of diabetic retinopathy in selected primary care centers during the 3-year screening intervals." *J. Fam. Med. Prim. Care.* 2018;7(5):975.
DOI: 10.4103/JFMPC.JFMPC_85_18
12. Hussain S, Qamar MR, Iqbal MA, Ahmad A, Ullah E. "Risk factors of retinopathy in type 2 diabetes mellitus at a tertiary care hospital, Bahawalpur Pakistan." *Pak J Med Sci.* 2013;29(2):536–539.
DOI: 10.12669/pjms.292.3066
13. Sahiledengle B, et al. "Prevalence and factors associated with diabetic retinopathy among adult diabetes patients in southeast

- ethiopia: A hospital-based cross-sectional study." Clin. Ophthalmol. 2022;16:3527–3545.
DOI: 10.2147/OPHTH.S385806
14. Kawasaki R, et al. "Risk of cardiovascular diseases is increased even with mild diabetic retinopathy: the Japan diabetes complications Study." Ophthalmology. 2013;120(3):574–82.
DOI: 10.1016/j.ophtha.2012.08.029
15. KNBS. "A survey report; 2019.
16. Yau JWY, et al. "Global prevalence and major risk factors of diabetic retinopathy." Diabetes Care. 2012;35(3):556–564.
DOI: 10.2337/DC11-1909
17. Khalaf FR, et al. "Does a diabetic retinopathy educational program raise awareness among elderly diabetic patients?." Diabetes, Metab. Syndr. Obes. Targets Ther. 2019;12:1867.
DOI: 10.2147/DMSO.S208072.
18. Mersha GA, Alimaw YA, Woredekal AT. "Prevalence of diabetic retinopathy among diabetic patients in Northwest Ethiopia—A cross sectional hospital based study." Plos One, 2022;17(1):e0262664.
DOI: 10.1371/JOURNAL.PONE.0262664
19. Zegeye AF, Temachu YZ, Mekonnen CK. "Prevalence and factors associated with Diabetes retinopathy among type 2 diabetic patients at Northwest Amhara comprehensive specialized hospitals, Northwest Ethiopia 2021." BMC Ophthalmol. 2023;23(1):1–8.
DOI:10.1186/S12886-022-02746-8/TABLES/3
20. Gudlavalleti A, et al. "Establishing peer support groups for diabetic retinopathy in India: lessons learned and way ahead." Indian J. Ophthalmol. 2020;68(13):S70–S73.
DOI: 10.4103/IJO.IJO_1928_19
21. Silverberg EL, Sterling TW, Williams TH, Castro G, de la Vega PR, Barengo NC. "The association between social determinants of health and self-reported diabetic retinopathy: An exploratory analysis." Int. J. Environ. Res. Public Heal. 2021;18(2):792-792.
DOI: 10.3390/IJERPH18020792.
22. Lee CC, et al. "Association between alcohol consumption and diabetic retinopathy and visual acuity-the AdRem Study." Diabet. Med. 2010;27(10):1130–1137.
DOI: 10.1111/J.1464-5491.2010.03080.X
23. Koseva D, Zlatarova Z, Kaneva M. "Is there an association between smoking and diabetic retinopathy?" J. Union Sci. - Varna. Med. Ecol. Ser. 2020;25(2):12–18.
DOI: 10.14748/ISUVSME.V25I1.7191
24. Zhu W, Wu Y, Meng YF, Xing Q, Tao JJ, Lu J. "Association of obesity and risk of diabetic retinopathy in diabetes patients: A meta-analysis of prospective cohort studies." Medicine (Baltimore). 2018;97(32).
DOI: 10.1097/MD.00000000000011807.
25. Jingi AM, Tankeu AT, Ateba NA, Noubiap JJ. "Mechanism of worsening diabetic retinopathy with rapid lowering of blood glucose: The synergistic hypothesis." BMC Endocr. Disord. 2017;17(1):1–4.
DOI: 10.1186/S12902-017-0213-3/PEER-REVIE

© 2023 Nyakaba et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/104697>