



Phytoremediation Indices of Water Hyacinth (*Eichhornia crassipes*) Growing in Panteka Stream, Kaduna, Nigeria

Victoria Moltong Yilwa ^a, Nwankwo Cornelius Tochukwu ^{a,b*},
Emere Matthew Chika ^a, Adejo Peter Ojodale ^{a,b}
and Danfulloh Tundeno Barde ^{a,b}

^a Department of Biological Sciences, Nigerian Defence Academy, Kaduna, PMB 2109, Nigeria.

^b Department of Biotechnology, Nigerian Defence Academy, Kaduna, PMB 2109, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJOB/2023/v19i3369

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/107968>

Original Research Article

Received: 24/08/2023

Accepted: 31/10/2023

Published: 07/11/2023

ABSTRACT

The aquatic environment is incessantly polluted by the release of high toxic concentrations of heavy metals which are bio-accumulative and persistent in nature. This investigation was conducted to assess the phytoremediation potential of water hyacinth (*Eichhornia crassipes*) growing in Panteka stream, Kaduna where mechanic and farming activities are carried out. Having three sampling points (A, B and C) and pond water, where farming is predominant served as the control site (D) using phytoremediation indices. The root and shoot samples of *E. crassipes* grown at the sampling points (A, B, C) and the control were analyzed to determine heavy metal concentrations of

*Corresponding author: E-mail: corneliustochukwu173@gmail.com;

cadmium (Cd), chromium (Cr), copper (Cu), nickel (Ni), lead (Pb), and zinc (Zn) through Atomic Absorption Spectrophotometry (AAS). The phytoaccumulation indices were calculated via bioaccumulation coefficient and translocation factor. The results showed that the root samples had higher accumulation of heavy metals than the shoot samples. During the analysis the heavy metal Zn was noted to be accumulated the highest in roots and shoots (335.32 ± 23.6 and 256.52 ± 30.82) mg/kg at sampling point B respectively. In this present analysis heavy metals were translocated efficiently and had bioaccumulation coefficient and translocation factor greater than one. Nickel had the highest bioaccumulation coefficient and translocation factor, and Zn had a translocation factor that was less than one across all samples. This implies that *E. crassipes* is a potential hyperaccumulator plant for phytoremediation.

Keywords: *Eichhornia crassipes*; heavy metals; phytoremediation and phytoremediation indices.

1. INTRODUCTION

One of the key priorities of this present age is the preservation of the environment's quality. The biosphere is being degraded as a result of the emission of natural and man-made substances that can harm living organisms. Heavy metals, among all pollutants, are easily transported and accumulated in the environment [1]. Water body contamination increases at an alarming rate by human activities such as the release of pesticides from agricultural sectors, radionuclides and hydrocarbons from petrochemical industry, and hazardous metals from untreated wastewater are all instances of pollutants. Metals, unlike organic trash, are not biodegradable. Arsenic, Cadmium, Lead, Manganese, Mercury, Selenium and Zinc are examples of trace and heavy metals that must be eradicated from the environment [2]. Once heavy metals have contaminated the environment, they will pose a long-term threat to humans and animals [3]. Biological decontamination approaches, such as phytoremediation, are thought to be safe for eliminating these toxins from the environment, especially from water and soil.

A green technology known as phytoremediation uses plants to clean up contaminated areas or to dislodge pollutants from the environment. It is an environmentally benign and affordable technology [4]. On contaminated terrain, many plant species may thrive, and some may even accumulate significant amounts of heavy metals in their tissues. Hyper-accumulator plants with the capacity to develop promptly and accumulate large metal levels are required to facilitate phytoremediation [5]. Currently known metal hyper-accumulator plant species number over four hundred [6].

Phytoextraction, phytostabilization, rhizofiltration, and phytovolatilization are some of the

phytoremediation approaches for treating metal-contaminated environments [7,8]. Phytoextraction refers to the absorption of metals, coupled with subsequent translocation and accumulation in vegetative organs, whereas phytostabilization refers to the hazardous ions that remain immobilized in the contaminated medium [9-12]. As a result, heavy metals become stable in the rhizosphere (horizontal underground stem region in the plant). The roots of phytoremediating plants absorb, concentrate or discharge contaminants out of polluted effluents during the rhizofiltration process [13]. Harvesting plant tissues that has accumulated hazardous metals is required for phytoextraction and rhizofiltration. The plant material is normally burned, and the ash generated can be utilized to recycle metal [14,7]. The creation of "eco-catalysts" for the chemical sectors is a recent method of utilizing biomass that has been gathered from plants that highly accumulate metal [15,16]. Volatilization is another method that plants use to remove inorganic contaminants. Metals undergo biological conversion to gaseous states, which are then discharged into the air during this process [17].

Recent strategy for phytoextraction of heavy metal contaminated soil includes the use microbial-assisted phytoremediation which is a potential approach, where a broad range of plant species coexist closely with arbuscular mycorrhizal fungi (AMF), promoting healthy growth and resilience to contaminants. AMF inhabit plant roots and expand their hyphae within the rhizosphere, helping the plants take up nutrients and minerals, control the buildup of heavy metals and stress tolerance [18]. [19], carried out a study using arbuscular mycorrhizal fungi to aid phytoextraction. Arbuscular mycorrhizal fungi (AMF) were used in this work to enhance the growth of *Zea mays* L. in heavy metal-rich tannery slush [20].

Eichhornia crassipes is an aquatic macrophyte belonging to the class Liliopsida and the family Pontederiaceae. It originated in tropical South America, but it can now be found in Africa, Australia, India, and a variety of other places. This plant may be used as animal feed and has a fast growth rate as well as the ability to withstand various forms of pollution [21]. An excellent plant for phytoremediation has a high biomass, high tolerance to heavy metals stress, and a high capacity for metal accumulation [22]. The said plant has these qualities; therefore, it is well-known for their ability to absorb toxic metals from water. Water hyacinth, as a good and capable hyper-accumulator plant, can accumulate a very high level of heavy metals [23].

The sole aim of this investigation was to determine the heavy metals content of the native aquatic macrophyte (*E. crassipes*) in Panteka stream and its ability to accumulate heavy metals and potency for phytoremediation. As well as, to ascertain more information on the heavy metal status of water bodies and *E. crassipes* which was seen as noxious species.

2. MATERIALS AND METHODS

2.1 Study Area

The study area is the stream that has its entry point from Rafin Guza, and is flowing through the Panteka mechanic village, located between latitudes 10° 32'37" and 10° 33'56" N and longitude 7° 24'29" and 7° 25'7" E, at the Northern part of Kaduna, Nigeria. The Panteka stream flows westward, it usually decreases in volume during dry season and *Eichhornia crassipes* grows in the stream. Similarly, Panteka is a mechanic village where all kinds of cars and motorcycle spare parts are sold, and their maintenance is carried out. Alongside this, agricultural activities take place in this area and runoff from the farms and debris washed off from these anthropogenic activities mentioned above enters the stream. The control samples were collected from an enclosed pond, where farming activities are carried out, behind Mal. Abdulrakeem Fish Farm, Farin-Gida Mando Kaduna. The stream was divided into three sampling points, A, B, and C. Sampling point A is 4,218m away from B, and B is 4000m away from C. At sampling point A, the predominant activity done in this area is farming, dumping of car worn-out parts and repairs, while at sampling points B and C, there is more of mechanic repair activities.

2.2 Plant Sample Collection

Plant samples were collected from Panteka's contaminated water and the control samples from the enclosed pond. Plant samples from the three sampling points and the control were wrapped in labeled polybags for each of the plants and brought to the Biological Sciences Department Laboratory Nigerian Defence Academy, Kaduna.

2.3 Plant Sample Processing and Analysis

The plant samples were washed to remove the unwanted debris from the plant with deionized water. Followed by herbarium techniques (which include pressing and drying the plant sample using the plant presser, labeling, and storage) and full authentication of the plant. The plant samples were identified and authenticated in the herbarium, Botany laboratory NDA. The specimen with voucher number NDABIOH202030 was kept in the herbarium for reference purposes.

2.4 Water Sample Collection

In both dry (April 2020) and wet (August 2020) seasons, water was sampled twice at early hours of the morning in sterile plastic containers from the four sampling points. Using the method of collecting water grab samples as described by Thomas [24], and was done by holding the uncapped sterile plastic container upside down and submersed it. And, the tip of the container upright and then allowed water to fill the container and was removed from the water, screw-on cap and carefully labeled respectively. These water samples were subjected to laboratory investigation for heavy metals.

2.5 Heavy Metal Analysis of Water

To determine the heavy metals in water samples, [25] approach was employed. 50ml of the water samples from each sampling point for each of the seasons were digested with 10ml of HNO₃ on a hot plate. The resulting clear solutions were filtered using Whatman filter papers and reconstituted to 50ml in volumetric flasks with deionized water. 50ml of the filtrates were analyzed for the heavy metals using Atomic Absorption Spectrophotometer (Model: PG-990). The reference solutions of metals to be assayed were prepared and atomic absorption spectrophotometer was allowed to run for 10




Fig. 1. Panteka stream study area map with sampling points

(Source: GIS, KADSU)

minutes before immersing the sipper into the reference metal solutions to calibrate the AAS and acetylene was used as the carrier gas. Flame absorption method at the following conditions; Cadmium (wavelength 228.8nm, slit 0.4nm, high voltage of 315.0V and 2.0mA lamp current), Chromium (wavelength 357.9nm, slit 0.4nm, high voltage of 287.50V and 4.0mA lamp current), Copper (wavelength 324.7nm, slit 0.4nm, high voltage of 283.50V and 3.0mA lamp current), lead (wavelength 283.3nm, slit 0.4nm, high voltage of 370V and 2.0mA lamp current), Nickel (wavelength 232.0nm, slit 0.2nm, high voltage of 420.0V and 4.0mA lamp current) and Zinc (wavelength 213.9nm, slit 0.4nm, high voltage of 339.75V and 3.0mA lamp current) were utilized in the process respectively. The absorbance was read before placing the sipper into the solution of the digested samples and the concentrations were taken from the AAS.

2.6 Heavy Metals Analysis of Plant

The method described by Shaayau and Garba [26] was adopted to analyze heavy metals from the plant samples. Exactly 0.25g of each of the

grounded plant samples (roots and shoots) were weighed after which it was placed in a 100ml beaker and digested with 30ml of mixed acid containing the following; 650ml, 80ml, and 20ml of concentrated, Trioxonitrate (v) acid (HNO₃), Perchloric acid (HClO₄) and Tetraoxosulphate (vi) acid (H₂SO₄) respectively on a hot plate. Following completion of the digestion, samples were cooled, made up to 50ml with distilled water, filtered using Whatman filter paper in volumetric flasks, and transferred into a sterile labeled plastic container. Using an Atomic Absorption Spectrophotometer (Model: PG-990), 50ml of the filtrates were utilized to determine heavy metal contents [27].

2.7 Characterization of Phytoremediation Potential

Bioaccumulation Coefficient (BAC): Is the plant/water concentration quotient [28]. It is calculated as follow;

$$BAC = C_{\text{root}} / C_{\text{water}}$$

$$BAC = C_{shoot} / C_{water}$$

Where C_{root} = Concentration of heavy metal in *E. crassipes* root (mg/kg), C_{shoot} = Concentration of heavy metal in *E. crassipes* shoot (mg/kg) and C_{water} = Concentration of heavy metals in water (mg/kg).

Translocation Factor (TF): The translocation from shoot to root was determined by translocation factor, which is given by:

$$TF = \frac{\text{Metal Concentration in the shoot (mg/kg)}}{\text{Metal Concentration in the root (mg/kg)}}$$

2.8 Statistical Analysis

To confirm the variability and validity of the results, One-way analysis of variance (ANOVA) and the T-Test were used to analyze the study's data. Using the Statistical Package for Social Sciences (SPSS-version 23.) and Excel version 2016, all data were presented as LSD and given a significance level of $p < 0.05$.

3. RESULTS AND DISCUSSION

3.1 Heavy Metals Concentration in the Study Area

The levels of these heavy metals in samples of water across the wet season from sampling points A, B, C, and control increased except those of Zinc, Lead and Copper which were also, of increased level during the dry season. This might be due to the runoff from farm lands and debris washed off from other anthropogenic activities depositing more of these heavy metals, and the increase in Zinc and Lead at sampling point A and B, during the dry season might be due to the intense deposits of engine oils, wastes from car parts and other items coated with zinc not being washed off in the stream during the wet season and those deposited by wind during the dry season. Among the heavy metals analyzed at Panteka Stream during the wet season, Cadmium was observed to have the highest concentration (0.78mg/kg) and this was more than the maximum concentration standards of FEPA (0.01mg/L) and USEPA (0.01mg/L). The high content of these heavy metals might be as a result of the release of effluents from car paints, car batteries, and phosphate fertilizers in the stream [29]. The heavy metals concentration conforms with the findings of Ganesh [30] who reported that heavy metals were observed from

paper dumping sites. Cadmium concentration in the water samples was the highest among the heavy metals studied. However, it was discordant with the reports of Astatkie et al. [31], who in their study reported highest concentration of heavy metal Lead in water collected from the Awetu watershed stream. However, the other heavy metals analyzed were within the recommended limits.

3.2 Heavy Metal Concentration in the Roots and Shoots of *E. crassipes*

Heavy metals concentration was entirely higher in roots than in shoots. Different plant components metals accumulation indicated heavy metals in varying concentrations, at the sampling points as shown in Tables 1 and 2. There was more accumulation of Zinc in root and shoot at sampling point B (335.32 ± 23.60 and 256.52 ± 30.82) mg/kg respectively. Other heavy metals accumulated significantly in *E. crassipes* growing at all the sampling points except in the control samples (Pb and Zn high in control). The results were in line with the findings of Veschasit et al. [32], who studied heavy metals in aquatic plant species *Ipomoea aquatica* and *Neptunia oleracea* growing in Tha Chin River, where they found out that all metals accumulated higher in the root than in the shoot. Also, [33], observed the same higher concentration of the Cadmium in the root than in the shoot. Ganesh [30] recorded a higher accumulation of heavy metals in root than in shoot of *E. crassipes* growing at paper dumping site and further stated that, in terms of plant components, the ratio of metal concentrations does not appear to change in a way that is reliant on exposure. Hence, there was a substantial accumulation of heavy metals in roots and shoots of *E. crassipes* at Panteka stream ($P < 0.05$).

3.3 Correlation of Heavy Metals, Roots and Shoots, and the Seasons of Sampling

The roots and shoots of *E. crassipes* and concentration of heavy metals in the water samples revealed a positive correlation as shown in Table 3. This agrees with the reports of Veschasit et al. [32], who reported a significant relationship between the native aquatic plant species *Ipomoea aquatica* and *Neptunia oleracea* that were found to accumulate high concentrations of the same heavy metals. Statistical analysis revealed that copper in

I. aquatica had a positive correlation with copper in the surrounding water in their report. Similarly, the total concentration of heavy metals in the root and shoot samples of *E. crassipes* increased in conjunction with the total concentration of these metals in the water, indicating a positive correlation. In addition, heavy metals at sampling points increased as these metals increased during the dry and wet seasons.

3.4 Bioaccumulation Coefficient and Translocation Factor of *E. crassipes*

To assess *E. crassipes'* capability for phytoremediation of heavy metals, two phytoremediation indices were applied. These include Bioaccumulation Coefficient (BAC) and Translocation Factor (TF). Bioaccumulation Coefficient (BAC) was used to indicate the degree of enrichment of heavy metal in the

plants relative to that in its habitat. Nickel showed the highest and Copper the least bioaccumulation coefficient in the root and shoot samples of *E. crassipes*. This stood in accordance with the study of [34]. Who in their study reported the highest bioaccumulation coefficient in Nickel in *Salvinia molesta* followed by Mn, Fe, Zn, Pb, Co, Cu, Cr and Cd.

Al-Farraj et al. [35] reported that Fungi like *Aspergillus niger* grow on roots of water hyacinth and can grow on an alkaline media (pH greater than 7) that can adsorb metals. The adsorption of metals onto the roots of water hyacinth may have been possible due to *A. niger* as was reported by Mahmood et al. [36]. In this study, the bioaccumulation coefficient was greater than one for all the heavy metals. This might be due to the high accumulation in root and shoot of *E. crassipes*.




Fig. 2. Mean Concentration of heavy metals at the sampling points during dry and wet seasons




Fig. 3. Mean Bioaccumulation Coefficient of Heavy Metals in the roots of *E. crassipes*. The descending order of bioaccumulation coefficient in the roots from maximum to the minimum at the sampling points is as follows Ni>Cr> Pb>Zn>Cd>Cu

Table 1. Heavy metal concentration in root samples

Sampling points	Metals (mg/kg)					
	Cd	Cr	Cu	Ni	Pb	Zn
A	27.65 ± 30.06 ^a	47.20 ± 21.69 ^a	36.88 ± 22.09 ^a	50.90 ± 16.48 ^a	55.05 ± 61.88 ^a	224.45 ± 30.24 ^a
B	54.76 ± 13.49 ^a	57.96 ± 13.11 ^a	127.12 ± 19.45 ^a	34.52 ± 11.88 ^{ab}	BDL	335.32 ± 23.60 ^b
C	46.88 ± 11.27 ^a	163.68 ± 129.62 ^a	90.72 ± 7.54 ^b	31.20 ± 8.11 ^{ab}	23.32 ± 49.69 ^a	195.56 ± 22.10 ^a
Control	BDL	12.95 ± 3.34 ^a	0.70 ± 3.70 ^a	15.00 ± 5.35 ^b	72.45 ± 7.99 ^a	227.70 ± 40.87 ^a

Values are expressed as mean ± standard error mean. Means with different superscripts down the column are significantly different ($P < 0.05$) at 95% confidence level using least significance difference (L.S.D), BDL-below detectible limit. The descending order of the heavy metal's accumulation in the root samples from the maximum to the minimum at the sampling points is; Zn>Cr>Cu>Pb>Cd>Ni

Table 2. Heavy metal concentration in shoot samples

Sampling points	Metals (mg/kg)					
	Cd	Cr	Cu	Ni	Pb	Zn
A	29.70 ± 27.67 ^a	34.70 ± 18.53 ^a	28.55 ± 17.97 ^{ab}	51.25 ± 18.49 ^{ac}	3.70 ± 25.21 ^{ab}	194.00 ± 30.39 ^a
B	58.08 ± 9.13 ^a	39.88 ± 14.94 ^a	50.24 ± 17.57 ^b	24.00 ± 4.59 ^{ab}	BDL	256.52 ± 30.82 ^a
C	54.96 ± 10.79 ^a	51.96 ± 17.88 ^a	74.44 ± 16.90 ^b	68.28 ± 18.89 ^c	33.60 ± 38.47 ^{ab}	195.56 ± 22.10 ^a
Control	BDL	10.20 ± 3.75 ^a	BDL	5.45 ± 2.26 ^b	47.30 ± 5.17 ^a	113.80 ± 1.49 ^b

Values are expressed as mean ± standard error mean. Means with different superscripts down the column are significantly different ($P < 0.05$) at 95% confidence level using least significance difference (L.S.D), BDL-below detectible limit. Zn>Cu>Ni>Cd>Cr>Pb is the descending order of the heavy metals in the shoot samples from the maximum to the least accumulation at the sampling points

Table 3. Pearson Correlation Coefficient between Heavy metals in the roots and shoots of *E. crassipes* and between the heavy metals in the water samples during seasons at the sampling points

Heavy Metals	Roots and Shoots	Dry and Wet
Cadmium (mg/kg)	0.998*	0.784*
Chromium (mg/kg)	0.862*	0.833*
Copper (mg/kg)	0.836*	0.883*
Nickel (mg/kg)	0.605*	0.895*
Lead (mg/kg)	0.826*	0.457*
Zinc (mg/kg)	0.646*	0.855*

*Correlation is significant at $P < 0.05$




Fig. 4. Mean Bioaccumulation Coefficient of Heavy Metals in the shoots of *E. crassipes*. The descending order of bioaccumulation coefficient in the shoots from the maximum to the minimum at the sampling points is as follows; Ni>Pb>Cr>Cd>Zn>Cu




Fig. 5. Mean Translocation Factor of Heavy metals in *E. crassipes*. The descending order of Translocation factor ranged from the maximum to the minimum at the sampling points is as follows; Ni>Cu>Cd>Pb>Cr>Zn

If the translocation factor is less than one ($TF < 1$) (also known as excluder plant), whereas if the TF is greater than one ($TF > 1$) the plant is classified as a phytoaccumulator.

as a phytoextractor or accumulator [37]. This study was in agreement with the findings of Shingadgaon and Chavan [34] in evaluation of phytoremediation indices and abilities of aquatic macrophyte (*Salvinia molesta*) and other Species exposed to metal contaminated wastewater. Where the translocation factor in nickel was the highest and was greater than one. But was in contrast with the finding of Arifin et al. [33] who in their study, had the highest TF recorded in Zinc in the stems and leaves of *Rhazya stricta*.

The phytoremediation potential of a plant is characterized not just by its capacity to acquire and accumulate significant concentration of metal, as well, by the plant possessing the enabling capacity to translocate the metals to aerial components while also producing huge biomass [38]. This was observed in *E. crassipes* in this study. The Translocation factor in this investigation demonstrated that there was no significant difference ($P < 0.05$) in the different heavy metals at the sampling points. The translocation factor of heavy metals from shoot to root was greater than one. Zinc was not significantly translocated. Hence, its translocation factor was less than one in all the sampling points.

4. CONCLUSION

From this finding, the *E. crassipes* could be said to be phytostabilizer for Zinc and is a phytoextractor of Nickel, Lead, Cadmium, Chromium, and Copper. This is suggesting that *E. crassipes* is a good hyperaccumulator plant and serve as better candidate for phytoremediation.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Ekta C, Praveen S. Duckweed plant: A better future option for phytoremediation, International Journal of Emerging Science and Engineering (IJESE). 2014; 2:2319–6378.
- Alluri HK, Ronda SR, Settalluri VS, Bondili JS, Suryanarayana V, Venkateshwar P, et al. Biosorption: An eco-friendly alternative for heavy metal removal, African Journal of Biotechnology. 2007;6(25): 2924-2931.
- Meagher RB. Phytoremediation of toxic elemental and organic pollutants, Current Opinion Plant Biology. 2000;3:153-162.
- Nwoko CO. Trends in phytoremediation of toxic elemental and organic pollutants, African Journal Biotechnology. 2010;9:6010-6016.
- Khan AG, Kuek C, Chaudhry TM, Khoo CS, Hayes WJ, et al. Role of plants mycorrhizae and phytochelators in heavy metal contaminated land remediation, Chemosphere. 2000;41:197-207.
- Freeman JL, Persans MW, Nieman K, Albrecht C, Peer W, Pickering IJ, et al. Salt, increased glutathione biosynthesis plays a role in nickel tolerance in *Thlaspi* nickel hyperaccumulators, Plant Cell. 2004;16:2176-2191.
- Ali H, Khan E, Sajad MA. Phytoremediation of heavy metals concepts and applications, Chemosphere. 2013;91:869–881.
- Lee JH. An overview of phytoremediation as a potentially promising technology for environmental pollution control, Biotechnology Bioprocess Engineering. 2013;18(3):431–439.
- Ximenez-Embun P, Rodriguez-Sanz B, Madrid-Albarran Y, Camara C. et al. Uptake of heavy metals by lupin plants in artificially contaminated sand. Preliminary results, International Journal of Environmental and Analytical Chemistry. 2001;82:80-81.
- Federal Environmental Protection Agency (FEPA) Act. Guidelines and standards for Industrial effluent, gaseous emissions, and hazardous waste management in Nigeria. National Environmental Protection Regulations, Federal Republic of Nigeria. Supplement to Official Gazette Extraordinary - Part B. 1991;78(42):B15–31.
- US. Environmental Protection Agency (EPA). Arsenic in drinking water. Washington DC; 2006. Available: <http://www.epa.gov/safewater/arsenic/index.html> [Accessed on September 14; accessed 2009 March]
- US. EPA. Environmental criteria and assessment office. Cincinnati, OH: the United States Environmental Protection Agency, Integrated Risk Information System (IRIS); 1992.
- Raskin I, Smith RD, Salt DE. Phytoremediation of metals: using plants

- to remove pollutants from the environment, *Current Opinion Biotechnology*. 1997;8(2):221–226.
14. Jabeen R, Ahmad A, Iqbal M. Phytoremediation of heavy metals: Physiological and molecular mechanisms, *Botany Review*. 2009;75:339–364.
 15. Escande V, Olszewski TK, Grison C. Preparation of ecological catalysts derived from Zn hyperaccumulation plants and their catalytic activity in Diels–Alder reaction, *CR Chim*. 2014;17:731–737.
 16. Escande V, Olszewski T, Grison C. From biodiversity to catalytic diversity: how to control the reaction mechanism by the nature of metallophytes, *Environmental Science Pollution Resources*. 2015;22:5653–5666.
 17. Kang JW. Removing environmental organic pollutants with bioremediation and phytoremediation, *Biotechnology Letter*. 2014;36(6):1129–1139.
 18. Khalid M, Saeed UR, Hassani D, Hayat K, Pei ZHOU, Nan HUI. Advances in fungal-assisted phytoremediation of heavy metals: A review. *Pedosphere*. 2021;31:475–495. [CrossRef]
 19. Singh G, Pankaj U, Chand S, Verma RK. Arbuscular mycorrhizal fungi-assisted phytoextraction of toxic metals by *Zea mays* L. from tannery sludge. *Soil Sediment Contamination International Journal*. 2019;28:729–746. [CrossRef]
 20. Venegas-Rioseco J, Ginocchio R, Ortiz-Calderón C. Increase in phytoextraction potential by genome editing and transformation: A review. *Plants*. 2022;11:86.
Available:<https://doi.org/10.3390/plants11010086>
 21. Yan S, Wang Y, Wang Z, Guo JY. Remediation effects of experimental projects using water hyacinth for pollution control in the lake Caohai Dianchi Jiangsu, *Journal of Agricultural Sciences*. 2012;28(5):1025-1030.
 22. Aubert S, Schwitzguébel JP. Screening of plant species for the phytotreatment of wastewater containing sulphonated anthraquinones, *Water Resources*. 2004;38:3569-3575.
 23. Upadhyay AR, Tripathi BD. Principle and process of biofiltration of Cd, Cr, Co, Ni, and Pb from Tropical Opencast Coalmine Effluent, *Water, Air, and Soil Pollution*. 2007;180(1-4):213- 223.
 24. Thomas JD. Protocols for collecting water grab samples in rivers, streams and freshwater wetlands, Division of Environmental Assessment Biological Monitoring Program, DEPLWO. 2014;06374.
 25. Wahab OO, Liasu AO. Phytoremediation potential of some heavy metals by water hyacinth, *International Journal Biological Chemical Science*. 2010;4(2):347-353.
 26. Shaayau S, Garba NA. Phytoextraction and bioaccumulation potentials of *Ipomoea asarifolia* (Desr.) Roem. and Schult on Chromium-polluted Soils, *Nigerian Journal of Pure and Applied Science*. 2019;32:3469-3479:0794-0378.
 27. Blaylock MJ, Huang JW. Phytoextraction of Metals in phytoremediation of toxic metals: Using plants to clean up the environment. John Wiley and Sons Inc, New York. 1999;53-70.
 28. Environmental Protection Agency (EPA). Introduction to phytoremediation. 600/R-99/10. February. U.S. Cincinnati. Ohio; 2000.
 29. Mukesh Parmar, Lokendra Singh Thakur. Heavy metal Cu, Ni and Zn: Toxicity, Health Hazards and their Removal Techniques by low-Cost Adsorbents: A Short Overview, *International Journal of Plant, Animal and Environmental Sciences*. 2013;3(3):2231-4490.
 30. Ganesh T. Cloning and functional characterization of putative heavy metal stress-responsive (Echmr) gene from *Eichhornia crassipes* (Mart. Solm L.). [Doctoral Thesis Department of Biotechnology, Indian Institute of Technology Guwahati]. 2012;781-039.
 31. Higemengist Astatkie, Argaw Ambelu, Embialle Mengistie Beyene. Sources and level of heavy metal contamination in the water of Awetu watershed streams, southwestern Ethiopia. Published by Elsevier Ltd. 2021;2405-8440.
Available:<https://doi.org/10.1016/j.heliyon.2021.e06385>
 32. Oning Veschasit, Shettapong Meksumpun, Charumas Meksumpun. Heavy metals contamination in water and aquatic plants in the Tha Chin River, Thailand, *Kasetsart Journal of Natural Sciences*. 2012;(46):931–943.
 33. Arifin A, Parisa A, Hazandy AH, Mahmud TM, Junejo N, Fatemeh A, Mohsen S, Wasli ME, Majid NM, et al. Evaluation of cadmium bioaccumulation and

- translocation by *Hopea odorata* grown in contaminated soil, African Journal of Biotechnology. 2012;11(29):7472-7482: 1684–5315.
34. Al-Farraj AS, Al-Wabel MI, Al-Shahrani TS, El-Maghraby SE, Al-Sewailem MAS, et al. Accumulation coefficient and translocation factor of heavy metals through *Rhazya stricta* grown in the mining area of Mahad AD'Dahab, Saudi Arabia, Waste Management and the Environment. 2010;325:1743-3541.
35. Mahmood T, Syed TH, Salman AM. Phytoremediation technologies for Ni⁺⁺ by water hyacinth, African Journal of Biotechnology. 2010;9(50):8648-8660. DOI: 10.5897/AJB10.070, 1684–5315
36. Tabassum A. Studies on the role of microbes in bioremediation of metals, [Dissertation of master of philosophy in Biology (Microbiology), submitted to Department of Biological Sciences, Quaid-i-Azam University, Islamabad, Pakistan]. 2003:61-63.
37. Shingadgaon SS, Chavan BL. Evaluation of Bioaccumulation Factor (BAF), Bioconcentration Factor (BCF), Translocation Factor (TF) and Metal Enrichment Factor (MEF) Abilities of Aquatic Macrophyte Species Exposed to Metal Contaminated Wastewater, International Journal of Innovative Research in Science, Engineering and Technology. 2019;8(1):ISSN(Online):2319-8753 ISSN (Print): 2347-6710 DOI: 10.15680/IJIRSET
38. Labanowski J, Monna F, Bermond A. Kinetic extractions to assess mobilization of Zn, Pb, Cu, and Cd in a metal-contaminated soil: EDTA vs. Citrate. Environmental Pollution. 2008;152(3): 693–701.

© 2023 Yilwa et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/107968>