



# Synthesis and Characterization of Co(II), Ni(II), Cu(II) and Zn(II) Complexes with N'-( phenyl ( pyridin-2-yl)methylene)Nicotinohydrazide

Ndèye Fatou Ndiaye <sup>a</sup>, Fatou Diouf <sup>a</sup>, Amadou Gueye <sup>b</sup>,  
Farba Bouyagui Tamboura <sup>a\*</sup>, Moussa Dieng <sup>a</sup>,  
Sofia Zazouli <sup>c</sup>, Nathalie Gruber <sup>d</sup>  
and Mohamed Lamine Gaye <sup>b</sup>

<sup>a</sup> Department of Chemistry, UFR SATIC, University Alioune DIOP, Bambey 21400, Senegal.

<sup>b</sup> Department of Chemistry, University Cheikh Anta Diop, Dakar 10700, Senegal.

<sup>c</sup> Sultan Moulay Slimane University, Faculty of Sciences and Technologies, BP 523, 23000 Beni-Mellal, Morocco.

<sup>d</sup> Laboratoire de Tectonique Moléculaire du Solide (UMR 7140), Chimie de la Matière Complexe, Université de Strasbourg, Institut Le Bel, 4 rue Blaise Pascal, F-67008 Strasbourg, France.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/CSJI/2023/v32i6869

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/109487>

Original Research Article

Received: 14/09/2023

Accepted: 19/11/2023

Published: 22/11/2023

## ABSTRACT

The Schiff base (HL), derived from 2-benzoylpyridine and nicotinic hydrazide, and its complexes with some metal transition (Co, Ni, Cu, Zn) have been synthesized. These compounds have been characterized by means of elemental analysis, <sup>1</sup>H and <sup>13</sup>C NMR (for HL), FTIR spectroscopy, UV–

\*Corresponding author: E-mail: farba.tamboura@uadb.edu.sn;

Vis spectroscopy, molar conductance and room temperature magnetic measurements. The compounds are found isostructural and are formulated as {[Co(L)<sub>2</sub>].2H<sub>2</sub>O} (1), {[Ni(L)<sub>2</sub>].2H<sub>2</sub>O} (2), {[Cu(L)<sub>2</sub>].3H<sub>2</sub>O} (3), {[Zn(L)<sub>2</sub>].H<sub>2</sub>O} (4). The two ligand molecules act in their monodeprotonated forms through one azomethine nitrogen atom, one pyridine nitrogen atom and one iminolate oxygen atom yielding six coordinate metal center. The complex 4 crystallizes in the monoclinic space group P2<sub>1</sub> with cell parameters of a = 9.3429(8) Å, b = 10.3034(9) Å, c = 16.6349(14) Å, β = 101.109(4)°, V = 1571.3(2) Å<sup>3</sup>, Z = 2, R<sub>1</sub> = 0.027, wR<sub>2</sub> = 0.074. The zinc atom is six-coordinated, and the coordination polyhedron is best described as a distorted octahedral geometry. The aromatic rings of the ligand molecules are twisted each other with dihedral angle values in the range 10.168(2)°–74.430(1)°.

**Keywords:** Hydrazide; cobalt; nickel; copper; zinc; crystal; X-ray diffraction.

## 1. INTRODUCTION

Schiff bases resulting from the condensation reaction of nicotinic hydrazide and a ketoprecursor such as ketones[1,2,3] are widely used in the synthesis of organic ligands for the preparation of coordination complexes of lanthanide [4–6] and transition metals [7,8,9]. These complexes are particularly interesting in various fields such as magnetism [10], catalysis [11,12], in medicine [13,14] and luminescence [15,16]. The presence of phenolic nuclei suggests that these complexes can be antioxidants, as reported in the literature [17–18] but also as antibacterial agents [19–23]. “In recent studies, nicotinic hydrazide complexes have been prepared and shown to have antimicrobial-like activities” [24-26]. “It is in this context that we set out to synthesize complexes at room temperature using nicotinic hydrazide and 2-benzoylpyridine in the presence of transition metal salts. Several complexes have been isolated in powder form and characterized by various techniques”. [24-26]

## 2. METHODOLOGY

### 2.1 Materials and Procedures

Nicotinic acid hydrazide, 2-benzoylpyridine, M(NO<sub>3</sub>)<sub>2</sub>.nH<sub>2</sub>O (M = Co(II), Ni(II), Cu(II) or Zn(II)) were commercial products (from Alfa and Aldrich) and were used without further purification. Solvents were of reagent grade and were purified by the usual methods. Elemental analyzes were performed in a Carlo- Erba EA microanalyser. Infrared spectra were recorded as KBr discs on a Bruker IFS-66V spectrophotometer. LSI-MS were recorded using a Micromass Autospec spectrometer with 3-nitrobenzyl alcohol as the matrix. The <sup>1</sup>H and <sup>13</sup>C NMR spectra were recorded in DMSO-d<sub>6</sub> on a

BRUKER 500 MHz spectrometer at room temperature using TMS as an internal reference. The UV-Vis spectra were run on a Shimadzu UV-2501 PC Recording spectrophotometer (1000–200 nm). The molar conductance of 10<sup>-3</sup> M in DMF solutions of the metal complexes was measured at 25 °C using a WTW LF-330 conductivity meter with a WTW conductivity cell. Room temperature magnetic susceptibilities of the powdered samples were measured using a Johnson Matthey scientific magnetic susceptibility balance Hg[Co(SCN)<sub>4</sub>] used as calibrant. Melting points were recorded on a Büchi apparatus and are uncorrected [27].

### 2.2 Synthesis of the HL Ligand and its Transition Metal Complexes

#### 2.2.1 Synthesis of the ligand N'-( phenyl ( pyridin-2-yl)methylene)nicotinohydrazide (HL)

In a 250 mL round bottomed flask containing 20 mL of absolute ethanol, benzoylpyridine (2.67 g, 14.58 mmol) was added. After dissolution, nicotinic hydrazide (2.00 g, 14.58 mmol) previously dissolved in 20 mL of methanol was added. The mixture was heated under reflux for four hours. After filtration, the resulting clear brown solution was left under slow evaporation. Crystals suitable for X-ray analyze were collected after three days. IR (ν, cm<sup>-1</sup>) : 3062 (N–H), 3044 (C<sub>Ar</sub>–H), 1684 (C=O), 1583 (C=N), [1541-1422] (C<sub>Ar</sub>=C<sub>Ar</sub>), 1282 (CN); 1142 (N–N); 804; 749; 699; 650; 614 (C<sub>Ar</sub>–H). <sup>1</sup>H NMR (dmso-d<sub>6</sub>, δ, ppm): 8.73 (s, 1H, H–C=N); [7.39-8.20] (m, 13H, H–Ar), 9.01 (s, 1H, N–H). <sup>13</sup>C NMR (dmso-d<sub>6</sub>, δ, ppm): 161.46 (C=O), 152.30 (C=N), 151.68, 148.72, 138.11, 137.71, 136.78, 135.39, 135.22, 130.61, 129.23, 128.34, 124.91, 124.08, 123.76.

## 2.2.2 Synthesis of transition metal complexes from the ligand (HL)

The ligand HL (0.1g, 0.33 mmol) was dissolved in 15 mL of methanol before adding (0.17 mmol) of transition metal nitrate  $M(\text{NO}_3)_2 \cdot x\text{H}_2\text{O}$  with  $M = \text{Co}, \text{Ni}, \text{Cu}$  and  $\text{Zn}$ . The resulting solution was stirred for two hours at room temperature, then filtered and allowed to slowly evaporate. One week later we obtained powders which were recovered by filtration then recrystallized in a methanol/DMF mixture (50/50). After three weeks suitable yellow crystals for X-ray diffraction were collected in the case of the  $\text{Zn}(\text{II})$  complex. The physico-chemical data were collected using the powder or the crystals collected after the recrystallization process.

$\{\{\text{Co}(\text{L})_2\} \cdot 2\text{H}_2\text{O}\}$  (1). Brown. IR ( $\nu$ ,  $\text{cm}^{-1}$ ): 3672, 3236, 2968, 2969, 2929, 1610, 1568, 1524, 1383, 1291, 1264, 1191, 1072, 1067, 1052, 1028, 969, 815, 798, 758, 696. Yield: 78%. M.P. ( $^\circ\text{C}$ ) > 260.  $\Lambda$  ( $\Omega^{-1} \text{cm}^2 \text{mol}^{-1}$ ) 6 (fresh solution) and 9 (after two weeks).  $\mu_{\text{eff}}$ : 3.88  $\mu\text{B}$ . UV-visible (MeOH,  $\lambda_{\text{max}}$ , nm): 325, 361, 383, 418, 465, 499. Anal. calcd. for  $\text{C}_{36}\text{H}_{30}\text{CoN}_8\text{O}_4$ : % C, 61.98; % H, 4.33; % N, 16.06. Found % C, 61.93; % H, 4.30; % N, 16.09.

$\{\{\text{Ni}(\text{L})_2\} \cdot 2\text{H}_2\text{O}\}$  (2). Green. IR ( $\nu$ ,  $\text{cm}^{-1}$ ): 3383, 2900, 2835, 1598, 1548, 1408, 1368, 1301, 1249, 1196, 1151, 1086, 1019, 825, 748, 727, 696. Yield: 80%. M.P. ( $^\circ\text{C}$ ) > 260.  $\Lambda$  ( $\Omega^{-1} \text{cm}^2 \text{mol}^{-1}$ ): 10 (fresh solution) and 15 (after two weeks).  $\mu_{\text{eff}}$ : 2.74  $\mu\text{B}$ . UV-visible (MeOH,  $\lambda_{\text{max}}$ , nm): 361, 419, 444, 499, 869. Anal. calcd. for  $\text{C}_{36}\text{H}_{30}\text{NiN}_8\text{O}_4$ : % C, 62.00; % H, 4.34; % N, 16.07. Found % C, 61.90; % H, 4.30; % N, 16.09.

$\{\{\text{Cu}(\text{L})_2\} \cdot 3\text{H}_2\text{O}\}$  (3). Green. IR ( $\nu$ ,  $\text{cm}^{-1}$ ): 3396, 2888, 2825, 1618, 1595, 1569, 1531, 1478, 1385, 1197, 1161, 1099, 1056, 1030, 827, 761, 696, 656. Yield: 80%. M.P. ( $^\circ\text{C}$ ) > 260,  $\Lambda$  ( $\Omega^{-1} \text{cm}^2 \text{mol}^{-1}$ ): 8 (fresh solution) and 12 (after two weeks).  $\mu_{\text{eff}}$ : 1.82  $\mu\text{B}$ . UV-visible (MeOH,  $\lambda_{\text{max}}$ , nm): 386, 418, 439, 459, 498, 678. Anal. calcd. for  $\text{C}_{36}\text{H}_{32}\text{CuN}_8\text{O}_5$ : % C, 60.03; % H, 4.48; % N, 15.56. Found % C, 60.06; % H, 4.54; % N, 15.61.

$\{\{\text{Zn}(\text{L})_2\} \cdot \text{H}_2\text{O}\}$  (4). Yellow. IR ( $\nu$ ,  $\text{cm}^{-1}$ ): 3649, 2364, 1559, 1507, 1457, 1360, 1302, 1257, 1193, 1150, 1082, 1053, 1033, 1014, 918, 834, 792, 742, 697. Yield: 72%. M.P. ( $^\circ\text{C}$ ) > 260,  $\Lambda$  ( $\Omega^{-1} \text{cm}^2 \text{mol}^{-1}$ ): 3 (fresh solution) and 5 (after two

weeks).  $\mu_{\text{eff}}$ : 0  $\mu\text{B}$ . UV-visible 361, 419, 441, 499. Anal. calcd. for  $\text{C}_{36}\text{H}_{28}\text{N}_8\text{O}_3\text{Zn}$ : % C, 63.03; % H, 4.11; % N, 16.33. Found % C, 62.97; % H, 4.14; % N, 16.39.

## 2.3 Crystal Structure Determination

"Crystals suitable for X-diffraction, of the reported compound, were grown by slow evaporation of MeOH /DMF mixture solution of the complex. Details of the X-rays crystal structure solution and refinement are given in Table 1. Diffraction data were collected using the Bruker SHELXTL diffractometer with graphite monochromatized Mo  $K\alpha$  radiation ( $\lambda = 0.71073 \text{ \AA}$ ). All data were corrected for Lorentz and polarization effects. No absorption correction was applied. Using the structures were solved by intrinsic phasing methods with SHELXT and SHELXL was used for full matrix least squares refinement". [28,29] "The hydrogen atoms of water molecules and NH groups were located in the Fourier difference maps and refined. Other H atoms were geometrically optimized and refined as riding model by AFIX instructions. Molecular graphics were generated using ORTEP-3" [30].

## 3. RESULTS AND DISCUSSION


### 3.1 General Study

Reaction of hydrated  $M(\text{II})$  nitrate ( $M = \text{Co}^{2+}, \text{Ni}^{2+}, \text{Cu}^{2+}$  and  $\text{Zn}^{2+}$ ) and HL in 1:2 ratio in methanol produces mononuclear complexes formulated as  $\{\{\text{Co}(\text{L})_2\} \cdot 2\text{H}_2\text{O}\}$  (1),  $\{\{\text{Ni}(\text{L})_2\} \cdot 2\text{H}_2\text{O}\}$  (2),  $\{\{\text{Cu}(\text{L})_2\} \cdot 3\text{H}_2\text{O}\}$  (3) and  $\{\{\text{Zn}(\text{L})_2\} \cdot \text{H}_2\text{O}\}$  (4) (Scheme 1)

The infrared spectrum of the ligand presents two bands pointed at  $3062 \text{ cm}^{-1}$  and  $3004 \text{ cm}^{-1}$  attributed, respectively to  $\nu_{\text{N-H}}$  and to  $\nu_{\text{C}_{\text{Ar}}-\text{H}}$  vibrations. The  $\nu_{\text{C=O}}$  vibration of the carbonyl group is located at  $1684 \text{ cm}^{-1}$  and the  $\nu_{\text{C=N}}$  of the imine group is pointed at  $1635 \text{ cm}^{-1}$  [31]. The  $\nu_{\text{C}_{\text{Ar}}=\text{C}_{\text{Ar}}}$  bands appear in the range  $1541\text{--}1422 \text{ cm}^{-1}$ . The characteristic band due to pyridine ring is pointed at  $1583 \text{ cm}^{-1}$ . The band pointed at  $1282 \text{ cm}^{-1}$  and  $1142 \text{ cm}^{-1}$  are, respectively, attributed to  $\nu_{\text{C-N}}$  and  $\nu_{\text{N-N}}$  vibrations. The deformation bands of the  $\text{C}_{\text{Ar}}-\text{H}$  bonds are located in the range  $804\text{--}614 \text{ cm}^{-1}$ .  $^1\text{H}$  and carbon  $^{13}\text{C}$  NMR spectra are carried out using  $\text{dmsO-d}_6$  solutions.  $^1\text{H}$  NMR spectrum indicates a set of signals in the form of multiplets between 7.39 and 8.20 ppm representing the eight

protons of the aromatic rings. In addition, the two appearing as singlets at 8.73 and 9.01 ppm are, respectively, due to the azomethine H-C=N and to the H-N protons. <sup>13</sup>C NMR spectrum indicates two characteristic signal at 160.10 and 148.91 ppm due to the carbon atoms of the C=O and H-C=N moieties. Signals due to aromatic carbon

atoms appears in the range 151.68–123.76 ppm. “The mass spectrum confirms the formation of the HL ligand with the presence of the basic peak at  $m/z = 303.12$  corresponding to the molar mass of the molecular ion (M+1). In our previous work the crystallographic structure of HL and its antioxidant activity were reported” [17].



**Scheme 1. Synthesis procedure of the ligand HL and its Co<sup>2+</sup>, Ni<sup>2+</sup>, Cu<sup>2+</sup> and Zn<sup>2+</sup>**

**Table-1. Crystal data and details of the structure determination of the complex [Zn(L)<sub>2</sub>]-H<sub>2</sub>O**

Chemical Formula	C <sub>36</sub> H <sub>26</sub> N <sub>8</sub> O <sub>2</sub> Zn·H <sub>2</sub> O
Shape and Color	Prism, colorless
Molar mass (g/ mol )	686.05
Crystal system,	Monoclinic,
Space group	P2 <sub>1</sub>
Crystal size (mm)	0.13 × 0.12 × 0.12
Temperature (K)	173
Mo Kα (Å)	0.71073
a (Å)	9.3429 (8)
b (Å)	10.3034 (9)
c (Å)	16.6349 (14)
β (°)	101.109 (4)
V (Å <sup>3</sup> )	1571.3 (2)
Z	2
D <sub>calc</sub> g/cm <sup>3</sup>	1.450
F(000)	354
μ (mm <sup>-1</sup> )	0.83
θ (°)	2.3-27
h, k, l ranges	-9 ≤ h ≤ 12, -14 ≤ k ≤ 14, -23 ≤ l ≤ 23
Measured reflections	65719
Independent reflections	8260
Reflections [I > 2σ(I)]	7596
R <sub>int</sub>	0.031
R[I > 2σ(I)]	0.027
wR <sub>2</sub>	0.074
Goodness-of-Fit	0.99
Δρ <sub>max</sub> , Δρ <sub>min</sub> (e Å <sup>-3</sup> )	0.50, -0.28

The infrared spectra of the four complexes 1–4 indicate shift of the  $\nu_{C=N}$  indicating the involvement of the azomethine nitrogen atom in the coordination. The absence of the  $\nu_{C=O}$  and the  $\nu_{N-H}$  indicate that iminisation undergoes during the reaction. The absences of ionic and coordinating nitrate bands on the spectra of the complexes are indicative of coordination through nitrogen atoms and oxygen atoms from the ligand molecules. On the infrared spectrum of the crystal complex of  $[ZnL_2] \cdot H_2O$ , the  $\nu(C=N)$  bands of imine and pyridine are located at  $1558\text{ cm}^{-1}$  and  $1506\text{ cm}^{-1}$ , respectively. However, we note the iminization of the ligand which is confirmed by the disappearance of the  $\nu_{C=O}$  band of the carbonyl and the appearance of new bands at  $1620\text{ cm}^{-1}$  and  $1250\text{ cm}^{-1}$  attributable to  $\nu(C=N)$  in the hydrazone and  $\nu(CO)$  of iminole [32]. In each of the spectra of the complexes the presence of uncoordinated water molecule is attested by the presence of one strong and broad band in the range  $3650\text{--}3300\text{ cm}^{-1}$ . Conductometric measurements of four complexes were carried out in a millimolar solution of Dimethylformamide (DMF). The values obtained in fresh solution and after two weeks of storage are fall in the range equal to  $3\text{--}15\ \Omega^{-1}\cdot\text{cm}^2\cdot\text{mol}^{-1}$ . These low values are typical of neutral electrolytes [33]. The low of the conductivity values over time show the good stability of these complexes in the DMF solution.

The electronic spectra of the complexes (Table 2) which are recorded in DMF solution show absorptions in the range  $257\text{--}275\text{ nm}$  and  $315\text{--}361\text{ nm}$  regions. "These bands are attributed, respectively, to the  $\pi \rightarrow \pi^*$  transitions of the aromatic nuclei and  $n \rightarrow \pi^*$  transitions of the azomethine moiety of the organic ligand molecules. The bands in the range  $418\text{--}441\text{ nm}$  are attributed to ligand-metal charge transfers" [34]. "For the cobalt(II) complex (1), the bands pointed at  $465$  and  $499\text{ nm}$  are attributable, respectively, to the transitions  ${}^4T_{1g}(F) \rightarrow {}^4T_{2g}(P)$  and  ${}^2E_g \rightarrow {}^2T_{1g}(P)$ . In the UV region, a band appears at  $383\text{ nm}$  due to the transition  ${}^4T_{1g}(F) \rightarrow {}^4A_{2g}(F)$ . These bands are typical of an octahedral environment around Co(II) cation [35]. For the complex (1), the value of the magnetic moment is  $5.88\ \mu_B$ . On the spectrum of the Ni(II) complex (2), the bands are located at  $444$ ,  $499$  and  $869\text{ nm}$  (Table 4). They are attributed to the transitions  ${}^1A_{1g} \rightarrow {}^1B_{1g}$ ,  ${}^1A_{1g} \rightarrow {}^1E_g$  and  ${}^3A_{2g}(F) \rightarrow {}^3T_{2g}$  in agreement with an octahedral geometry around nickel(II)" [35]. "This observation is confirmed by the magnetic moment the value of

$2.74\ \mu_B$  corresponding to an octahedral Ni(II) cation. The spectrum of the Cu(II) complex (3) presents an absorption band at  $678\text{ nm}$  (Table 4) which is assigned to the transition band  ${}^2E_g \rightarrow {}^2T_{2g}$  indicating an octahedral geometry around Cu(II) cation" [36]. "This observation is confirmed by magnetic value of  $1.82\ \mu_B$  which indicates the presence of a  $Cu^{2+}$  ion in octahedral environment" [37]. "The spectra of the Zn(II) complex (4) show strong absorption bands at  $499\text{ nm}$  due to charge transfers Ligand  $\rightarrow$  Metal" [38].

### 3.2 Structure Description of the Complex $C_{36}H_{26}N_8O_2Zn \cdot H_2O$

The mononuclear complex crystallizes in the monoclinic system  $P2_1$ . Labelled plot of the mononuclear structure of Zn(II) complex is shown in Fig. 1. Selected interatomic distances are listed in Table 3. The structure of the complex is consistence with the  $[ZnL_2] \cdot H_2O$  formulation. The asymmetric unit contains one  $Zn^{2+}$ , two monoanionic organic ligand, and one water molecule. Each Schiff base molecules is its iminol form acts in tridentate fashion through one azomethine nitrogen atom, one pyridine nitrogen atom and one anionic oxygen atom resulting in two membered chelating rings  $ZnNCCN$  and  $ZnNNCO$  with bite angles of  $N1\text{--}Zn1\text{--}N2 = 74.23(7)^\circ$ ,  $O1\text{--}Zn1\text{--}N2 = 75.72(7)^\circ$ ,  $N5\text{--}Zn1\text{--}N6 = 74.82(7)^\circ$ ,  $O2\text{--}Zn1\text{--}N6 = 75.41(7)^\circ$ . The Zn1 center is octacoordinated and is situated in a severely distorted octahedral environment. The basal plane is occupied by N1, N2, O1 and N6 atoms, the apical positions being occupied by N3 and N8 atoms. The angles in the basal plan  $O1\text{--}Zn1\text{--}N2 = 75.72(7)^\circ$ ,  $N6\text{--}Zn1\text{--}O1 = 112.98(7)^\circ$ ,  $N1\text{--}Zn1\text{--}N6 = 97.36(7)^\circ$ ,  $N1\text{--}Zn1\text{--}N2 = 74.23(7)^\circ$ ,  $O1\text{--}Zn1\text{--}N1 = 149.66(7)^\circ$ ,  $N6\text{--}Zn\text{--}N2 = 168.73(7)^\circ$ . The sum of the cisoids angles subtended by the atoms in the basal plane is  $360.3^\circ$ . The value of the angle between the apical atoms is  $O2\text{--}Zn1\text{--}N5 = 150.21(7)^\circ$ . All the angles deviate severely from the ideal angle values of  $90^\circ$  and  $180^\circ$  for octahedral geometry. These facts are indicative of a severely distorted octahedral polyhedron around the zinc(II) cation center. The aromatic rings  $C1/C2/C3/C4/C5/N31$  and  $C7/C8/C9/C10/C11/C12$  of one of the ligand molecule are twisted with dihedral angle of  $52.55(1)^\circ$ . They are respectively twisted toward the third aromatic ring  $C14/C15/N4/C17/C18/C19$  of the ligand molecule with dihedral angle of  $21.72(2)^\circ$  and  $33.35(2)^\circ$ . The five membered

rings Zn1O1C13N3N2 and Zn1N2C6C5N1 formed by the ligand, upon coordination to the Zn(II), are not coplanar. Their mean planes form a dihedral angle of 4.04(1)°. For the second ligand molecule the mean planes of the five membered rings formed upon coordination [Zn1O2C32N7N6 and Zn1N6C25C24N5] are quite coplanar with a dihedral angle of 1.86(1)°. The aromatic rings C20/C21/C22/C23/C24/N5 and C26/C27/C28/C29/C30/C31 of the second ligand molecule are twisted with dihedral angle of 64.792(1)°. They are respectively twisted toward the third aromatic ring C34/C33/C38/C37/C36/N8 of the ligand molecule with dihedral angle of 10.168(2)° and 74.430(1)°.

The Zn—O bonds are 2.0710 (18) and 2.1061(17) Å and are similar to those found for octahedral Zn(II) complex reported by Kane et al.[39]. The Zn—N<sub>imine</sub> bond distances (Zn—N2 and Zn—N6) are observed to be 2.1004 (17) and 2.0812 (18) Å, respectively. They are shorter than those of the Zn—N<sub>pyridine</sub> bonds (Zn—N1 = 2.189 (2) Å and Zn—N5 = 2.227 (2) Å). These Zn—N bonds are similar to those observed for octahedral mononuclear complexes obtained with a similar ligand (E)-N'-(1-(pyridin-2-yl)ethylidene)nicotinohydrazide [40]. Intramolecular hydrogen bond involving the oxygen atom of the uncoordinated water molecule and nitrogen pyridine atom [O3—H3A...N8] is observed.

**Table 2. UV-visible and Magnetic moments of the complexes**

Compound	UV-visible $\lambda$ (nm)	$\mu_{\text{eff}}$ ( $\mu_B$ )
HL	257, 275, 315, 361	
1	257, 275, 315, 325, 361, 383, 418, 465, 499	3.88
2	257, 275, 315, 361, 419, 444, 499, 869	2.74
3	257, 275, 315, 386, 418, 439, 459, 498, 678	1.82
4	257, 275, 315, 361, 419, 441, 499	Diam

**Table 3. Selected interatomic bond distances and bond angles around Zn**

<b>Zn1—O1</b>	<b>2.0710 (18)</b>
Zn1—N6	2.0812 (18)
Zn1—N2	2.1004 (17)
Zn1—O2	2.1061 (17)
Zn1—N1	2.189 (2)
Zn1—N5	2.227 (2)
O1—Zn1—N5	92.42 (8)
N6—Zn1—N5	74.82 (7)
N2—Zn1—N5	98.26 (7)
O2—Zn1—N5	150.21 (6)
N1—Zn1—N5	95.93 (8)
O1—Zn1—N6	112.98 (7)
O1—Zn1—N2	75.72 (7)
N6—Zn1—N2	168.73 (7)
O1—Zn1—O2	98.24 (7)
N6—Zn1—O2	75.41 (7)
N2—Zn1—O2	111.29 (7)

**Table 4. Hydrogen-bond geometry (Å, °)**

D—H...A	D—H	H...A	D...A	D—H...A
O3—H3A...N8	0.87	2.10	2.917(7)	157
C3—H3...N4	0.95	2.51	3.280(4)	138
C12—H12...N3	0.95	2.51	2.912(3)	105
C15—H15...N3	0.95	2.46	2.796(3)	101
C19—H19...N4	0.95	2.56	3.296(4)	134
C28—H28...O1	0.95	2.45	3.161(3)	132
C34—H34...N7	0.95	2.53	3.476(4)	172
C35—H35...O3	0.95	2.22	2.930(8)	131




Fig. 1. ORTEP plot (30% probability ellipsoids) showing the structure 4. H atoms are omitted for clarity




Fig. 2. The packing of the compound in the crystal structure

In the crystal, unclassical intramolecular C—H...N [C3—H3...N3, C12—H12...N3, C15—H15...N3, C19—H19...N4, C34—H34...N7] and C—H...O [C28—H28...O1, C35—H35...O3] interactions are observed in the crystal (Table 3, Fig. 2).

#### 4. CONCLUSION

This present paper describes the results of the preparation of coordination complexes of transition metal (II) nitrate salt (Co, Ni, Cu and Zn) with the ligand N'-(phenyl(pyridin-2-yl)methylene)nicotinohydrazide (HL). The ligand is characterized by elemental analyze, mass spectroscopy, and different spectrophotometric methods such as <sup>1</sup>H and <sup>13</sup>C NMR, FT-IR and UV-visible. The structures of the complexes 1-4 are elucidated by element analyze, molar conductivity, room temperature magnetic measurements, FT-IR, and UV-Visible spectroscopies. X-ray crystallographic diffraction technic is used to solve the structure of complex 4. In all the structures of the compounds, the two ligand molecules act in their iminol forms yielding six-coordinated metal ions complexes. The coordination geometry can be described as a severely distorted octahedral polyhedron.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

- Juyal VK, Pathak A, Panwar M, Thakuri SC, Prakash O, Agrwal A, Nand V. Schiff base metal complexes as a versatile catalyst: A review. *Journal of Organometallic Chemistry*. 2023;999:122825. Available:https://doi.org/10.1016/j.jorganchem.2023.122825
- Despaigne AAR, Silva JGD, Carmo ACMD, Piro OE, Castellano EE, Beraldo H. Copper(II) and zinc(II) complexes with 2-benzoylpyridine-methyl hydrazone. *Journal of Molecular Structure*. 2009;920(1):97–102. Available:https://doi.org/10.1016/j.molstruc.2008.10.025
- Despaigne AAR, Silva JGD, Carmo ACM, Do Piro OE, Castellano EE, Beraldo H. Structural studies on zinc(II) complexes with 2-benzoylpyridine-derived hydrazones. *Inorganica Chimica Acta*. 2009;362(7):2117–2122. Available:https://doi.org/10.1016/j.ica.2008.09.040
- Woods M, Kovacs Z, Sherry AD. Targeted Complexes of Lanthanide(III) Ions as Therapeutic and Diagnostic Pharmaceuticals. *Journal of Supramolecular Chemistry*. 2002;2(1):1–15. Available:https://doi.org/10.1016/S1472-7862(02)00072-2
- Tamboura FB, Diop M, Gaye M, Sall AS, Barry AH, Jouini T. X-ray structure and spectroscopic properties of some lanthanides(III) complexes derived from 2,6-diacetylpyridine-bis(benzoylhydrazone). *Inorganic Chemistry Communications*. 2003;6(8):1004–1010. Available:https://doi.org/10.1016/S1387-7003(03)00167-9
- Curtius Th, Heidenreich K. Ueber Hydraziund Azocarbonester. *Berichte der deutschen chemischen Gesellschaft*. 1894;27(1):773–774. Available:https://doi.org/10.1002/cber.189402701150
- Fallah-Mehrjardi M, Kargar H, Munawar KS. Tridentate ONO hydrazone Schiff base complexes in organic transformations: Catalytic and mechanistic studies. *Inorganica Chimica Acta*. 2024;560:121835. Available:https://doi.org/10.1016/j.ica.2023.121835
- Abou-Melha KS. Transition metal complexes of isonicotinic acid (2-hydroxybenzylidene)hydrazide. *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy*. 2008;70(1):162–170. Available:https://doi.org/10.1016/j.saa.2007.07.023
- Aly SA, Fathalla SK. Preparation, characterization of some transition metal complexes of hydrazone derivatives and their antibacterial and antioxidant activities. *Arabian Journal of Chemistry*. 2020;13(2):3735–3750. Available:https://doi.org/10.1016/j.arabjc.2019.12.003
- Kahn O. *Molecular Magnetism*. Courier Dover Publications; 2021.
- Catalysis of transesterification reactions by lanthanides - Unprecedented acceleration of methanolysis of aryl and alkyl esters



- promoted by La(OTf)<sub>3</sub> at neutral pH and ambient temperatures | Request PDF. (n.d.). ResearchGate. Available:<https://doi.org/10.1139/cjc-79-11-1704>
12. Organolanthoid Chemistry, Synthesis, Structure, Catalysis. (n.d.). Retrieved from Available:<https://link.springer.com/book/10.1007/BFb0015593>
  13. Londoño-Mosquera JD, Aragón-Muriel A, Cerón DP. Synthesis, antibacterial activity and DNA interactions of lanthanide(III) complexes of N(4)-substituted thiosemicarbazones. *Universitas Scientiarum*. 2018;23(2):141–169. Available:<https://doi.org/10.11144/Javeriana.SC23-2.saqa>
  14. Hijazi AK, Taha ZA, Ajlouni AM, Al-Momani WM, Idris IM, Hamra EA. (n.d.). Synthesis and Biological Activities of Lanthanide (III) Nitrate Complexes with N-(2-hydroxynaphthalen-1-yl) methylene) Nicotinohydrazone Schiff Base. *Medicinal Chemistry*. 13(1):77–84.
  15. (PDF) Synthesis, characterization, luminescence and biological activities of lanthanide complexes with a hydrazone ligand. (n.d.). ResearchGate. Available:<https://doi.org/10.3233/MGC-180251>
  16. Panayiotidou L, Drouza C, Arabatzis N, Lianos P, Stathatos E, Viskadourakis Z, Keramidis A D. Structure, reactivity, luminescence and magnetism of dinuclear Ln<sup>3+</sup> complexes produced by the Ln<sup>3+</sup>-assisted hydrolysis of 3,6-bis(2-pyridyl)tetrazine. *Polyhedron*. 2013;64:308–320. Available:<https://doi.org/10.1016/j.poly.2013.05.037>
  17. Seck TM, Faye FD, Gaye AA, Thiam IE, Diouf O, Gaye M, Retailleau P. Synthesis of mono and bis-substituted asymmetrical compounds, (1-(pyridin-2-yl)ethylidene)carbonohydrazone and 1-(2'-hydroxybenzylidene)-5-(1'-pyridylethylidene)carbonohydrazone: Structural characterization and antioxidant activity study. *European Journal of Chemistry*. 2020;11(4):285–290. Available:<https://doi.org/10.5155/eurjchem.11.4.285-290.2023>
  18. Haba PM, Tamboura FB, Diouf O, Gaye M, Sall AS, Baldé CA, Slebodnick C. Preparation, spectroscopic studies and x-ray structure of homobinuclear lanthanide(III) complexes derived from 2,6-diformyl-4-chlorophenol-bis-(2'-hydroxybenzoylhydrazone). *Bulletin of the Chemical Society of Ethiopia*. 2006;20(1):45–54. Available:<https://doi.org/10.4314/bcse.v20i1.21142>
  19. IK, MT, Vk R. New lanthanide complexes with antioxidant activity. *Medicinal chemistry (Sharifah (United Arab Emirates))*. 2008;4(4). Available:<https://doi.org/10.2174/157340608784872181>
  20. Tamboura FB, Gaye M, Sall AS, Barry AH, Jouini T. Synthesis, properties and X-ray structure for the mononuclear complex of [(1-methyl imidazol-2-yl)methylene-2-aminoethylpyridine]-dichloro copper (II) monohydrate. *Inorganic Chemistry Communications*. 2002;5(4):235–238. Available:[https://doi.org/10.1016/S1387-7003\(02\)00348-9](https://doi.org/10.1016/S1387-7003(02)00348-9)
  21. Dongli C, Handong J, Hongyun Z, Deji C, Jina Y, Jian LB. Studies on acetylferrocenyl nicotinoyl and isonicotinoyl hydrazones and their coordination compounds with transition metals—II. *Polyhedron*. 1994;13(1):57–62. Available:[https://doi.org/10.1016/S0277-5387\(00\)86638-3](https://doi.org/10.1016/S0277-5387(00)86638-3)
  22. Madanhire T, Davids H, Pereira MC, Hosten EC, Abrahams A. Synthesis, characterisation and anticancer activity screening of lanthanide(III) acetate complexes with benzohydrazone and nicotinohydrazone ligands. *Polyhedron*. 2020;184:114560. Available:<https://doi.org/10.1016/j.poly.2020.114560>
  23. Mohamad ADM, Abualreish MJA, Abu-Dief AM. Antimicrobial and anticancer activities of cobalt (III)-hydrazone complexes: Solubilities and chemical potentials of transfer in different organic co-solvent-water mixtures. *Journal of Molecular Liquids*. 2019;290:111162. Available:<https://doi.org/10.1016/j.molliq.2019.111162>
  24. Patil P, Zangade S. Synthesis, characterization, antimicrobial screening and cytotoxic properties of Cu(II) and Zn(II) complexes with bidentate hydroxylated 1,3-diaryl-2-propene-1-ones ligand. *Journal of the Serbian Chemical Society*. 2021;86(2):153–164. Available:<https://doi.org/10.2298/JSC200901068Z>

25. Fasina TM, Dueke-Eze CU, Idika FN. Synthesis, spectroscopic and antimicrobial properties of Co(II), Ni (II) and Cu(II) complexes of (E)-N'-(2-hydroxy -5-nitrobenzylidene)isonicotinohydrazide). *Journal of Applied Sciences and Environmental Management*. 2017;21(1): 120–125. Available: <https://doi.org/10.4314/jasem.v21i1.13>
26. Wehrmann M, Billard P, Martin-Meriadec A, Zegeye A, Klebensberger J. Functional Role of Lanthanides in Enzymatic Activity and Transcriptional Regulation of Pyrroloquinoline Quinone-Dependent Alcohol Dehydrogenases in *Pseudomonas putida* KT2440. *mBio*; 2017. Available: <https://doi.org/10.1128/mbio.00570-17>
27. SOW MM, DIENG M, Bouyagui F. Synthesis and characterization of novel M (II)(M= Co (II), Ni (II), Cu (II), Zn (II) or Cd (II)) complexes with N, O-donor ligand (E)-1-(1-(1H-pyrrol-2-yl) ethylidene) carbonohydrazide (H5L).
28. Sheldrick GM. it SHELXT – Integrated space-group and crystal-structure determination. *Acta Crystallographica Section A*. 2015;71(1):3–8. Available: <https://doi.org/10.1107/S2053273314026370>
29. Sheldrick GM. Crystal structure refinement with it SHELXL. *Acta Crystallographica Section C*. 2015;71(1):3–8. Available: <https://doi.org/10.1107/S2053229614024218>
30. Farrugia LJ. it WinGX and it ORTEP for Windows: an update. *Journal of Applied Crystallography*. 2012;45(4):849–854. Available: <https://doi.org/10.1107/S0021889812029111>
31. Moksharagni B, Reddy KH. Spectral Characterization And Dna Binding Properties Of Lanthanide (Iii) Complexes With 2acetylpyridine Nicotinoylhydrazone (APNH). *World Journal of Pharmaceutical Research*. 2017;6(2):866–876. Available: <https://doi.org/10.20959/wjpr20172-7747>
32. Znovjyak K, Fritsky IO, Sliva TY, Amirkhanov VM, Seredyuk M. Crystal structure of the mixed methanol and ethanol solvate of bis\3,4,5-trimeth\oxy-it N^\prime-[1-(pyridin-2-yl)ethyl\idene]benzohydrazidatozinc(II). *Acta Crystallographica Section E*. 2020;76(3): 303–308. Available: <https://doi.org/10.1107/S2056989020000857>
33. Geary WJ. The use of conductivity measurements in organic solvents for the characterisation of coordination compounds. *Coordination Chemistry Reviews*. 1971;7(1):81–122. Available: [https://doi.org/10.1016/S0010-8545\(00\)80009-0](https://doi.org/10.1016/S0010-8545(00)80009-0)
34. Gueye A, Tamboura FB, Sy A, Gaye M, Gruber N, Jouaiti A. Six New Transition Metal Mononuclear Complexes Of N'-(5-bromo-2-hydroxybenzylidene)nicotinohydrazide Schiff Base. Synthesis, Spectroscopic Characterization And X-ray structure Determination of the Zinc(II) Complex. *IOSR Journal of Applied Chemistry*. 2019;12:24–30. Available: <https://doi.org/10.9790/5736-1204012430>
35. Alaghaz ANMA, Ammar YA, Bayoumi HA, Aldhlmani SA. Synthesis, spectral characterization, thermal analysis, molecular modeling and antimicrobial activity of new potentially N2O2 azo-dye Schiff base complexes. *Journal of Molecular Structure*. 2014;1074:359–375. Available: <https://doi.org/10.1016/j.molstruc.2014.05.078>
36. Elaaraj I, Raouan S, ER, Nakkabi A, Es-sounni B, Koraichi I, El moualij N, Fahim M. Synthesis, characterization and antioxidant, antibacterial activity Zn<sup>2+</sup>, Cu<sup>2+</sup>, Ni<sup>2+</sup> and Co<sup>2+</sup>, complexes of ligand [2-(thiophen-2-yl)-1-(thiophen-2-ylmethyl)-1H-benzo[d]imidazole]. *Journal of the Indian Chemical Society*. 2022;99(5):100404. Available: <https://doi.org/10.1016/j.jics.2022.100404>
37. Gutierrez K, Corchado J, Lin S, Chen Z, Piñero Cruz DM. A non-innocent salen naphthalene ligand and its Co<sup>2+</sup>, Ni<sup>2+</sup> and Cu<sup>2+</sup> metal complexes: Structural, electrochemical, and spectroscopic characterization and computational studies. *Inorganica Chimica Acta*. 2018; 474:118–127. Available: <https://doi.org/10.1016/j.ica.2018.01.002>
38. Alyass JM, Mohammed AF. Preparation and characterization of some new complex salts of Cr(III), Fe(III), Mn(II), Co(II), Ni(II), Cu(II) and Zn(II) ions with tetra propyl ammonium iodide. *Egyptian Journal of Chemistry*. 2022;65(1):239–244.

- Available:<https://doi.org/10.21608/ejchem.2021.80143.3953>
39. Kane CH, Tinguiano D, Tamboura FB, Thiam IE, Barry AH, Gaye M, Retailleau P. Synthesis and characterization of novel M(II) (M = Mn(II), Ni(II), Cu(II) or Zn(II)) complexes with tridentate N<sub>2</sub>O-donor ligand (E)-2-amino-N'-[1-(pyridin-2-yl)-ethylidene]benzohydrazide. Bulletin of the Chemical Society of Ethiopia. 2016;30(1): 101–110.
- Available:<https://doi.org/10.4314/bcse.v30i1.9>
40. Shen S, Chen H, Zhu T, Ma X, Xu J, Zhu W, Peng C. Synthesis, characterization and anticancer activities of transition metal complexes with a nicotinothiohydrazone ligand. Oncology Letters. 2017;13(5): 3169–3176.
- Available:<https://doi.org/10.3892/ol.2017.5857>

## SUPPLEMENTARY MATERIALS

CCDC-2304399 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge via <https://www.ccdc.cam.ac.uk/structures/>, or by e-mailing [data\\_request@ccdc.cam.ac.uk](mailto:data_request@ccdc.cam.ac.uk), or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: +44(0)1223-336033.

© 2023 Ndiaye et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/109487>