



Evaluation of Proximate Composition and Seasonal Variations of the Muscle Tissue of Mouth-Brooding Tilapia *Oreochromis mossambica* from Meghadrigedda Reservoir, Visakhapatnam, Andhra Pradesh, India

**Krishna P. V. ^{a*}, P. Varalakshmi ^a, B. V. L. Aradhya Sarma ^a,
P. Dedeepya ^a and K. Durga Rao ^a**

^a *Department of Zoology and Aquaculture, Acharya Nagarjuna University, Nagarjuna Nagar – 522510, Andhra Pradesh, India.*

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJFAR/2023/v25i6719

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/110277>

Original Research Article

Received: 13/10/2023

Accepted: 15/12/2023

Published: 27/12/2023

ABSTRACT

This study aimed to investigate the proximate composition and seasonal variations in the muscle of mouth-brooding tilapia *O. mossambica* from Meghadrigedda reservoir, Visakhapatnam. The biochemical analysis such as total protein, total fat, total carbohydrates, ash and moisture content in the muscle of fish was taken for seasonal variations in relation to reproductive characters during 2021-22. The total protein percent of the fish *O. mossambica* ranged from 24.5 to 36.8 in females and males was 24.5 and 37.5. The total fat percentage ranged from 5.4 to 8.6 in females and

*Corresponding author: Email: drpvkrishna@gmail.com;

males was 6.4 to 8.7. Total Carbohydrates percentage was 4.1 to 5.5 in females and males was 4.8 and 5.9. Moisture percentage was 80.2 to 84.5 in females and males was 75.5 to 85.5. Ash percentage was 4.1 to 4.9 in females and males was 4.1 to 4.8. The results of the present clearly indicate that there was differences in the proximate composition of the male and female species.

Keywords: *O. mossambica*; proximate composition; maturation and spawning.

1. INTRODUCTION

“Fish and fishery products are dynamic source of food items for mankind all over the globe due to its high-quality proximate composition. It contains the most important protein and low fat components serve as a vital energy for human beings and their domestic animals” [1]. FAO [2] estimated that nearly 11% of the total population of the world is suffering from poverty and do not have access to enough nutritious food for proper growth and human health. WHO [3] has reported tripled number of obese people since 1975 which accounts for 13% of the total adult population of the world. Also, the occurrence of cardiovascular diseases related ailment to obesity and malnutrition has increased globally to 17.8 million deaths every year by 2017. It has been predicted that the population will rise by 2.4 billion people in developing countries of Sub-Saharan Africa and South Asia by 2050 [3]. The four major biological constituents of muscle are protein, fat, moisture and ash, also called the proximate composition of fish. Fish is a particularly important part of the traditional food, and its consumption has been proven to have many health benefits [4]. Nutritional and economic factors were taken into account in the selection of the fish types. According to our findings, dry fish has high-protein content and low-fat which can contribute greatly to children's as well as adult's recommended protein intake. Fish nutritional value varies significantly depending on the species, size, and age of the fish, but understanding the proximate analysis of the fish can provide us with a general idea of its nutritional profile [4].

“Fish is a supreme quality food due to its great nutritional value. Proximate analysis of different fish is carried out by many different researchers worldwide reporting minor discrepancies in them because of the differences in their feeding habits, surrounding temperature, age, season and sex. It is reported that fish provides high quality protein containing all essential amino acids, polyunsaturated omega-3 fatty acids and several micronutrients, including vitamins and minerals”

[5,6]. Fish or fish based products are a very rich source of high quality protein and there are many evidences suggesting their beneficial effects on human health [5].

The consumption of fishery products are preventing cardiovascular and other diseases [6] due to their low fat [5]. “Regularly eating of fish can reduce the risk of cancer and other nutrition related diseases” [7]. “And also lesser risk of Alzheimer's diseases and prevent the cardiovascular diseases” [8]. According to Love [9] “the fish containing 70 to 80% of water, 20 to 30% of protein and 2 to 12% of lipid”. “In different environmental conditions, the fish biochemical composition may influence food and feeding conditions and size & sex of the fish” [10]. The present study deals with the monthly variations of total carbohydrates, total protein, fat, moisture and ash contents in muscle of the proximate composition of *O. mossambica* tissue was investigated with respect to different seasons.

2. MATERIALS AND METHODS

Fish samples were collected from Meghadrigedda reservoir, Visakhapatnam district of Andhra Pradesh and geographical coordinates of 17°45'54"N; 83°11'27"E. And samples were washed with running tap water and the then length and weight was recorded. Non-edible portions of fish were removed immediately to avoid decomposition. The muscles washed, dried and powder for various analyses. To find out the concentrations of the total proteins, total carbohydrates, total fats, moisture and ash in *O. mossambica* samples were collected at regular monthly intervals for a period of one year (June, 2021- May, 2022) at Meghadrigedda reservoir in Visakhapatnam District.

Total Proteins: The protein content of the muscles was estimated by following method of Lowry's [11]. **Total Carbohydrates:** Anthrone in Sulphuric acid can be used for colorimetric determination of sugar; methylated sugars and polysaccharides [12]. **Total Fats:** The total lipids

were extracted from the dry tissues, by following the method of Floch et al.,[13]. Moisture and Ash: Moisture and Ash was determined by AOAC methods [14].

3. RESULTS AND DISCUSSION

The proximate composition of the *O. mossambica* were presented Figs. 1 to 5.




Fig. 1. Monthly variations of total protein content in muscle of male and female *O. mossambica*




Fig. 2. Monthly variations of total Fats content in muscle of male and female *O. mossambica*




Fig. 3. Monthly variations of total Carbohydrates content in muscle of male and female *O. mossambica*




Fig. 4. Monthly variations of Moisture content in muscle of male and female *O. mossambica*

“The muscle composition of total protein, total fat, total carbohydrates, moisture and ash contents showed variations in their different seasons in both females and males”. [49] The highest protein 36.8 ± 3.4 in the month of August in South west monsoon and lowest 24.5 ± 3.2 recorded in May in summer in females and the highest in males was 37.5 ± 3.4 in the month of August in South west monsoon and lowest 24.5 ± 2.9 in the month of April in summer. The highest content of fat was 8.6 ± 1.4 was recorded in Post monsoon in the month of November and the lowest 5.4 ± 1.9 in South west monsoon in the month of September

in case of females and the highest in males was 8.7 ± 1.8 in post monsoon in the month of November and lowest 6.4 ± 0.9 in south west monsoon in the month of July. Total carbohydrates was observed highest 5.5 ± 1.2 in South west monsoon in the month of September and lowest 4.1 ± 0.6 recorded in the North east monsoon in the month of January in case of females. In males, highest 5.9 ± 1.2 was recorded in the South west monsoon in the month of August and the lowest 4.8 ± 1.1 was recorded in North east monsoon in the month of February.




Fig. 5. Monthly variations of Total Ash content in muscle of male and female *O. mossambica*

The highest moisture 84.5 ± 6.5 was observed in Post monsoon in the month of October and minimum 80.2 ± 5.9 was recorded in South west monsoon in the month of July in case of females and in males the highest moisture 85.5 ± 6.9 was recorded in post monsoon in the month of October and the lowest recorded 75.5 ± 5.6 in summer in the month of May. The ash content was highest 4.9 ± 1.5 in North east monsoon and the lowest 4.1 ± 1.2 in South west monsoon in the month of June in females and also males was highest 4.8 ± 1.6 in North East monsoon in the month of December and lowest 4.1 ± 1.7 in South west monsoon in the month of June. The muscle composition of the different fish species showed variation depending on water quality, sexual, feeding habit and these factors are responsible for fish proximate composition [15]. Fishes of various species do not provide the same nutrient profile [16] and the nutritive quality of a fish varies with season [17]. The protein content of the fish is considered as an important tool for evaluation of the physiological status of fishes [18].

3.1 Total Protein

Proteins are high molecular weight polypeptides and fundamental compounds of living things. It contains in addition, non-amino acid materials, lipid or carbohydrate. Proteins play a major role in cell function and cell structure. Amino acids are fundamental bio-molecules in all aspects of cell structure and function [19]. Fish protein is highly digestible and contains a good number of

essential amino acids, thereby having a high biological value. In the present study highest protein recorded in South west monsoon in males and females. The protein content in showed variations and ranged between 24.5 and 36.8% female *O. mossambica*. The nutrient composition of selected food fishes and their protein and fat values studied by Ahmed et al., [20]. Mustafa et al., [21] observed protein content in muscle of *P. sophore* and he stated that protein showed a correlate with spawning period. Banik and Saha, [22] explained a sharp decline in the weight of the gonad after spawning.

3.2 Total Fats

Fats are heterogeneous group of compounds and important dietary constituents with high energy value. They are essential nutrient for the supply of energy for the body. Fatty acids are the building blocks and essential fatty acids like linolenic acid. Nahid et al., [23] recorded that the fat value in muscle of *X. canila*. Bogard et al., [24] reported that the nutrient analysis of different food fishes. Siddique et al., [25] also reported "the fat values of *Puntius sarana* and *Puntius sophore*". Mustafa et al., [26] studied on "fat levels of *Hilsa ilisha* and *P. sarana* during different processing method". Begum et al., [27] reported "the muscle fat high value in *P. sarana*". Maria Zaman et al., [28] recorded "the nutrient levels of selected fish species" and Sarojnalini and Sarjubala Devi [29] studied on "nutritive values of *Puntius sophore*". Bijyalakshmi et al., [30] recorded "low values of total lipid of *Puntius*

sophore". Prabhakara Rao et al., [31], recorded "fat levels in high commercial fishes like rohu (*Labeo rohita*) and channa (*Channa striata*)". "Ecological factors and water condition are influence on the proximate composition particularly fat content of selected fishes" was studied by Iverson, et al., [32].

3.3 Total Carbohydrates

Total Carbohydrates are a group of organic compounds including sugars, starches and fibre, which is a major source of energy for animals. Carbohydrates regarding fishery products are not dietary fibre but only glucides, the majority consist of glycogen. They contain traces of glucose, fructose, sucrose and other mono and disaccharides [33]. The carbohydrate in the muscle of *P. vigil*, was varied from 0.3 to 0.63% [34], 2.4 to 3.4% in *C.smithii* [39], 0.24% in claw meat of *S. serrata* [36] and other crab was studied by Balasubramanian et al., [35]. In the present study the highest (5.9) percentage carbohydrate recorded in South west monsoon in the month of August in males and females was highest 5.5 in South west monsoon in the month of September in *O. mossambica*.

3.4 Moisture

Moisture forms the major component of proximate composition. The moisture values are ranged from 73.43 to 78.86%. Mustafa et al., [26,] find out "moisture values in *P. sophore* and *Hilsa sps*". Maria Zaman et al.,[28] recorded "moisture content of *P. sophore* was 75.71". Bijayalakhami et al., [30] studied "the moisture value of *P.sophore* is 78.95 to 81.34%. In the present study highest moisture values were observed in August 2021 during spawning season of south west monsoon season. The results of the present study revealed that changes in moisture content in the muscles of *O. mossambica* could be attributed to changes in moisture level in spawning period".

3.5 Ash

The concentrations of minerals and trace elements that are contribute to the total ash composition in fishes. Its depending on their food availability and feeding habits, water quality and sex & size of the fish [15] and also season, environment and ecosystem [37,38]. "The ash content in the muscle of marine fishes was little bit higher than that of the freshwater fishes" [39]. Panchakshari et al.,[39] observed "biochemical observations of *Lates calcarifer*". "Sometimes

storage is also influence the ash content changes in the muscle tissue of fishes" [40].

"The intensity of feeding, quality and quantity of natural food, temperature may impact on the protein and fat content in the muscles of the fishes" [41]. Luzia et al., [42] stated that "differences was observe in fat content in the muscle of fish collected in different seasons". "Seasons and food quality may also impact on the muscle of rainbow trout" [43]. Skatecki et al., [44] studied "the effect of the season on fat content in muscle tissue in perch". "In the autumn season contained significantly more fat (0.29%) than those caught in the spring (0.11%)" [45].

"The protein content of muscle has been showed to fluctuate in relation to the maturation and spawning was explained" by Jafri and Khawaja [46]. Piska and Warghray [47] stated that "the seasonal variations in the protein content are well marked and striking correlation with maturation, spawning and feeding activity in fishes." "The high content of protein in pre-spawning may due to the fundamental nitrogen required for the maturation". [49] Piska et al., [48], reported that muscles protein was highest in pre-spawning of *S.bacaila*. The result obtained in this study has provided the detailed knowledge of the proximate composition of the fish *O. mossambica* in relation to spawning season.

4. CONCLUSION

The present study revealed that the seasonal changes in proximate analysis of the muscle of *O.mossambica* regarding the reproductive cycle, storage and utilization of reserves. It's observed that the protein was used for body building and gonadal development during different phases of maturation and fats has been used for energy. From the current study it can be concluded that due to the presence of appreciable amount of protein, fat and carbohydrates and also suitable moisture this fish used for feeds. Hence the fish is suitable as potential industrial material for possible utilization for different products.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Sutharshiny S, Sivashanthini K. Total lipid and cholesterol content in the flesh of the

- five important commercial fishes from waters around Jaffina Peninsula, Sri Lanka. *Int J Biological Chem.* 2011;6:161-169.
2. FAO. The state of world fisheries and aquaculture. Rome: Food and Agriculture Organization of the United Nation. 2018.
 3. World Health Organization (WHO). Obesity and overweight; 2020. Available:<https://www.who.int/news-room/fact-sheets/details/obesity-and-overweight>.
 4. Indur BD, Vishwajit V, Prashant, V. Ambrish. Evaluation of muscle proximate analysis and nutritional value of six freshwater fish species. *Int. J. Environ. Clim. Change.* 2023;13(4):156-168.
 5. Krishna P.V. Haematological and Biochemical observations of *Channa striatus* infected with *Acanthocephala (Pallisentis)* sp). *Indian Journal of Comparative Animal physiology.* Tirupathi. 2006;24(1):38-41.
 6. Marques I, Botelho G, Guiné R. Comparative study on nutritional analysis of fish available in Portugal. *Nutrition & Food Science.* 2019;49(5):925-41.
 7. Sidhu KS. Health benefits and potential risks related to consumption of fish or fish oil. *Regul Toxicol Pharmacol.* 2003; 38:336-344.
 8. Cahu C, Salen E, Lorgeril MD. Farmed and wild fish in the prevention of Cardiovascular diseases: Assessing possible differences in lipid nutritional values. *NutrMetab Cardiovascular Dis.* 2004;14:34-41.
 9. Love RM. *The Chemical Biology of Fishes.* Brown ME (Edn), Academic press New York, USA; 1980.
 10. Oliveira ERN, Agostinho AA, Matsushita M. Effect of biological variables and capture period on the proximate analysis and fatty acid analysis of the dorsal muscle tissue of *Hypophthalmu sedentatus* (Spix, 1829). *Brazilian Arch. Biol Technol.* 2003;46:105-114.
 11. Lowry OH, Rosebrough JN, Farr AL, Rose J. Randall. Protein measurement with the Folin phenol reagent. *J. Biol. Chem.*1951:193:265-275.
 12. Dubois M, Gilles KA, Hamilton JK, Rebers PA, Smith F. Colorimetric method for determination of sugars and related substances. *Analytical Chemistry.* 1956; 28:350-356. Available:<http://dx.doi.org/10.1021/ac60111a017>
 13. Folch J, Lees J, Sloane Stanley, G. H. A simple method for the isolation and purification of total lipid from animal tissues. *J. Biolo. Chem.* 1957;266:497-509.
 14. Association of Official Analytical Chemists. Official methods of analysis, 17th Edition, The association of official analytical chemists, gaitersburg, MD, USA; 2000.
 15. Krishna PV, Gopi G, Hemanth Kumar V, K. Bhanu Prakash. Comparative study of meat analysis of Catlacatla, *Labeo rohita*, *Cirrhinus mrigala* and *Pangasius hypophthalmus* under different treatments. *International Journal of Advanced Research.* 2015;3(10):480–485.
 16. Ravishankar P, Arivind Kumar. Seasonal changes in the biochemical analysis of muscle in relation to breeding cycle in carp minnow, *Salmostoma bacaila* (Ham.) from Himayat Sagar reservoir, Hyd. (India). *U.P. J. Zool.* 1989;9(1):39-42.
 17. Medford BA Mackay, WC. Protein and lipid content of gonads, liver and muscle of northern pike (*Esox lucius*) in relation to gonad growth. *J. Fisheries Res. Board of Canad.* 1978;35:213-219.
 18. Jafri, A.K. and Khawaja, D.K. Seasonal changes in the biochemical analysis of the freshwater murrel *Ophiocephalus punctatus* (Bloch). *Hydrobiologia.*1968; 32:206-213.
 19. Krishna PV. Biochemical observations on *Heteropneustis fossilis*. Infected with epizootic ulcerative syndrome (EUS). *J. Ecobiology.* 2002;14(2):125–129.
 20. Ahmed S, Arifur Rahman, AFM, Mustafa MG, Hossain MB, Nahar N. Nutrient analysis of indigenous and exotic fishes of rain fed waterlogged paddy fields in Lakshmipur, Bangladesh. *World J. Zool.* 2012;7(2):135-140.
 21. Mustafa MG, Yasmin T, Sharif Nilla, S Rahman Khan, Rahman Khan MM. M.A. Effect of freezing period on the microbial and nutrient quality of pool barb, *Puntius sophore* from Dhaka city retail market, Bangladesh. 2013;41(1):1-11.
 22. Banik S, Saha, S. Reproductive physiology and feeding biology of *Puntius sophore* of Tripura, India. *Res. Environ. Life Sci.* 2013;6 (4):115-118.
 23. Nahid MN, Latifa GA, Farid FB, Begum M. Evaluation of biochemical analysis of salt and garlic treated smoke- dried Chapila (*Gudusia chapra* Hamilton, 1822) and Kaika (*Xenentodon cancila* Hamilton-

- Buchanan,1822) Fish at laboratory condition(27-31o C). Res. J. Animal, Veterinary and Fishery Sci. 2014;2(10):10-15.
24. Bogard JR, Thilsted SH, Marks GC, Wahab MA, Hossain MAR, Jakobsen J, Stangoulis J. Nutrient analysis of important fish species in Bangladesh and potential contribution to recommended nutrient intakes. J. Food Analysis and Analysis. 2015;42:120–133.
 25. Siddique MN, Hasan MJ, Rahman MZ, Islam MR, Boduruzaman M, Reza MF, Ray S. Effect of freezing time on nutritional value of Jatpunti (*Puntius sophore*), Sarpunti (*P. sarana*) and Thaisarpunti (*P. gonionotus*). Bangladesh Res. Pub. J. 2011;5(4):387-392.
 26. Mustafa MG, Begum SR, Abdul Khaleque M, Jannat M, Ahsan DA. Nutritional qualities of Hilsha and Sarpunti in different salt curing methods. DhakaUniv. J. Biol. Sci. 2012;21(1): 97-104.
 27. Begum M, Minar MH. Comparative study about body analysis of different SIS, Shell fish and Ilish; commonly available in Bangladesh. Trends in Fisheries Research, 2012;1(1):38-42.
 28. Maria Zaman, Naser MN, Abdullah ATM, Nasima Khan. Nutrient contents of some popular freshwater and marine fish species of Bangladesh. Bangladesh J. Zool. 2014;42(2):251-259.
 29. Sarojnalini CH, Sarjubala Devi W. Antioxidant properties and nutritive values of raw and cooked pool barb (*Puntius sophore*) of Eastern Himalayas. Int. J. Biol., Food, Veterinary and Agricul. Eng. 2014;8(1):8-12.
 30. Bijayalakshmi Ch, Romen Singh N, Indira N, Shomorendra M. Estimation of moisture and total lipid content of some small indigenous fishes of Manipur. Inter. J. Sci. and Res. 2014;3(12):1091-1094.
 31. Prabhakara Rao PG, Jyothirmayi T, Karuna MSL, Prasad RBN. Studies of lipid profiles and fatty acid analysis of roe from Rohu (*Labeorohita*) and murrel (*Channa striatus*). J. Oleo Sci. 2010;59(10):515-519.
 32. Iverson SJ, Frost KJ, Lang SLC. Fat content and fatty acid analysis of forage fish and invertebrates in prince william sound, Alaska: factors contributing to among and within species variability. Marine Ecol. Progress series. 2002;241:161–181.
 33. Pilla S, Konathala RMR, KSR. Histology and histopathology of the lutjanus johnii and lutjanus russelli from Visakhapatnam coast. IOSR journal of pharmacy and Biological Sciences (IOSR-JPBS). 2014;9(3):35-42.
 34. Ravichandran S, Kumaravel K, Pamela Florence E. Nutritive analysis of some edible fin fishes. International Journal of zoological Research. 2011;7(3):241-251.
 35. Balasubramanian CP, Suseelan C. Biochemical analysis of the deep water crab *charybdis smithii*. Indian J. Fish. 2001;48(3):333-335.
 36. Rao PY, Sirisha IR. Changes in the muscle biochemical analysis of *Lagocephalus spadiceus* (Richardson, 1845) and *Lagocephalus lunaris* (Bloch and Schneider, 1801) off Visakhapatnam, East Coast of India. International Journal of Scientific and Research Publications. 2013;7(3):1-6.
 37. Andres S, Ribeyre F, Toureneq JN, Boudou. Interspecific comparison of cadmium and zinc contamination in the organs of four fish species along a polymetallic pollution gradient (Lot River, France). Sci Total Environ. 2000;284:11-25.
 38. Abdallah, Mohamed. M.A. Speciation of trace metals in coastal sediments of El- max bay south mediterranean sea-west of alexandria, (Egypt). Environ. Monitor.Assess. 2007;132:111-113.
 39. Panchakshari V PV, Krishna K, Madhusudhana Rao, Prabhavathi K. Microbiological and biochemical observations of *Lates calcarifer* in different types of processing with Hurdle technology. International Journal of Advanced Research. 2015;3(4):1232-1235.
 40. Hassan, MN Rahman, Hossain M, MM Newsad AAKM, Hossain MB. Post-harvest loss and shelf life of traditionally smoked shrimp products produced in Bangladesh. World J. of Fish. and Marine Sci. 2013;5(1):14-19.
 41. Puchala R, Pilarczyk M. Nutrition effect on chemical analysis of carp meat. Agri.Eng. 2007; 93(5),363–368.
 42. Luzia LA, Sampaio GR, Castellucci CMN, Torres EAFS. The Influence of season on the lipid profiles of five commercially

- important species of brazilian fish. Food Chem. 2003;83(1):93–97.
DOI: 10.1016/S0308-8146(03)00054-2.
43. Skąłeckı P, Staszowska A, Kaliniak A, Florek M. Utility value and meat quality of rainbow trouts (*Oncorhynchus mykiss* Walb.) from extensive and intensive farming. Sci. Ann. Polish Soc. Animal Product. 2013;9(3):59–67.
44. Skąłeckı P, Florek M, Litwińczuk A, Zaborska A. Utility value and meat quality of rainbow trout (*Oncorhynchus mykiss*) with regard to the weight of fish. Sci. Ann. Polish Soc. Animal Product. 2013;9(1):69–73.
45. Skąłeckı P, Florek M, Staszowska A. effect of fishing season on value in use, intrinsic properties, proximate analysis and fatty acid profile of perch (*Perca fluviatilis*) muscle tissue. Arch. Pol. Fish. 2013;21(4): 249–257.
46. Jafri AK, Khawaja DK. Seasonal changes in the biochemical analysis of the freshwater murrel *Ophiocephalus punctatus* (Bloch). Hydrobiologia. 1968;32:206-213.
47. Piska RS, Waghray, S. Some aspects of reproductive biology of amblypharynaadon mola (Ham) geobios. 1986;13(5):204-207.
48. Piska RS, Naik BR, Diwakara Chary K. Variation in Biochemical analysis in muscles of AmblypharynaadonMola (Ham) with reference to its breeding cycle. J. Aqua. Biol.1998;13(1 &2):86-88.
49. Krishna PV, Kumar MN, Sarma BA, Dedeepya P. Proximate composition and its seasonal variations of the muscle tissue of *Channa striata* from Krishna river, Andhra Pradesh. International Journal of Fisheries and Aquatic Studies. 2023;11(1):109-115.

© 2023 Krishna et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/110277>