

Effect of Titanium Dioxide Nanoparticles on Growth and Biomass Accumulation in Lettuce (*Lactuca sativa*)

Channa B. Rajashekar*, Brenden Armstrong

Department of Horticulture and Natural Resources, Kansas State University, Manhattan, KS, USA

Email: *crajashe@ksu.edu

How to cite this paper: Rajashekar, C.B. and Armstrong, B. (2024) Effect of Titanium Dioxide Nanoparticles on Growth and Biomass Accumulation in Lettuce (*Lactuca sativa*). *American Journal of Plant Sciences*, 15, 1-13.

<https://doi.org/10.4236/ajps.2024.151001>

Received: November 19, 2023

Accepted: January 2, 2023

Published: January 5, 2023

Copyright © 2024 by author(s) and Scientific Research Publishing Inc. This work is licensed under the Creative Commons Attribution International License (CC BY 4.0).

<http://creativecommons.org/licenses/by/4.0/>



Open Access

Abstract

The use of titanium dioxide nanoparticles (nTiO₂) is gaining interest in agriculture because of their impact on many aspects of plant growth. The present study examines the effects of nTiO₂ (5 nm and 10 nm) applied to seeds and the seedlings as a foliar application on various aspects of growth characteristics and biomass accumulation in lettuce (*Lactuca sativa*, cv. Grand Rapids). Application of 10 nm nTiO₂ to seeds through imbibition resulted in a significant reduction in shoot biomass accumulation while 5 nm nTiO₂ did not affect the biomass accumulation in lettuce. The application of 10 nm nTiO₂ reduced the fresh shoot biomass accumulation by about 18% compared to the control plants. Other growth characteristics such as shoot dry biomass, root fresh and dry biomass, plant height, and leaf area were not affected by the application of both 5 nm and 10 nm nTiO₂. In addition, foliar application of these nanoparticles to the lettuce seedlings did not have a significant effect on most of the growth parameters examined, and the increasing concentration ranging from 5 nm/L to 400 mg/L did not produce a consistent response in lettuce. Thus, nTiO₂ application to lettuce seeds had a notable negative impact on shoot growth while foliar application did not have a significant effect on many plant growth characteristics. However, foliar applications produced some symptoms of toxicity to the foliage in the form of necrotic or chlorotic patches on the leaves, which were more pronounced with increasing concentrations of both 5 nm and 10 nm nTiO₂. However, these symptoms were apparent at a concentration as low as 50 mg/L of nTiO₂. Thus, foliar application of nTiO₂ may not have a significant impact on many of the growth characteristics in lettuce, but it can result in foliar toxicity.

Keywords

Growth Characteristics, Lettuce, Nanoparticles, Titanium Dioxide, Toxicity

1. Introduction

There is an increasing interest in using titanium dioxide nanoparticles (nTiO₂) in crop production to improve the overall growth of plants and enhance the yield potential of many cultivated crops [1]. Many studies show a positive impact of nTiO₂ on various aspects of plant growth characteristics such as improved seed germination, photosynthetic efficiency, root growth, biomass accumulation, nutrient assimilation, plant resilience under stress, and the overall growth and performance of many crop plants [1] [2] [3] [4] [5].

Titanium dioxide is a naturally occurring mineral and is considered safe and hence extensively used in many industrial and household products such as paints, construction materials, cosmetics, medicine, and many food products [1] [6]. TiO₂ nanoparticles show photocatalytic properties and produce reactive oxygen species in the presence of UV and even under visible light with some modification of pristine TiO₂ [7]. The photocatalytic property of nTiO₂ and the resulting production of oxidative species has been harnessed extensively in the reclamation of soil, water, and air contaminated with pollutants, and as an antimicrobial agent to improve food safety and in the self-sterilization of surfaces [6] [8] [9] [10] [11].

Despite the increasing interest in the application of nTiO₂ in agriculture to improve plant growth and performance, its effects on many aspects of plant growth are not clearly understood because plant responses to nTiO₂ are complex and widely variable. Several plant factors can impact the effects of nTiO₂ depending on plant species, type of plant tissues, growing conditions, and plant-environment interactions [4] [5] [12] [13]. Several studies have shown a positive impact of nTiO₂ on many aspects of plant growth [1] [14] [15] [16] [17]. However, some studies show either negative effects [18] [19] [20] or no effects [14] [21] [22] [23] of nTiO₂ in many crop species. Thus, there is no clear understanding of the factors that may contribute to the differential responses to nTiO₂ in crop plants including lettuce.

In addition to plant factors, many nanoparticle characteristics can notably influence the impact of nTiO₂ on plant growth. For example, particle size can significantly influence the absorption, translocation, and accumulation of nTiO₂ in various plant parts, which can affect the plant responses to nTiO₂. Smaller molecules are typically taken up by the plants and translocated more readily than the larger ones [24]. Studies on wheat by Larue *et al.* [24] showed that plants may have a limiting threshold of nanoparticle size for absorption and the subsequent translocation of nTiO₂ to other parts of the plants. Similarly, the concentration of nTiO₂ also has a significant impact on plants' responses. Typically, lower concentrations of nTiO₂ are known to produce a positive impact on many plant functions [1] [14] while on the other hand, higher concentrations have been shown to produce a negative impact on plant growth [20] [25]. Furthermore, high concentrations of nTiO₂ can also lead to the production of oxidative stress and toxicity in plants due to their photocatalytic activity [26] [27] [28].

The negative impact of the photocatalytic activity of nTiO₂ and the resulting oxidative stress has been demonstrated in many plant species and soil microbiota [13] [29] [30] [31] [32] [33]. However, there is very little known about the impact of nTiO₂ nanoparticle size and its concentration on the growth of lettuce plants.

Therefore, the focus of the present study was to examine the effect of various concentrations of two different sizes of nTiO₂, applied to seeds and to seedlings as foliar sprays, on various growth characteristics and biomass accumulation in lettuce plants.

2. Materials and Methods

The study consisted of two greenhouse trials to determine the impact of nTiO₂ nanoparticles on the growth of lettuce plants (*Lactuca sativa*, cv. Grand Rapids) in Manhattan, KS during 2023. Lettuce seeds were obtained from Ferry-Morse Seed Co. (Fulton, KY, USA), and the TiO₂ nanoparticles (anatase; 5 nm and 10 - 30 nm, referred to herein as 10 nm) were obtained from SkySpring Nanomaterials, Inc. (Houston, TX, USA) and were stored in the dark in desiccators until use. Treatments of nTiO₂ consisting of two nanoparticle sizes were applied either through seed imbibition or as foliar sprays to the seedlings.

2.1. Seed Imbibition

Aqueous suspensions of nTiO₂ (5 nm and 10 nm) were used for all the treatments. Lettuce seeds were imbibed with water (control) or nTiO₂ suspensions (250 mg/L) for up to 12 hours and placed on moist paper towels in Petri dishes for germination. The nTiO₂ suspensions were freshly prepared and thoroughly agitated before all the treatment applications. Following the germination (after 10 days of sowing), the seedlings were transplanted into a peat-based mix growing medium, BM6 (Berger, Quebec, Canada) contained in 15 cm plastic pots (1 seedling/pot) (HC Companies, Twinsburg, OH, USA). The plants were grown in a greenhouse maintained approximately at 20°C with solar radiation as the light source. Plants were watered every other day and fertigated approximately once a week with 100 mL of solution/plant containing 250 ppm N (Jack's Peat-Lite, NPK: 20-10-20, JR Peters, Inc., Allentown, PA, USA). Plants were harvested after 4-5 weeks of transplanting. The experiment was conducted on a completely randomized design with 6 replications for each nanoparticle size.

2.2. Foliar Spray Treatments

In the first greenhouse study, the lettuce seeds were imbibed in water for up to 12 hours and directly sown in the moist peat-based growing medium (BM6) on July 17, 2023. After germination, the seedlings were thinned to retain one plant/pot. Seedlings were watered every other day and fertigated once a week as described above. The 20-day-old lettuce seedlings were sprayed with freshly prepared nTiO₂ suspensions (5 nm and 10 nm) at 5, 10, 50, 100, 150, mg/L. Each

plant was evenly sprayed with water (control) or nTiO₂ suspension ensuring its uniform coverage of the canopy and avoiding nTiO₂ contamination of the growing medium. The plants were irrigated before the foliar applications of nTiO₂ suspensions. The 5-week-old plants were harvested to measure their growth characteristics. The experiment was conducted on a completely randomized design with 6 replications.

A second greenhouse study was conducted to assess the impact of higher concentrations of nTiO₂ on the growth of lettuce plants. Lettuce seeds were imbibed in water for 12 hours and were germinated and then, transplanted into the peat-based growing medium as described above. After thirteen days of transplanting, seedlings were treated with a foliar spray of freshly prepared nTiO₂ (5 nm and 10 nm) suspensions at 500 mg/L, 2500 mg/L, and 4000 mg/L as described above. With the high concentration of TiO₂, agglomeration can be a problem, so, care was taken to apply well-dispersed nTiO₂ evenly over the foliage. The control plants were sprayed with water. Plants were irrigated every other day and fertigated once a week as described above. Plants were irrigated before the foliar applications of nTiO₂ suspensions. The experiment was conducted on a completely randomized design with 6 replications for each nanoparticle size.

2.3. Plant Growth Measurements

All plant growth measurements were made on the plants grown from seeds imbibed with nTiO₂ and on those treated with nTiO₂ through foliar sprays before harvest.

As the lettuce variety has a loose-leaf canopy, plant height was measured from the base of the stem to the tallest leaf. The other growth characteristics measured were the number of fully opened leaves/plant and the leaf area of all the fully opened leaves using a leaf area meter (LI-3100, LI-COR, Inc., Lincoln, NE). To measure the plant biomass, plant shoots were separated from roots. The roots were washed free of the growing medium and were blotted dry with paper towels to remove the surface moisture. Shoot and root fresh weights from each plant were measured. They were then transferred to a drying oven at 70°C and were dried for 4 days or until a constant weight was achieved before measuring the dry weights.

2.4. Statistical Analyses

All the experiments were conducted on a completely randomized design with 6 replications with nTiO₂ nanoparticle sizes and their various concentrations as treatment variables. The Tukey HSD test was conducted at alpha 0.05 and *p-values* were obtained to perform multiple comparisons of the treatment means.

3. Results and Discussion

3.1. Seed Imbibition and Plant Growth

The response of lettuce plants grown from seeds treated with 5 nm and 10 nm

nTiO₂ in relation to various plant growth characteristics including shoot fresh and dry biomass, root fresh and dry biomass, plant height, leaf area, and leaf number was studied at the time of harvest.

Lettuce plants grown from the seeds imbibed with nTiO₂ at 250 mg/L produced variable responses. The fresh shoot biomass of lettuce plants was significantly reduced by the seed treatment with 10 nm nTiO₂ by 18.1% compared to the control while, on the contrary, there was no significant impact on the fresh shoot biomass in plants receiving seed treatment with 5 nm nanoparticles (**Figure 1(a)**). This was partly due to a significant reduction in the number of leaves/ plant with 10 nm nTiO₂ seed treatment (**Figure 1(b)**). This shows that larger nanoparticle size had a significant negative impact on shoot biomass in lettuce compared to the smaller particle size. However, it should be noted that the size of nanoparticles can impact not only their absorption but more importantly their translocation leading to their differential accumulation in various plant parts [24].

Many growth characteristics including plant height, dry shoot biomass, leaf area, and both fresh and dry root biomass were not affected by the seed treatment of either 5 nm or 10 nm nanoparticles (**Table 1** and **Figure 1(b)**). Also, there was no significant impact on the seed germination in response to the different nanoparticle sizes (data not shown). Similar results were observed in kidney beans where nTiO₂ did not affect the germination but had a significant impact on early root growth [14].

Although most of the growth characteristics were unaffected by nTiO₂ treatment, it is noteworthy that the negative impact of nTiO₂ (10 nm) on fresh shoot biomass has practical implications for the growers as it can notably affect the marketable yield. The reduction of shoot growth (biomass accumulation) when




Figure 1. Growth characteristics of lettuce plants grown from seeds treated with nTiO₂ nanoparticles of different sizes. Lettuce seeds were imbibed with nTiO₂ (5 nm and 10 nm; 250 mg/L) or water (control) for up to 12 hours, and the shoot fresh weight (FW) (a), leaf area and number of leaves/plant (b) were measured at the time of harvest. The mean values with their standard deviations (SD) are presented. Significant differences at $p \leq 0.05$ are indicated by letters.

nTiO₂ was applied to seeds may indicate its negative impact on the early growth of lettuce [14]. Similar observations on the negative effects of seed treatment of nTiO₂ have been made in other species including corn, cucumber, oats, onion, and tomato [15] [34].

3.2. Foliar Spray and Plant Growth

In the first greenhouse study, lettuce seedlings were treated with both 5 nm and 10 nm nTiO₂ nanoparticles at concentrations ranging from 5 mg/L to 150 mg/L as a foliar spray.

Foliar spray of 5 nm nTiO₂ did not have any significant impact on many of the growth characteristics of lettuce except the number of leaves per plant (Table 2). Plants treated with 10 mg/L and 150 mg/L of nTiO₂ (5 nm) produced more leaves/plant compared to the control plants, the highest leaf number/plant being at 150 mg/L of nTiO₂. The responses to other nTiO₂ concentrations were not

Table 1. Growth characteristics of lettuce plants grown from seeds treated with nTiO₂ nanoparticles of different sizes. Lettuce seeds were imbibed with nTiO₂ (5 nm and 10 nm; 250 mg/L) or water (control) for up to 12 hours and the growth measurements of plants were made at the time of harvest. The mean values with their SD and *p-values* are presented.

	Plant height (cm)	Shoot DW* (g)	Root FW* (g)	Root DW* (g)
Control	20.6 ± 2.1	7.2 ± 1.4	29.8 ± 5.5	3.3 ± 1.0
nTiO ₂ , 5 nm	21.3 ± 2.7	6.4 ± 0.8	34.3 ± 8.5	3.5 ± 0.7
nTiO ₂ , 10 nm	19.3 ± 3.9	5.8 ± 0.9	35.6 ± 3.4	3.5 ± 0.7
<i>p-value</i>	0.953	0.150	0.530	0.925

*FW and DW represent the fresh weight and dry weight, respectively.

Table 2. Effect of foliar spray of lettuce plants with nTiO₂ (5 nm) on growth characteristics of lettuce plants. Lettuce plants were sprayed with nTiO₂ (5 nm) or water (control). The growth measurements were made at the time of harvest. The mean values with their SD and *p-values* are presented.

nTiO ₂ 5 nm	Shoot FW* (g)	Shoot DW* (g)	Plant height (cm)	Leaf number/ plant	Leaf area (cm ²)
Control	43.6 ± 12.0	2.9 ± 0.8	20.5 ± 1.7	8.6 ± 1.1	636 ± 158.8
5 mg/L	42.3 ± 2.0	3.8 ± 0.5	21.7 ± 2.8	10.2 ± 0.4	674 ± 191.1
10 mg/L	47.8 ± 12.5	4.5 ± 1.6	22.2 ± 3.7	12.8 ± 2.3*	812 ± 131.7
50 mg/L	45.2 ± 9.9	3.6 ± 0.9	26.2 ± 4.7	11.4 ± 2.0	606 ± 89.5
100 mg/L	38.0 ± 5.9	3.9 ± 0.6	22.2 ± 3.7	11.2 ± 1.7	601 ± 97.9
150 mg/L	54.8 ± 9.6	4.9 ± 1.1	25.0 ± 4.8	13.8 ± 2.4*	796 ± 109.0
<i>p-value</i>	0.278	0.067	0.174	0.003	0.063

*FW and DW represent the fresh weight and dry weight, respectively.

different from those in the control plants. Unlike the seed treatment of nTiO₂ (10 nm), the foliar sprays did not result in the reduction of shoot fresh biomass compared to the control plants (**Figure 2(a)**). Although there was an increase in shoot dry biomass in response to nTiO₂ (10 nm) at 10 mg/L, both shoot fresh and dry biomass accumulations at higher concentrations of nTiO₂ were similar to the those in the control.

Similarly, nTiO₂ concentrations did not have an impact on the leaf area compared to the control (**Figure 2(b)**). Overall, these observations suggest that the foliar application of nTiO₂ up to 150 mg/L did not significantly impact most of the growth characteristics in lettuce plants. Therefore in a separate study, we used higher concentrations of nTiO₂ as a foliar spray, to determine their impact on the growth of lettuce plants. Foliar sprays of both 5 nm and 10 nm nTiO₂ at concentrations ranging from 500 mg/L to 4000 mg/L were applied to the lettuce seedlings. Again, these treatments produced variable results on the shoot and root biomass with regard to the nanoparticle concentrations. The notable response of lettuce at higher concentrations of nTiO₂ was the reduction in fresh shoot biomass with 5 nm nTiO₂ and in dry shoot biomass with 10 nm nTiO₂ (**Figure 3(a)** and **Figure 3(b)**).

Fresh shoot biomass was reduced by approximately 20% compared to that in the control plants with the application of 5 nm nTiO₂ at 500 mg/L. However, 10 nm nTiO₂ at 500 mg/L reduced the shoot dry biomass by about 29% compared to that in the control plants (**Figure 3(a)** and **Figure 3(b)**).

However, these responses were variable at higher concentrations. Foliar applications of 5 nm or 10 nm nTiO₂ did not have any impact on other growth characteristics such as plant height, leaf area, root fresh and dry biomass (**Table 3** and **Table 4**). Our results support the previous studies that show no impact of nTiO₂ on many aspects of plant growth [3] [14]. Furthermore, previous studies have also shown that nTiO₂ can have a negative impact on plant growth and even be toxic, especially at higher concentrations [17]. Hu *et al.* [25] showed that




Figure 2. Effect of foliar spray with 10 nm nTiO₂ on growth characteristics of lettuce plants. Lettuce seedlings were sprayed with nTiO₂ (10 nm), and the growth measurements of plants were made at the time of harvest. Control plants were treated with water. The mean values for the shoot fresh weight (FW) and dry weights (DW) (a) and leaf area (b) with their SD are presented. Significant differences at $p \leq 0.05$ are indicated by letters.




Figure 3. Effect of foliar spray with nTiO₂ (5 nm and 10 nm) on growth characteristics of lettuce plants. Lettuce seedlings were sprayed with nTiO₂ (5 nm and 10 nm) or water (control), and the growth measurements of plants were made at the time of harvest. The mean values for the leaf area and shoot fresh weight (FW) for 5 nm nTiO₂ (a), shoot dry weight (DW) and leaf number for 10 nm nTiO₂ (b) with their SD are presented. Significant differences at $p \leq 0.05$ are indicated by letters.

Table 3. Effect of foliar spray of lettuce plants with 5 nm nTiO₂ on growth characteristics of lettuce plants. Lettuce seedlings were sprayed with nTiO₂ (5 nm) or water (control). The growth measurements were made at the time of harvest. The mean values with their SD and *p-values* are presented.

nTiO ₂ 5 nm	Shoot DW* (g)	Plant height (cm)	Leaf number/plant	Root FW* (g)	Root DW* (g)
Control	7.2 ± 1.4	20.6 ± 2.1	13.2 ± 0.7	29.8 ± 5.5	3.3 ± 1.0
500 mg/L	5.3 ± 0.9	21.3 ± 2.9	12.3 ± 0.8	34.8 ± 12.9	3.6 ± 1.6
2500 mg/L	6.2 ± 0.6	21.7 ± 2.1	14.5 ± 2.8	25.4 ± 2.4	3.6 ± 1.6
4000 mg/L	6.1 ± 1.0	18.8 ± 2.3	12.3 ± 1.4	31.7 ± 4.1	3.2 ± 0.5
<i>P-value</i>	0.048	0.683	0.096	0.511	0.560

*FW and DW represent the fresh weight and dry weight, respectively.

Table 4. Effect of foliar spray of lettuce plants with 10 nm nTiO₂ on growth characteristics of lettuce plants. Lettuce seedlings were sprayed with nTiO₂ (10 nm) or water (control). The growth measurements were made at the time of harvest. The mean values with their SD and *p-values* are presented.

nTiO ₂ 10 nm	Shoot FW* (g)	Plant height (cm)	Leaf area (cm ²)	Root FW* (g)	Root DW* (g)
Control	69.8 ± 5.9	20.6 ± 2.1	1403.8 ± 139.1	29.8 ± 5.5	3.3 ± 1.0
500 mg/L	58.0 ± 5.2	21.2 ± 4.1	1310.7 ± 149.8	41.6 ± 11.2	4.3 ± 1.5
2500 mg/L	61.9 ± 8.7	19.3 ± 2.4	1290.3 ± 254.9	31.6 ± 1.5	3.1 ± 0.5
4000 mg/L	58.6 ± 8.8	20.7 ± 3.4	1374.5 ± 258.7	32.3 ± 1.7	3.1 ± 0.3
<i>p-value</i>	0.109	0.809	0.531	0.240	0.419

*FW and DW represent the fresh weight and dry weight, respectively.

the effects of nTiO₂ were dose-dependent in lettuce plants in that its higher concentrations suppressed chlorophyll content, biomass, and the nutritional quality of lettuce.

Considering the observation that the negative impact of plant growth (biomass accumulation) by nTiO₂ when it was applied to the seeds while foliar sprays produced little or no impact on overall growth may suggest that early stages of seedling growth including the germination process are more sensitive to nTiO₂ [14]. This is further supported by previous studies that find early growth in many plants including corn, oats, cabbage, and oilseed rape is sensitive to nTiO₂ [14] [15].

Furthermore, foliar applications of nTiO₂ to lettuce seedlings, regardless of their size, showed some signs of toxicity. The foliar applications resulted in many brown necrotic spots or etiolated blotches on the leaves, which were more pronounced at higher concentrations of nTiO₂ (Figure 4). However, these symptoms were apparent with the foliar application of nTiO₂ even at concentrations as low as 50 mg/L. It is noteworthy that although the growth responses were variable with foliar application of nTiO₂, the higher concentrations of nTiO₂ produced signs of toxicity in lettuce. These observations support many previous studies involving the applications of high concentrations of nTiO₂ that result in injury caused by oxidative stress in several plant species [25] [28] [31] [35]. As nTiO₂ is a photocatalytic substance, it can generate reactive oxygen species in the presence of light, especially UV, which can be highly damaging to cells [23]. The primary effect of oxidative species induced by nTiO₂ is the damage to the chloroplast membranes resulting not only in reduced photosynthetic function but also in promoting autophagy, a process through which impaired cell organelles could be eliminated [31] [36].



Figure 4. Effect of nTiO₂ foliar spray on lettuce plants. Foliar damage to leaves (highlighted in red rectangle) by nTiO₂ (5 nm) at 50 mg/L (left) and 100 mg/L (right).

Similar findings of the negative impact of nTiO₂ on photosynthetic function and the overall growth in many plant species including lettuce have been reported [20] [25] [31] [32]. Our results on lettuce show that the application of both 5 nm and 10 nm nTiO₂ as a foliar spray can cause toxicity to the leaves in lettuce even at moderate concentrations of nTiO₂ (50 mg/L or higher).

4. Conclusion

The impact of two nanoparticle sizes of nTiO₂ applied through seed imbibition

and as a foliar spray at various concentrations (5 mg/L to 4000 mg/L) on the growth characteristics of lettuce was studied in a greenhouse trial. Seed imbibition of nTiO₂ (10 nm) significantly reduced shoot biomass accumulation in lettuce plants. However, the other growth characteristics including dry shoot biomass and both fresh and dry root biomass were not affected by nTiO₂ treatments. The foliar applications of nTiO₂ showed variable responses with regard to many of the growth parameters studied. Considering the observation that there was a significant negative impact on shoot growth with seed treatment but with no significant effects on many growth characteristics with foliar applications, early growth in lettuce appears to be more sensitive to nTiO₂ than the later stages of plant growth. However, foliar spray, especially at higher concentrations (≥50 mg/L), showed signs of toxicity as evidenced by localized leaf necrotic and etiolated patches. Thus, the results show that lettuce plants are more sensitive to nTiO₂ when applied to seeds, suppressing shoot growth, while the foliar applications had no significant effect on many of the growth characteristics but can produce foliar toxicity.

Conflicts of Interest

The authors declare no conflicts of interest regarding the publication of this paper.

References

- [1] Lyu, S., Wei, X., Chen, J., Wang, C., Wang, X. and Pan, D. (2017) Titanium as a Beneficial Element for Crop Production. *Frontiers in Plant Science*, **8**, Article No. 597. <https://doi.org/10.3389/fpls.2017.00597>
- [2] Yang, F., Hong, F., You, W., Liu, C., Gao, F., Wu, C. and Yang, P. (2006) Influences of Nano-Anatase TiO₂ on the Nitrogen Metabolism of Growing Spinach. *Biological Trace Element Research*, **110**, 179-190. <https://doi.org/10.1385/BTER:110:2:179>
- [3] Daghan, H., Gulmezoglu, N., Koleli, N. and Karakaya, B. (2020) Impact of Titanium Dioxide Nanoparticles (TiO₂-NPs) on Growth and Mineral Nutrient Uptake of Wheat (*Triticum vulgare* L.). *Biotech Studies*, **29**, 69-76. <https://doi.org/10.38042/biost.2020.29.02.03>
- [4] Choi, H.G. (2021) Influence of TiO₂ Foliar Spray Application on Photosynthesis and Chlorophyll Fluorescence of Strawberry during Low Light Intensity. *Acta Horticulturae*, **1309**, 247-252. <https://doi.org/10.17660/ActaHortic.2021.1309.36>
- [5] Vatankhah, A., Aliniaiefard, S., Moosavi-Nezhad, M., Abdi, S., Mokhtarpour, Z., Reezi, S., Tsaniklidis, G. and Fanourakis, D. (2023) Plants Exposed to Titanium Dioxide Nanoparticles Acquired Contrasting Photosynthetic and Morphological Strategies Depending on the Growing Light Intensity: A Case Study in Radish. *Scientific Reports*, **13**, Article No. 5873. <https://doi.org/10.1038/s41598-023-32466-y>
- [6] Mohajerani, M., Burnett, L., Smith, J.V., Kurmus, H., Milas, J., Arulrajah, A., Horpiulsuk, S. and Kadir, A.A. (2019) Nanoparticles in Construction Materials and Other Applications, and Implications of Nanoparticle Use. *Materials*, **12**, Article No. 3052. <https://doi.org/10.3390/ma12193052>
- [7] Etacheri, V., Valentin, C., Schneider, J., Bahnemann, D. and Pillai, S.C. (2015) Visible-Light Activation of TiO₂ Photocatalysts: Advances in Theory and Experiments. *Journal of Photochemistry and Photobiology C: Photochemistry Reviews*, **25**, 1-29. <https://doi.org/10.1016/j.jphotochemrev.2015.08.003>

- [8] Wang, T.C., Lu, N., Li, J. and Wu, Y. (2011) Plasma-TiO₂ Catalytic Method for High-Efficiency Remediation of *p*-Nitrophenol Contaminated Soil in Pulsed Discharge. *Environmental Science and Technology*, **45**, 9301-9307. <https://doi.org/10.1021/es2014314>
- [9] Long, M., Wang, J., Zhuang, H., Zhang, Y. and Wu, H. (2014) Performance and Mechanism of Standard Nano-TiO₂ (P-25) in Photocatalytic Disinfection of Foodborne Microorganisms—*Salmonella typhimurium* and *Listeria monocytogenes*. *Food Control*, **39**, 68-74. <https://doi.org/10.1016/j.foodcont.2013.10.033>
- [10] Kühn, K., Chaberny, I.F., Massholder, K., Stickler, M., Benz, V.W., Sonntag, H. and Erdinger, L. (2003) Disinfection of Surfaces by Photocatalytic Oxidation with Titanium Dioxide and UVA Light. *Chemosphere*, **53**, 71-77. [https://doi.org/10.1016/S0045-6535\(03\)00362-X](https://doi.org/10.1016/S0045-6535(03)00362-X)
- [11] Zhan, C., Li, Y., Sharma, P.R., He, H., Sharma, S.K., Wang, R. and Hsiao, B.S. (2019) A Study of TiO₂ Nanocrystal Growth and Environmental Remediation Capability of TiO₂/CNC Nanocomposites. *RSC Advances*, **9**, 40565-40576. <https://doi.org/10.1039/C9RA08861J>
- [12] Jiang, F., Shen, Y., Ma, C., Zhang, X., Cao, W. and Rul, Y. (2017) Effects of TiO₂ Nanoparticles on Wheat (*Triticum aestivum* L.) Seedlings Cultivated under Super-Elevated and Normal CO₂ Conditions. *PLOS ONE*, **12**, e0178088. <https://doi.org/10.1371/journal.pone.0178088>
- [13] Gohari, G., Mohammadi, A., Akbari, A., Panahirad, S., Dadpour, M.R., Fotopoulos, V. and Kimura, S. (2020) Titanium Dioxide Nanoparticles (TiO₂ NPS) Promote Growth and Ameliorate Salinity Stress Effects on Essential Oil Profile and Biochemical Attributes of *Dracocephalum moldavica*. *Scientific Reports*, **10**, Article No. 912. <https://doi.org/10.1038/s41598-020-57794-1>
- [14] Song, U., Shin, M., Lee, G., Roh, J., Kim, Y. and Lee, E.J. (2013) Functional Analysis of TiO₂ Nanoparticle Toxicity in Three Plant Species. *Biological Trace Element Research*, **155**, 93-103. <https://doi.org/10.1007/s12011-013-9765-x>
- [15] Andersen, C.P., King, G., Plocher, M., Storm, M., Pokhrel, L.R., Johnson, M.B. and Rygielwicz, P.T. (2016) Germination and Early Plant Development of Ten Plant Species Exposed to Titanium Dioxide and Cerium Oxide Nanoparticles. *Environmental Technology and Chemistry*, **35**, 2223-2229. <https://doi.org/10.1002/etc.3374>
- [16] Debnath, K., Das, A., Das, B. and Karfoma, J. (2020) TiO₂ Nanoparticles Enhancing Germination, Growth and Yield of Rice. *International Research Journal of Pure and Applied Chemistry*, **21**, 25-30. <https://doi.org/10.9734/irjpac/2020/v21i630173>
- [17] Sebesta, M., Kolencik, M., Sunil, B.R., Illa, R., Mosnacek, J., Ingle, A. and Urik, M. (2021) Field Application of ZnO and TiO₂ Nanoparticles on Agricultural Plants. *Agronomy*, **11**, Article No. 2281. <https://doi.org/10.3390/agronomy11112281>
- [18] Asli, S. and Neuman, P.M. (2009) Colloidal Suspensions of Clay or Titanium Dioxide Nanoparticles Can Inhibit Leaf Growth and Transpiration via Physical Effects on Root Water Transport. *Plant, Cell and Environment*, **32**, 577-584. <https://doi.org/10.1111/j.1365-3040.2009.01952.x>
- [19] Ruffini Castiglione, M., Giorgetti, L., Ger., C. and Cremonini, R. (2010) The Effects of Nano-TiO₂ on Seed Germination, Development and Mitosis of Root Tip Cells of *Vicia narbonensis* and *Zea mays* L. *Journal of Nanoparticle Research*, **13**, 2443-2449. <https://doi.org/10.1007/s11051-010-0135-8>
- [20] Du, W., Sun, Y., Ji, R., Wu, J. and Gao, H. (2011) TiO₂ and ZnO Nanoparticles Negatively Affect Wheat Growth and Soil Enzyme Activities in Agricultural Soil. *Journal of Environmental Monitoring*, **13**, 822-828. <https://doi.org/10.1039/c0em00611d>

- [21] Larue, C., Khodja, H., Herlin-Biome, N., Brisset, F., Flank, A., Fayard, B., Chillou, S. and Carriere, M. (2011) Investigation of Titanium Dioxide Nanoparticles Toxicity and Uptake by Plants. *Journal of Physics*, **304**, Article ID: 012057. <https://doi.org/10.1088/1742-6596/304/1/012057>
- [22] Korenkova, L., Sebesta, M., Urik, M., Kolencik, M., Kratosova, G., Bujdos, M., Vavra, I. and Dobrocka, E. (2017) Physiological Response of Culture Media-Grown barley (*Hordeum vulgare* L.) to Titanium Dioxide Nanoparticles. *Acta Agriculturae Scandinavica, Section B—Soli and Plant Science*, **67**, 285-291. <https://doi.org/10.1080/09064710.2016.1267255>
- [23] Li, J., Naeem, M.S., Wang, X., Liu, L., Chen, C., Ma, N. and Zang, C. (2015) Nano-TiO₂ Is Not Phytotoxic as Revealed by the Oilseed Rape Growth and Photosynthetic Apparatus Ultrastructural Response. *PLOS ONE*, **10**, e0143885. <https://doi.org/10.1371/journal.pone.0143885>
- [24] Larue, C., Layurette, J., Herlin-Biome, N., Khodja, H., Fayard, B., Flank, A., Brisset, F. and Carriere, M. (2012) Accumulation, Translocation and Impact of TiO₂ Nanoparticles in Wheat (*Triticum aestivum* ssp.): Influence of Diameter and Crystal phase. *Science of the Total Environment*, **431**, 197-208. <https://doi.org/10.1016/j.scitotenv.2012.04.073>
- [25] Hu, J., Wu, X., Wu, F., Chen, W., Zhang, X., White, J.C., Li, J., Wan, Y., Liu, J. and Wang, X. (2020) TiO₂ Nanoparticle Exposure on Lettuce (*Lactuca sativa* L.): Dose-Dependent Deterioration of Nutritional Quality. *Environmental Science Nano*, **7**, 501. <https://doi.org/10.1039/C9EN01215J>
- [26] Mattiello, A., Filippi, A., Poscic, F., Musetti, R., Savatici, M.C., Giordano, S., Vischi, M., Bertolini, A. and Marchiol, L. (2015) Evidence of Phytotoxicity and Genotoxicity in *Hordeum vulgare* L. exposed to CeO₂ and TiO₂ Nanoparticles. *Frontiers in Plant Science*, **6**, Article No. 1043. <https://doi.org/10.3389/fpls.2015.01043>
- [27] Dogaroglu, Z.G. and Koleli, N. (2017) TiO₂ and ZnO Nanoparticles Toxicity in Barley (*Hordeum vulgare* L.). *Clean-Soil Air Water*, **45**, Article ID: 1700096. <https://doi.org/10.1002/clen.201700096>
- [28] Jurkow, R., Kalisz, A., Huska, D., Sekara, A. and Dastborhan, S. (2021) Sequential Changes in Antioxidant Potential of Oakleaf Lettuce Seedlings Caused by Nano-TiO₂ Treatment. *Nanomaterials*, **11**, Article No. 1171. <https://doi.org/10.3390/nano11051171>
- [29] Zheng, L., Hong, F., Lu, S. and Liu, C. (2005) Effect of Nano-TiO₂ on Strength of Naturally Aged Seeds and Growth of Spinach. *Biological Trace Element Research*, **104**, 83-91. <https://doi.org/10.1385/BTER:104:1:083>
- [30] Lin, D. and Xing, B. (2007) Phytotoxicity of Nanoparticles: Inhibition of Seed Germination and Root Growth. *Environmental Pollution*, **150**, 243-250. <https://doi.org/10.1016/j.envpol.2007.01.016>
- [31] Chen, Y., Wu, N., Mao, H., Zhou, J., Su, Y., Zhang, Z., Zhang, H. and Yuan, S. (2019) Different Toxicities on Nanoscale Titanium Dioxide Particles in the Roots and Leaves of Wheat Seedlings. *RSC Advances*, **9**, 19243-19252. <https://pubs.rsc.org/en/content/articlehtml/2019/ra/c9ra02984b> <https://doi.org/10.1039/C9RA02984B>
- [32] Wani, S., Irum, S., Gul, I., Yaqoob, K., Khalid, U., Ali, M., Manzoor, U., Noor, T., Ali, S., Rizwan, M. and Arshad, M. (2021) TiO₂ Nanoparticles Dose, Application Method and Phosphorous Levels Influence Genotoxicity in Rice (*Oryza sativa* L.), Soil Enzymatic Activities and Plant Growth. *Ecotoxicology and Environmental Safety*, **213**, Article ID: 111977. <https://doi.org/10.1016/j.ecoenv.2021.111977>

- [33] Lima, A., Carvalho, A., Pinheiro, S., Torres, Y., Miguel, T., Pireda, S., Fechine, P., Fergolente, L. and Miguel, E. (2023) Effect of TiO₂ Microparticles in Lettuce (*Lactuca sativa* L.) Seeds and Seedlings. *Bulletin of Environmental Contamination and Toxicology*, **110**, Article No. 116. <https://doi.org/10.1007/s00128-023-03752-2>
- [34] Song, U., Jun, H., Waldman, B., Roh, J., Kim, Y., Yi, J. and Lee, E. (2013) Functional Analyses of Nanoparticle Toxicity: A Comparative Study of the Effects of TiO₂ and Ag on Tomatoes (*Lycopersicon esculentum*). *Ecotoxicology and Environmental Safety*, **93**, 60-67. <https://doi.org/10.1016/j.ecoenv.2013.03.033>
- [35] Song, G., Gao, Y., Wu, H., Hou, W., Zhang, C. and Ma. (2012) Physiological Effect of Anatase TiO₂ Nanoparticles on *Lemna minor*. *Environmental Toxicology and Chemistry*, **31**, 2147-2152. <https://doi.org/10.1002/etc.1933>
- [36] Shull, T.E., Kurepa, J. and Smalle, J.A. (2019) Anatase TiO₂ Nanoparticles Induce Autophagy and Chloroplast Degradation in Thale Cress (*Arabidopsis thaliana*). *Environmental Science and Technology*, **53**, 9522-9532. <https://doi.org/10.1021/acs.est.9b01648>