



Prevalence and Forms of Intimate Partner Violence among People Living with HIV/AIDS in Sokoto State, Nigeria

**Oche Mansur Oche ^{a,b*}, Zainab Abdullahi Kontagora ^b,
Zainab Ahunna Ezenwoko ^b,
Abdulaziz Muhammad Danmadami ^b, Zainab Umaru ^a,
Isah Hudu Garba ^a and Adenrele Temitope Elijah ^a**

^a Department of Community Health, Usmanu Danfodiyo University, Sokoto, Nigeria.

^b Department of Community Medicine, Usmanu Danfodiyo University Teaching Hospital Sokoto, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. Author OMO was involved in conceptualization of the study, study design, write up. Author ZAK was involved in data entry and analysis. Author ZAE was involved in data entry and analysis and write up. Author AMD was involved in supervision of data collection, writing of draft paper. Author ZU was involved in data collection and entry. Author IHG was involved in data collection and entry. Author ATE was involved in data collection and entry. All the authors reviewed the final paper and approved same. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JAMMR/2024/v36i45406

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/114191>

Original Research Article

Received: 08/01/2024

Accepted: 12/03/2024

Published: 19/03/2024

ABSTRACT

Introduction: Intimate partner violence (IPV) is an aberrant behavior that occurs within an intimate relationship resulting in physical, sexual, and psychological harm to a partner. It transcends all strata of the society including socio-cultural groups, families and the community as a whole. Amongst people diagnosed with the Human Immuno-deficiency Virus (HIV), IPV could have

*Corresponding author: E-mail: ochedr@hotmail.com; oche.manusr@udusok.edu.ng;

adverse health consequences and this could have serious implications for disease progression, transmission and control. This study was therefore aimed at assessing the knowledge, prevalence and forms of IPV against people with HIV/AIDS in Sokoto metropolis, Nigeria.

Methods: A cross-sectional descriptive study design was used to recruit a total of 330 respondents which comprised adult male and female HIV/AIDS patients receiving comprehensive health care services in selected health facilities in Sokoto metropolis. Data on knowledge, prevalence and forms of IPV was elicited using a set of semi-structured interviewer-administered questionnaire downloaded into Open Data Kit (ODK). Data collected was entered into and analyzed using SPSS statistical software version 25 with level of statistical significance set at $p < 0.05$.

Results: More than two-thirds, 259 (78.5%) of the respondents were diagnosed with HIV within the last one to ten years with almost all, 323(97.9%) having good knowledge of IPV. Only 28 (8.5%) of the respondents experienced at least one form of IPV with more females (9.5%) compared to 4.5% of males experiencing IPV. The various forms of IPV experienced by the respondents included sexual (4.2%), physical (6.7%) and psychological (8.5%).

Conclusion: Although almost all the respondents were knowledgeable about IPV, only a few of them experienced one form of IPV or the other. This has wider public health implications for interventions and underscores the need to initiate IPV screening among people living with HIV, improved awareness of IPV and supportive in care services and counseling aimed at better clinical outcomes amongst PLWHA.

Keywords: Intimate partner violence; HIV; prevalence; PLWHA; Sokoto; public health; sexual harm; health consequences; immediate; intimate partner.

1. INTRODUCTION

Intimate Partner Violence (IPV) is often defined as any behavior within an intimate relationship that causes physical, psychological, or sexual harm to those within the relationship [1]. Globally, intimate partner violence (IPV) is a serious public health problem with a global prevalence of 30% for physical and/or sexual IPV among ever-partnered women [2]. Estimates by authors across different regions in Nigeria have reported prevalence of IPV ranging from 29% in South west [3], 41% in South south [4], 42% in the North [5] to 78.8% in Southeast [6].

Intimate partner violence (IPV) is associated with immediate and lifelong adverse health consequences, including impairment of multiple organ systems, physical injuries, permanent disabilities, and death [7,8]. Studies from South Africa and Rwanda have indicated that physical, sexual and psychological IPV were associated with higher levels of subsequent risk of HIV [9,10,11-15]. In sub-Saharan Africa, the prevalence of IPV among HIV-positive women ranged from 26 to 72% [16,17] It is to be noted that Intimate partner violence and HIV are overlapping/intersecting challenges, with a significantly high prevalence among women who are living with HIV/AIDS [18-20].

Intimate partner violence in all its forms considerably increases the risk of human

immunodeficiency virus (HIV) infection, especially in communities where the traditional patriarchal system operates, and violence against women perpetrated by an intimate partner is endorsed [21,22-29]. Significant evidence also indicated that violence against women may increase the risk of HIV transmission both directly and indirectly [30,31-33]. A systematic review has revealed that several persons living with HIV have experienced different forms of IPV [34,35-40] with coerced or forced sexual initiation having a significant contribution to a woman's risk for HIV infection [41]. Previous studies have observed that after the disclosure of HIV serostatus, one in three women experienced partner violence while some women experienced controlling behaviour by their partner, including emotional abuse, denial of communication, blame, abandonment, refusal to use safer sex methods, withdrawal of marital support, and marriage dissolution, stigma, and violence [42,43,44,45]. The Federal Government of Nigeria Official Gazette of November 2014 forbids discrimination against People living with HIV/AIDS (PLWHA) including any form of violence against such persons [46,47]. Most of the previous studies [48-54] carried out in several countries had focused on IPV against women with little attention to both men and women. To the best of the authors' knowledge, no work has been carried out to assess IPV against PLWHA in Sokoto state hence the need for this study which was aimed at assessing the knowledge,

prevalence and forms of IPV against people with HIV/AIDS in Sokoto metropolis, Nigeria.

2. MATERIALS AND METHODS

2.1 Study Area

Sokoto (also called the seat of the caliphate) is located in North-Western Nigeria near the confluence of the Sokoto River and Rima River. The inhabitants of the area are predominantly Muslim of Hausa and Fulani ethnic groups. It has a total of 23 local government areas (LGAs). The metropolis which is the capital city of the state is home to Usmanu Danfodiyo University Teaching Hospital, State Specialist Hospital, several primary health centers and private health facilities. Several of these health facilities render comprehensive HIV/AIDS services in Sokoto among which are seven centers within the metropolis including the following: (1) Primary Health Care: Military Barracks Hospital, Sokoto, Holy Family Trinity Hospital Sokoto and CHC Kofar Rini. (2) Secondary Health Care: Women and Children Welfare Centre (WCWC), Sokoto and Maryam Abacha Women and Children Clinic (MAWCC), Sokoto. (3) Tertiary Health Care: Usmanu Danfodio University Teaching Hospital (UDUTH), Sokoto and State Specialist Hospital (SHS), Sokoto. The services rendered include consultation, HIV testing, adherence counseling, home visits and drug dispensing.

2.2 Study Design

A cross-sectional descriptive study design was used.

2.3 Study Population

The target populations for this study were adult male and female persons with HIV/AIDS residing in Sokoto metropolis. The study populations were male and female persons with HIV/AIDS receiving treatment at selected health facilities in the metropolis offering comprehensive HIV/AIDS services during the study period. Study participants were individuals with HIV/AIDS sampled from among the study population.

2.4 Inclusion Criteria

Adult male and female HIV/AIDS patients receiving comprehensive health care services in Sokoto metropolis not later than one year before the commencement of the study.

2.5 Exclusion Criteria

Clients who are critically ill and were not into any relationship were excluded.

2.6 Sample Size Determination

The minimum sample size was determined using the formula for cross-sectional studies

$$n = \frac{Z^2 pq}{d^2}$$

where;

n = minimum sample size desired;

Z = standard normal deviation at 95% confidence interval =1.96;

p = prevalence of intimate partner violence in a previous study =28%=0.28²³

q = complementary probability of p = 1 - p = 0.72

d = tolerable alpha error or level of precision =5% =0.05;

$$\text{Therefore } n = \frac{1.96^2 \times 0.28 \times 0.72}{0.0025} = 310$$

Considering that the population of adults with HIV/AIDS is less than 10,000 in Sokoto state, the formula;

$$n_f = n/1+(n)/(N)$$

Where,

n_f = Desired sample size for a population less than 10,000

n = Minimal sample size for a population greater than 10,000

N = Estimated population of adult HIV/AIDS patients in Sokoto metropolis =7,322 (as obtained from registers of facilities).

$$n_f = = 297.41 \approx 297$$

Allowing for a response rate of 90%, n = 330

Three hundred and thirty (330) eligible participants were recruited into the study

2.7 Sampling Technique

A two-stage sampling technique was carried out. A line list of the health facilities rendering comprehensive HIV/AIDS services in Sokoto metropolis was obtained [7]. The health facilities were grouped into primary, secondary and tertiary.

Stage 1:

Using simple random sampling by balloting, one facility was selected from amongst primary, secondary and tertiary Health facilities (3 in all) namely, CHC Kofar Rini Sokoto, Abacha Women and Children Clinic, ad

Usmanu Danfodiyo University Teaching Hospital, Maryam

The patients' registers in each of the selected health facilities were assessed to determine the number of patients attending each facility weekly. Thereafter proportionate to size allocation was done.

Stage 2:

Systematic sampling technique was used at each of the selected health facilities to recruit eligible participants into the study after proportional allocation based on the number of patients attending each facility every week.

2.8 Study Instruments

A semi-structured interviewer-administered questionnaire was used to collect data from study participants comprising the following sections:

- Section A: socio-demographic profile
- Section B: partner's socio-demographic profile
- Section C: HIV/AIDS History
- Section D: knowledge of IPV
- Section E: experience and forms of IPV.

The Questionnaire was developed and uploaded to the researchers' server via the Open Data Kit (ODK) application. The validated form was downloaded from the internet server into the Android phones and tablets of all the data collectors that was used to collect data on the field.

Data collected from the field was sent to the researchers' servers via the internet for aggregation and real-time monitoring of data collection.

2.9 Training of Research Assistants

A two-day training on the conduct of research, the objectives of the study, protection of personal information, interpersonal communication skills, ethics related to fieldwork, and data collection using ODK was conducted by the researchers

involving resident doctors and final-year medical students of our university. After the training, the instrument for data collection was pretested among adult HIV/AIDS patients receiving care in another health facility outside the selected ones in Sokoto metropolis. This was followed by the administration of the questionnaire to the respondents which lasted for five days.

2.10 Data Management

Exploratory data analysis using SPSS statistical software version 25 was done to identify errors in the data entry and determine the distribution of the data, this involved running descriptive statistics for all the variables. Quantitative variables were summarized using mean and standard deviation while categorical variables were summarized using frequencies and percentages.

A chi-square test was done to test associations between independent categorical variables and the prevalence of IPV among the participants. The level of statistical significance (α) for the analyses was set at $P < 0.05$.

2.11 Ethical Consideration

Ethical approval to carry out the research was obtained from the Health Research Ethics Committee of the Usmanu Danfodiyo University Teaching Hospital. Permission was sought from various heads of the health facilities while written informed consent was obtained from individual participants. The respondents were assured of strict confidentiality of their responses and were informed that their participation was voluntary and could withdraw their participation at any stage of the study.

3. RESULTS

A total of 330 questionnaires were administered and completely filled, giving a response rate of 100 percent. The mean age of the respondents was 36.95 years \pm 9.93, with most 153 (46.4%) of them within the age group of 30-39 years. Most, 264 (80.0%) of the respondents were females, 285 (86.4%) were Muslims and 277 (83.9%) currently married. More than half 167 (60.3%) of the respondents were in monogamous family settings, and 310 (93.9%) had between 0-7 children. Less than half, 139 (42.1%) of the respondents had secondary school as their highest level of education (Table 1).

Table 2 shows the mean age of the respondents' partners was 43.23 ± 9.922 years; those within the age group of 40-49 years had the highest proportion 114 (34.5%). Most 285 (86.4%) were Muslims, less than half 145

(43.9%) had attained the tertiary level of education and 121 (36.7%) were civil servants. Only 10(3.0%) and 7(2.1%) of the respondents used alcohol and marijuana respectively.

Table 1. Socio-demographic characteristics of people with HIV/AIDS in Sokoto metropolis

Variables	Frequencies n = 330	Percentage (%)
Age in years		
<30	69	20.9
30-39	153	46.4
40-49	58	17.6
50-59	35	10.6
≥ 60	15	4.5
Sex		
Male	66	20.0
Female	264	80.0
Religion		
Islam	285	86.4
Christianity	45	13.6
Relationship Status		
Currently married	277	83.9
Divorced/Separated	25	7.6
Boyfriend/Girlfriend living together	10	3.0
Boyfriend/Girlfriend living apart	18	5.5
Family setting if married		
Monogamy	167	60.3
Polygamy	110	39.7
No. of children		
0-7	310	93.9
>7	20	6.1
No. of children alive		
0-7	317	96.1
>7	13	3.9
Tribe		
Hausa	235	71.2
Fulani	41	12.4
Igbo	21	6.4
Yoruba	27	8.2
¹ Others	6	1.8
Highest level of education		
None	36	10.9
Primary	45	13.6
Secondary	139	42.1
Tertiary	87	26.4
Quranic only	23	7.0
Occupation		
Unemployed	68	20.6
Petty trader	114	34.5
Civil servant	42	12.7
Business	80	24.2
Farmers	10	3.0
² Others	16	4.8
Which of these do you take		
Cigarette		
Yes	9	2.7
No	321	97.3
Alcohol		
Yes	2	0.6
No	328	99.4
Totulin*		
Yes	1	0.3
No	329	99.7
Marijuana		
Yes	2	0.6
No	328	99.4

¹Dakarkari, Idoma, Igala, ²Driver, Student, tailor *A cough syrup commonly used as drug of abuse

Table 2. Sociodemographic characteristics of the partners of people living with HIV/AIDS in Sokoto Metropolis

Variables	Frequencies n = 330	Percentage (%)
Age in years		
<30	18	5.5
30-39	98	29.7
40-49	114	34.5
50-59	74	22.4
60-69	24	7.3
>70	2	0.6
Religion		
Islam	285	86.4
Christianity	45	13.6
Highest level of education		
None	15	4.5
Primary	30	9.1
Secondary	130	39.4
Tertiary	145	43.9
Quranic only	10	3.0
Occupation		
Unemployed	42	12.7
Petty trader	37	11.2
Civil servant	121	36.7
Business	94	28.5
Farmers	18	5.5
*Others	18	5.5
Which of these does your partner take (multiple responses allowed)		
Cigarette		
Yes	64	19.4
No	266	80.6
Alcohol		
Yes	10	3.0
No	320	99.0
Tutolin ^a		
Yes	3	0.9
No	327	99.1
Marijuana		
Yes	7	2.1
No	323	97.9

**Carpenter, Driver, Mechanic, Student
a cough syrup used as drug of addiction*

Table 3. HIV/AIDS History of respondents living with HIV/AIDS in Sokoto metropolis

Variables	Frequencies n =330	Percentage (%)
Duration of HIV diagnosis (years)		
1-10	295	89.4
>10	35	10.6
On anti-retroviral medication		
Yes	330	100.0
No	0	0.0
Duration of treatment (years)		
1-9	267	80.9
10-19	57	17.3
≥20	6	1.8
Partner aware of his/her status		
Yes	286	86.7
No	25	7.6
Don't know	19	5.8
Use of condoms during sexual intercourse		
Yes	109	33.0
No	221	67.0
Do you know if your partner has other sexual partners		
Yes	61	18.5
No	108	32.7
Don't know	161	48.8

HIV – Human Immunodeficiency Virus

Table 4. Knowledge of Intimate Partner Violence among people with HIV/AIDS in Sokoto metropolis

Variables	Frequenciesn = 330	Percentage (%)
Have you heard of IPV?		
Yes	310	93.9
No	20	6.1
Have you witnessed IPV as a child growing up?		
Yes	66	20.0
No	264	80.0
Which of these acts constitute IPV?		
Being shoved		
Yes	317	96.1
No	13	3.9
Being slapped		
Yes	322	97.6
No	8	2.4
Being kicked		
Yes	323	97.9
No	7	2.1
Being pushed		
Yes	322	97.6
No	8	2.4
Being dragged		
Yes	323	97.9
No	7	2.1
Being beaten up		
Yes	321	97.3
No	9	2.7
Being scolded		
Yes	323	97.9
No	7	2.1
Being burnt on purpose		
Yes	7	2.1
No	323	97.9
Being threatened		
Yes	321	97.3
No	9	2.7
Using a weapon		
Yes	323	97.9
No	7	2.1
Being insulted		
Yes	320	97.0
No	10	3.0
Using abusive language		
Yes	322	97.6
No	8	2.4
Being belittled		
Yes	322	97.6
No	8	2.4
Being intimidated in front of other people		
Yes	320	97.0
No	10	3.0
Getting jealous that a partner is unfaithful when he/she relates with other women/men		
Yes	320	97.0
No	10	3.0
Getting suspicious that partner is unfaithful when he/she relates with other women/men		
Yes	321	97.3
No	9	2.7
Preventing or restricting partner from going to his/her parent's friends'/relatives' houses		
Yes	322	97.6
No	8	2.4
Physically forcing partner to have sexual intercourse when she/he did not want to		
Yes	291	88.2
No	39	11.8
Making the partner to do sexual activities they		

Variables	Frequenciesn = 330	Percentage (%)
find degrading or humiliating?		
Yes	323	97.9
No	7	2.1
Knowledge grade		
Good	323	97.9
Poor	7	2.1

IPV – Intimate Partner Violence




Fig. 1. Prevalence of Intimate Partner Violence amongst PLWHA in Sokoto metropolis




Fig. 2. Prevalence of Intimate Partner Violence by Sex amongst people with HIV/AIDS in Sokoto metropolis

In Table 3, more than two-thirds 259 (78.5%) of the respondents were diagnosed with HIV between 1-10 years ago. All 330 (100.0%) the respondents are currently on anti-retroviral medication with 267 (80.9%) having been on treatment for 1-9 years. Most 286 (86.7%) of the

respondents' partners knew their HIV status; close to one-third 109 (33.0%) used condoms during sexual intercourse and almost half 161 (48.8%) respondents did not know if their partners had other sexual partners.

Almost all 323 (97.9%) respondents had good knowledge of intimate partner violence with only 7 (2.1%) recording poor knowledge (Table 4).

Of all the respondents, only 28 (8.5%) had experienced at least one form of intimate partner violence while 302 (91.5%) had no experience (Fig. 1).

Findings from the study showed that more females (9.5%) experienced IPV more compared to males (4.5%) (Fig. 2)

3.1 Forms of Intimate Partner Violence Experienced by Respondents

The prevalence of physical violence was 6.7%. The commonest form of physical violence experienced by the respondents was being slapped by partner 18 (15.0%) followed by being pushed by partner 17 (14.2%). The least 2 (1.7%) forms of physical abuse experienced by respondents were being burnt on purpose, using sharp objects on them and being threatened with a gun by their partners (Table 5).

The prevalence of psychological violence was 8.5%. The commonest form of psychological violence experienced by the respondents was respondents feeling ignored by their partners 24 (16.4%), followed by respondents' partners monitoring their movements 17 (11.6%). The least 5 (3.4%) forms of psychological abuse experienced by respondents were that of the threat of hurting the respondent's family, children or pets (Table 6).

The prevalence of sexual violence was 4.2% with the commonest form of sexual violence experienced by the respondents being denial of sex as a punishment 10 (25.0%). The least forms of sexual abuse experienced by respondents included being forced or threatened to engage in oral sex, getting hurt or injured in respondents' private parts intentionally and being forced to watch pornographic films against respondents' will 3 (7.5%) (Table 7).

All the respondents reported various effects of IPV which included cuts and abrasions, miscarriages, abandonment, depression, and loneliness.

Table 5. Physical abuse among people with HIV/AIDS in Sokoto metropolis

*Variables	Frequency n = 28	Percentage (%)
Partner ever slapped you?	18	15.0
Partner ever pushed you?	17	14.2
Partner ever hit you with her hand?	14	11.7
Partner ever beaten you up?	14	11.7
Partner ever thrown something at you?	12	10.0
Partner ever hit you with anything that could hurt you?	10	8.3
Partner ever shoved you?	9	7.5
Partner ever dragged you?	6	5.0
Partner ever scalded you?	6	5.0
Partner ever kicked you?	5	4.2
Partner ever pulled your hair?	3	2.5
Partner ever burnt you on purpose?	2	1.7
Partner ever used a sharp object on you (e.g knife, razor, bottle)?	2	1.7
Partner ever threatened you with a gun?	2	1.7

**Multiple responses allowed*

Table 6. Psychological abuse among respondents

*Variables	Frequency n = 28	Percentage (%)
Partner ignores your feelings	24	16.4
Partner monitors your movement	17	11.6
Partner threatens to leave or make you leave	16	11.0
Partner monitors your phone calls with others	16	11.0
Partner goes through your phone without your permission	16	11.0
Partner stalks you when you are with other people	13	8.9
Partner threatens to take your children	8	5.5
Partner threatens to hurt your prized possessions	8	5.5
Partner isolates you from friends and family?	7	4.8
Partner goes through your documents without your permission?	6	4.1
Partner threatens to hurt your family?	5	3.4
Partner threatens to hurt your children?	5	3.4
Partner threatens to hurt your pets?	5	3.4

**Multiple responses allowed*

Table 7. Sexual abuse among respondents

*Variables	Frequency n = 28	Percentage (%)
Sexual abuse		
Partner ever denied you sex as a punishment?	10	25.0
Partner forced you to have sex with him/her against your will?	6	15.0
Partner used threats to make you have sex with her/him?	6	15.0
Partner made hurtful statements about your sexual performance?	6	15.0
Partner used force or threats to make you engage in oral sex?	3	7.5
Partner forced you to have sex with a condom against your judgment?	3	7.5
Partner ever hurt or caused injury to your private parts intentionally?	3	7.5
Partner forced you to watch pornographic film against your will?	3	7.5

*Multiple responses allowed

Table 8. Relationship between respondents' socio-demographic characteristics and experience of intimate partner violence

Variable	Experience of intimate partner violence		Test statistic p value	OR (95% CI)
	Yes n(%)	No n (%)		
Age group (years)				
19 – 30	7 (6.4)	102 (93.6)	$\chi^2 = 0.892$ $p = 0.345$	0.654 (0.269 – 1.588)
>30	21 (9.5)	200 (90.5)		
Sex (respondent)				
Male	3 (4.5)	63 (95.5)	$\chi^2 = 1.694$ $P = 0.199$	0.455 (0.133 – 1.556)
Female	25 (9.5)	239 (90.5)		
Relationship status				
Currently married	25 (9.0)	252 (91.0)	Fisher's exact = 1.555 $p = 0.654$	-
Divorced/separated	2 (8.0)	23 (92.0)		
Boyfriend/girlfriend living together	1 (10.0)	9 (90.0)		
Boyfriend/girlfriend living apart	0 (0.0)	18 (100.0)		
Family setting if married				
Monogamy	20 (13.2)	147 (86.8)	$\chi^2 = 4.460$ $p = 0.035$	2.875 (1.039 – 7.856)
Polygamy	5 (4.7)	105 (95.3)		
No. of children				
0-7	26 (8.4)	284 (91.6)	Fisher's exact $P = 0.682$	0.824 (0.181 – 3.749)
>7	2 (10.0)	18 (90.0)		
Educational status				
Informal	8 (13.6)	51 (86.4)	$\chi^2 = 2.383$ $P = 0.123$	1.969 (0.822 – 4.715)
Formal	20 (7.4)	251 (92.6)		
Occupation				
Unemployed	9 (11.5)	69 (88.5)	$\chi^2 = 1.141$ $p = 0.285$	1.580 (0.679 – 3.676)
Employed	18 (7.6)	218 (92.4)		
Substance abuse				
Yes	2 (16.7)	10 (83.3)	Fisher's exact $p = 0.270$	2.246 (0.467–10.799)
No	26 (8.2)	292 (91.8)		
Knowledge grade				
Good	27 (8.4)	296 (91.6)	Fisher's exact $p = 0.466$	0.547 (0.064 – 4.714)
Poor	1 (14.3)	6 (85.7)		

$\chi^2 =$ Pearson's chi square, $p = < 0.05$, OR = odds ratio, CI = Confidence Interval

The proportion of respondents 20 (13.2%), who were married and in a monogamous family setting that experienced intimate partner violence was less than those who experienced IPV 147 (86.8%) in the same family setting, and the association was statistically significant ($\chi^2 = 4.460$, $p = 0.035$). Among female respondents, the proportion of those who did not experience IPV 239 (90.5%) was higher than those who had experienced IPV 25 (9.5%), but the association was not statistically significant ($\chi^2 = 1.694$, $P = 0.199$) (Table 8).

The number of female respondents with good knowledge about IPV 261 (98.9%) was higher than the number of male respondents 62 (93.9%) and the association was statistically significant (Fisher's exact = NA, $p = 0.032$). Also, the proportion of the respondents with 0-7 children and had good knowledge of IPV 306 (98.7%) was higher than those with more than 7 children 17 (85.0%) and the association was statistically significant (Fisher's exact = NA, $p = 0.006$) (Table 9).

Table 9. Relationship between respondents' socio-demographic characteristics and knowledge of intimate partner violence

Variable	Knowledge grade of IPV		Test statistic p value	OR (95% CI)
	Good n(%)	Poor n (%)		
Age group (years)				
19 – 30	109 (100.0)	0 (0.0)	Fisher's exact = Not available p = 0.100	1.033 (1.008 – 1.058)
>30	214 (96.8)	7 (3.2)		
Sex (respondent)				
Male	62 (93.9)	4 (6.1)	Fisher's exact = Not available P = 0.032	0.178 (0.039 – 0.817)
Female	261 (98.9)	3 (1.1)		
Relationship status				
Currently married	251 (97.3)	7 (2.7)	Fisher's exact = 0.500 p = 1.000	-
Divorced/separated	22 (100.0)	0 (0.0)		
Boyfriend/girlfriend living together	10 (100.0)	0 (0.0)		
Boyfriend/girlfriend living apart	18 (100.0)	0 (0.0)		
Family setting if married				
Monogamy	147 (96.7)	5 (3.3)	Fisher's exact = Not available p = 0.704	0.565 (0.108 – 2.970)
Polygamy	104 (98.1)	2 (1.93)		
No. of children				
0-7	306 (98.7)	4 (1.3)	Fisher's exact = not available P = 0.006	13.500 (2.796 –65.189)
>7	17 (85.0)	3 (15.0)		
Educational status				
Informal	58 (98.3)	1 (1.7)	Fisher's exact = Not available P = 1.000	1.313 (0.155–11.117)
Formal	265 (97.8)	6 (2.2)		
Occupation				
Unemployed	78 (100.0)	0 (0.0)	Fisher's exact = Not available p = 0.199	1.031 (1.008 – 1.054)
Employed	229 (97.0)	7 (3.0)		
Substance intake				
Yes	12 (100.0)	0 (0.0)	Fisher's exact = Not available p = 1.000	1.023 (1.006 – 1.040)
No	311 (97.8)	7 (2.2)		

$\chi^2 =$ Pearson's chi square, $p = < 0.05$, OR = odds ratio, CI = Confidence Interval

Table 10. Relationship between respondents' HIV/AIDS history and experience of intimate partner violence

Variable	Experience of intimate partner violence		Test statistic p value	OR (95% CI)
	Yes n(%)	No n (%)		
Duration of HIV diagnosis (years)				
1-10	26 (8.8)	269 (91.2)	Fisher's exact p = 0.752	1.595 (0.362 – 7.027)
>10	2 (5.7)	33 (94.3)		
Duration on antiretroviral medication (years)				
1-10	26 (8.8)	271 (91.2)	Fisher's exact p = 1.000	1.487 (0.337 – 6.568)
>10	2 (6.1)	31 (93.9)		
Partner knows his/her HIV status?				
Yes	23 (8.0)	263 (92.0)	Fisher's exact = 1.118 P = 0.598	-
No	3 (12.0)	22 (88.0)		
Don't know	2 (10.5)	17 (89.5)		
Do you use any condoms during sexual intercourse?				
Yes	11 (10.1)	98 (89.9)	$\chi^2 = 0.541$ P = 0.462	1.347 (0.608 – 2.985)
No	17 (7.7)	204 (92.3)		
Aware if your partner has other sexual partners?				
Yes	6 (9.8)	55 (90.2)	$\chi^2 = 1.112$ p = 0.574	-
No	11 (10.2)	97 (89.8)		
Don't know	11 (6.8)	150 (93.2)		

$\chi^2 =$ Pearson's chi square, $p = < 0.05$, OR = odds ratio, CI = Confidence Interval
HIV – Human Immunodeficiency Virus

The proportion of respondents diagnosed with HIV/AIDS 269 (91.2%), within 1 – 10 years who did not experience intimate partner violence was higher than those who experienced IPV 26 (8.8%), however, the association was not statistically significant (Fisher's exact, $p = 0.752$) (Table 10).

4. DISCUSSION

It has been observed that the general perception of IPV used to be that of a female victim and a male perpetrator. However, this perception is gradually being adjusted to expose the increasing trend of a female perpetrator and a male victim, or a male-male or female-female victim and perpetrator with varying magnitudes, forms, and motivations [55,56]. Nevertheless, it is important to mention that there are reports that suggest that the motivations for perpetrating IPV by men and women are similar [57,58].

In this study, the magnitude of IPV among the respondents was low (8.4%) and this is a reflection of the socio-cultural milieu of the study area where acts of IPV are seen as normal within a relationship. In a similar study where investigators used data from the Medical Monitoring Project (MMP), an annual cross-sectional survey used to produce nationally representative estimates of sociodemographic, behavioral, and clinical characteristics of adult PWH in the U.S.A, it was observed that only 4.4% of their respondents experienced IPV in the past 12 months [59]. A study amongst the general population in China indicated that the prevalence of IPV among people living with HIV was 15.44%, a result lower than that of Canada (35%) [60]. In contrast to these findings, a similar study from Enugu, Southeast Nigeria recorded the magnitude of IPV among their respondents to be 52.5% in the last 12 months before the study [56]. Similarly, a study in rural US noted that 39% of their study subjects experienced IPV in the past one year while 66.7% experienced more than one form of IPV [61]. Also, findings from a study in Uganda, a country with a high burden of HIV observed that 65.29 and 72.22% of men and women with HIV respectively reported that they had experienced at least one form of IPV in their lifetime [17]. The difference in the magnitude between our study and others could be attributed to cultural differences as cases of IPV amongst partners should not be made public hence the low reportage of IPV in our society. In the different cultures of Nigeria, women are made to believe that IPV is a practice that has been passed from one generation to another therefore

the women must come to terms with it and the men must not report it otherwise it will be seen as a sign of weakness and lack of machoism.

Before now the general perception of IPV used to be that of a female victim and a male perpetrator. However, this perception is gradually being adjusted to expose the increasing trend of a female perpetrator and a male victim, or a male-male or female-female victim and perpetrator [62].

In this study the prevalence of IPV was found to be statistically higher among females (9.5%) than males (4.5%) living with HIV. This finding was not surprising, as IPV is a problem commonly occurring among women [61], and research has shown that gender is a prominent risk factor for IPV, with women being disproportionately affected when compared to men [62].

Low prevalence rates of IPV have been reported amongst women in Ogun state Nigeria, 5.02% [63], and Jos 12.6% [64]. Similar studies from Zaria, and Kano in northern Nigeria reported 22 and 28% among their study subjects respectively [65,50].

The national figure reported among the general populace during the Nigeria Demographic and Health Survey (NDHS) was 28% while it was 30% for women in the general population and these figures are by far greater than figures obtained in this study [66]. Furthermore, a higher prevalence of IPV was obtained from other studies in Nigeria (65.8%) [67], South Africa 67.3% [68], DRC Congo 51% [69], and Columbia Canada 59% [70].

The much higher figures reported in studies from other parts of Nigeria and Africa compared to our own figure may be due to socio-cultural differences and specific instruments used to collect data as studies that utilized the WHO questionnaire with behaviorally specific questions on acts of IPV, have been found to improve disclosure rates [71].

Men are usually more educated, often engaged in paid employment, and therefore economically more viable than women and this invariably explains the inequality in power hence it is expected that IPV should always be perpetrated by the men. However this is not always the case and in the absence of any visible compromise by men, women are now known to perpetrate violence against men as seen in this study where 4.5% of the men were known to have

experienced violence. The prevalence of IPV among men is low when compared to similar studies in rural Appalachia USA 39% [60], Enugu Nigeria 39% [56], Uganda 69.3% [17], and China 16.3% [72]. The high prevalence observed in other studies may not be unrelated to good reporting of IPV in those societies compared to our study area where abuse of men must not be heard as such trends if in public domain will demean men and see them as weak.

There is a huge body of literature that shows the various forms of IPV experienced across the globe. In this study, the commonest form of abuse experienced by our respondents was psychological (8.5%) followed by sexual and physical abuse. This is in consonance with findings from other studies [17,61,73,72].

It has been observed in developing countries that a woman is the property of the husband who handles her the way he likes, and there is a failure of authorities to treat sexual violence as a criminal offence, hence discouraging reporting on sexual violence by most women [73,74]. Although these abuses occurred in different locations, however, the forms and magnitude vary considerably. As reported in this study, the effects of IPV ranged from physical to emotional problems including cuts and abrasions, loneliness, abandonment, miscarriages by the women and depression. The study conducted in Enugu state Nigeria observed that their respondents suffered the same problems that includes both physical to mental health problems, and sometimes suicidal ideations which might have been heightened by the double burden of experiencing IPV while living with HIV [56]. Findings from this study indicated that being in a monogamous marriage was the only factor associated with the risk of IPV amongst our respondents. In contrast to this finding, the study from Uganda showed that the sociodemographic status of being married was associated with a higher risk of IPV [74], while another type of marital status, such as being divorced, was also associated with IPV in another context [75,76]. This calls for concerted efforts including improving public awareness and providing family counselling aimed at stemming the tide of IPV regardless of marital status and marriage settings.

5. CONCLUSION

The overall prevalence of IPV among People living with HIV in Sokoto is not high, however, IPV is disproportionately higher among women

compared to men. This underscores the need to initiate IPV screening among people living with HIV, improved awareness of IPV and supportive in care services and counseling aimed at better clinical outcomes amongst PLWHA.

6. LIMITATIONS OF THE STUDY

This study explored personal experiences of IPV and it was difficult extracting these information from the respondents for obvious reasons especially amongst the men as they would not want to be seen as weaklings in the hands of their female partners. Recall bias played out during interview sessions

CONSENT AND ETHICAL APPROVAL

Ethical approval for this study was obtained from the Health Research Ethics Committee of Usmanu Danfodiyo University Teaching Hospital Sokoto and written informed consent obtained from participants before the commencement of data collection.

CONSENT FOR PUBLICATION

All the authors went through the final version of the manuscript and have consented to the paper being published

DATA AVAILABILITY

Data for this research can be made available on reasonable request.

ACKNOWLEDGEMENT

We wish to acknowledge the Resident doctors who assisted in the data collection and also thank the participants who spared their precious time to respond to our questions

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. World Health Organization. World report on violence and health. Krug EG, Dahlberg LL, Mercy JA, Zwi AB, Lozano R, editors. Geneva: World Health Organization; 2002 [Accessed 18 Feb 2023].

- Available from: http://apps.who.int/iris/bitstream/handle/10665/42495/9241545615_eng.pdf?sequence=1].
2. World Health Organization. Global and regional estimates of violence against women: prevalence and health effects of intimate partner violence and non-partner sexual violence. WHO, 2013. Available: <https://www.who.int/reproductivehealth/publications/violence/9789241564625/en/>. Accessed 19 October, 2023.
 3. Benebo FO, Schumann B, Vaezghasemi M. Intimate partner violence against women in Nigeria: a multilevel study investigating the effect of women's status and community norms. *BMC Women's Health*. 2018;18(1):136
 4. Dienye P, Gbeneol P, Gbeneol IK. Intimate partner violence and associated coping strategies among women in a primary care clinic in Port Harcourt. *Niger J Fam Med Prim Care*. 2014;3(3):193-198
 5. Tanimu TS, Yohanna S, Omeiza SY. The pattern and correlates of intimate partner violence among women in Kano, Nigeria. *African J Prim Heal Care Fam Med*. 2016;8(1):e1-e6.
 6. Okemgbo CN, Omideyi AK, Odimegwu OC. Prevalence, patterns and correlates of domestic violence in selected Igbo communities of Imo state, Nigeria. *Afr J Reprod Heal*. 2002;6(2):101-114
 7. Smith SG, Chen J, Basile KC et al. The National Intimate Partner and Sexual Violence Survey (NISVS): state report. Atlanta, GA: National Center for Injury Prevention and Control, Centers for Disease Control and Prevention; 2010–2012. Available: <https://www.cdc.gov/violenceprevention/pdf/nisvs-staterreportbook.pdf>. Published April 2017. Accessed June, 25, 2021
 8. Black MC. Intimate partner violence and adverse health consequences: implications for clinicians. *Am J Lifestyle Med*. 2011;5(5):428–439. DOI:10.1177/1559827611410265
 9. Jewkes RK, Dunkle K, Nduna M, Shai N: Intimate partner violence, relationship power inequity, and incidence of HIV infection in young women in South Africa: a cohort study. *Lancet*. 2010, 376: 41-48. DOI:10.1016/S0140-6736(10)60548-X.
 10. Dude AM: Spousal intimate partner violence is associated with HIV and Other STIs among married Rwandan women. *AIDS Behav*. 2011, 15: 142-152. DOI:10.1007/s10461-009-9526-1.
 11. Kouyoumdjian FG, Findlay N, Schwandt M, Calzavara LM. A systematic review of the relationships between intimate partner violence and HIV/ AIDS. *PLoS ONE*. 2013;8(11): e81044
 12. Stockman JK, Lucea MB, Campbell JC. Forced sexual initiation, sexual intimate partner violence and HIV risk in women: a global review of the literature. *AIDS Behav*. 2013;17(3):832–4
 13. Colombini M, James C, Ndwiga C, Mayhew SH. The risks of partner violence following HIV status disclosure, and health service responses: narratives of women attending reproductive health services in Kenya. *J Int AIDS Soc*. 2016;19(1):20766.
 14. Maeri I, El Ayadi A, Getahun M, Charlebois E, Akatukwasa C, Tumwebaze D et al. "How can I tell?" Consequences of HIV status disclosure among couples in eastern African communities in the context of an ongoing HIV "test-and-treat" trial. *AIDS Care*. 2016;28(Suppl 3):59–66
 15. HIV and AIDS (Anti-Discrimination) Act, The Federal Republic of Nigeria Official Gazette. The Federal Government Printer, Lagos, Nigeria; 2014. FGP43/521015/1,200
 16. Kouyoumdjian FG, Findlay N, Schwandt M, Calzavara LM. A systematic review of the relationships between intimate partner violence and HIV/ AIDS. *PLoS ONE*. 2013;8(11): e81044.
 17. Ogbonnaya IN, Wanyenze RK, Reed E, Silverman JG, Kiene SM. Prevalence of and risk factors for intimate partner violence in the first 6 months following HIV diagnosis among a population-based sample in rural Uganda. *AIDS Behav*. 2020;24(4):1252–65
 18. Shi CF, Kouyoumdjian FG, Dushof J. Intimate partner violence is associated with HIV infection in women in Kenya: a cross-sectional analysis. *BMC Public Health*. 2013; 13:512.
 19. Durevall D, Lindskog A. Intimate partner violence and HIV in ten sub-Saharan African countries: What do the Demographic and Health Surveys tell us? *Lancet Glob Health*. 2015;3(1):e34–43
 20. Grant C., Lynn R. Addressing the intersection of HIV/AIDS, violence against women and girls, & gender-related health disparities: Interagency Federal Working

- Group Report. The White House Washington; 2013, September. Accessed 19 Oct. 2023
Available:<https://obamawhitehouse.archives.gov/blog/2013/09/06-addressing-intersection-hiv-aids-violence-against-women-and-girls-and-gender-related->
21. Kabwama SN, Bukenya J, Matovu JK, Gwokyalya V, Makumbi F, BeyezaKashesya J et al. Intimate partner violence among HIV positive women in care-results from a national survey, Uganda 2016. *BMC Womens Health*. 2019;19(1):1–10
 22. World Health Organization. World report on violence and health. Krug EG, Dahlberg LL, Mercy JA, Zwi AB, Lozano R, editors. Geneva: World Health Organization; 2002. [Accessed 18 Feb 2023].
Available:http://apps.who.int/iris/bitstream/handle/10665/42495/9241545615_eng.pdf?sequence=
 23. World Health Organization. Global and regional estimates of violence against women: prevalence and health effects of intimate partner violence and non-partner sexual violence. WHO, 2013. Accessed 19 October,2023
Available:<https://www.who.int/reproductivehealth/publications/violence/9789241564625/en/>
 24. Tanimu TS, Yohanna S, Omeiza SY. The pattern and correlates of intimate partner violence among women in Kano, Nigeria. *African J Prim Heal Care Fam Med*. 2016;8(1):e1-e6.
 25. Benebo FO, Schumann B, Vaezghasemi M. Intimate partner violence against women in Nigeria: a multilevel study investigating the effect of women’s status and community norms. *BMC Women's Health*. 2018;18(1):136
 26. Okemgbo CN, Omidayi AK, Odimegwu OC. Prevalence, patterns and correlates of domestic violence in selected Igbo communities of Imo state, Nigeria. *Afr J Reprod Heal*. 2002;6(2):101-114
 27. Dienne P, Gbeneol P, Gbeneol IK. Intimate partner violence and associated coping strategies among women in a primary care clinic in Port Harcourt. *Niger J Fam Med Prim Care*. 2014;3(3):193-198
 28. Smith SG, Chen J, Basile KC, et al. The National Intimate Partner and Sexual Violence Survey (NISVS): state report. Atlanta, GA: National Center for Injury Prevention and Control, Centers for Disease Control and Prevention; 2010–2012.
Available:<https://www.cdc.gov/violenceprevention/pdf/nisvs-staterreportbook.pdf>.
Published April 2017. Accessed June, 25, 2021
 29. Black MC. Intimate partner violence and adverse health consequences: Implications for clinicians. *Am J Lifestyle Med*. 2011;5(5):428–439.
DOI:10.1177/1559827611410265
 30. Jewkes R, Flood M, Lang J. From work with men and boys to changes of social norms and reduction of inequities in gender relations: a conceptual shift in prevention of violence against women and girls. *The Lancet*. 2015;385(9977):1580–9
 31. Olowookere SA FO, Adekanle DA, Adeleke NA, Abioye-Kuteyi EA. Patterns and Correlates of Intimate Partner Violence to Women Living With HIV/AIDS in Osogbo, Southwest Nigeria. *Violence Against Women*. 2015;21(11):1330–40.
 32. Shri N, Muhammad T. Association of intimate partner violence and other risk factors with HIV infection among married women in India: Evidence from National Family Health Survey 2015–16. *BMC Public Health*. 2021;21(1):2105.
Available:<https://doi.org/10.1186/s12889-021-12100-0>.
 33. Colombini M, James C, Ndwiga C, Team I, Mayhew SH. The risks of partner violence following HIV status disclosure, and health service responses: Narratives of women attending reproductive health services in Kenya. *Journal of the International AIDS Society*. 2016;19(1).
Available:<https://doi.org/10.7448/IAS.19.1.20766>.
 34. Kouyoumdjian FG, Findlay N, Schwandt M, Calzavara LM. A systematic review of the relationships between intimate partner violence and HIV/ AIDS. *PLoS ONE*. 2013;8(11): e81044
 35. Jewkes RK, Dunkle K, Nduna M, Shai N: Intimate partner violence, relationship power inequity, and incidence of HIV infection in young women in South Africa: a cohort study. *Lancet*. 2010, 376: 41-48.
DOI:10.1016/S0140-6736(10)60548-X.
 36. Dude AM: Spousal intimate partner violence is associated with HIV and Other STIs among married Rwandan women. *AIDS Behav*. 2011, 15: 142-152.
10.1007/s10461-009-9526-1.

37. Kouyoumdjian FG, Findlay N, Schwandt M, Calzavara LM. A systematic review of the relationships between intimate partner violence and HIV/ AIDS. PLoS ONE. 2013;8(11): e81044.
38. Ogonnaya IN, Wanyenze RK, Reed E, Silverman JG, Kiene SM. Prevalence of and risk factors for intimate partner violence in the first 6 months following HIV diagnosis among a population-based sample in rural Uganda. AIDS Behav. 2020;24(4):1252–65
39. Shi CF, Kouyoumdjian FG, Dushof J. Intimate partner violence is associated with HIV infection in women in Kenya: a cross-sectional analysis. BMC Public Health. 2013;13:512.
40. Durevall D, Lindskog A. Intimate partner violence and HIV in ten sub-Saharan African countries: what do the Demographic and Health Surveys tell us? Lancet Glob Health. 2015;3(1):e34–43
41. Grant C., Lynn R. Addressing the intersection of HIV/AIDS, violence against women and girls, & gender-related health disparities: interagency Federal Working Group Report. The White House Washington; 2013, September. Accessed 19 Oct. 2023
Available: <https://obamawhitehouse.archives.gov/blog/2013/09/06-addressing-intersection-hiv-aids-violence-against-women-and-girls-and-gender-related>.
42. Stockman JK, Lucea MB, Campbell JC. Forced sexual initiation, sexual intimate partner violence and HIV risk in women: a global review of the literature. AIDS Behav. 2013;17(3):832–4
43. Colombini M, James C, Ndwiga C, Mayhew SH. The risks of partner violence following HIV status disclosure, and health service responses: narratives of women attending reproductive health services in Kenya. J Int AIDS Soc. 2016;19(1):20766.
44. Maeri I, El Ayadi A, Getahun M, Charlebois E, Akatukwasa C, Tumwebaze D et al. "How can I tell?" Consequences of HIV status disclosure among couples in eastern African communities in the context of an ongoing HIV "test-and-treat" trial. AIDS Care. 2016;28(Suppl 3):59–66
45. Kabwama SN, Bukenya J, Matovu JK, Gwokyalya V, Makumbi F, Beyezakashesya J et al. Intimate partner violence among HIV positive women in care-results from a national survey, Uganda 2016. BMC Womens Health. 2019;19(1):1–10
46. Jewkes R, Flood M, Lang J. From work with men and boys to changes of social norms and reduction of inequities in gender relations: a conceptual shift in prevention of violence against women and girls. The Lancet. 2015;385(9977):1580–9
47. HIV and AIDS (Anti-Discrimination) Act, The Federal Republic of Nigeria Official Gazette. The Federal Government Printer, Lagos, Nigeria; 2014. FGP43/521015/1,200
48. Olowookere SA FO, Adekanle DA, Adeleke NA, Abioye-Kuteyi EA. Patterns and Correlates of Intimate Partner Violence to Women Living With HIV/AIDS in Osogbo, Southwest Nigeria. Violence Against Women. 2015;21(11):1330–40.
49. Federal Ministry of Health. Technical report: 2010 National HIV sero-prevalence sentinels survey. Abuja, Nigeria: Department of Public Health, National AIDS/STI Control Programme; 2010. Available: http://www.nigeriaaids.org/documents/2010_National%20HIV%20Sero%20Prevalence%20Sentinel%20Survey.pdf
50. Iliyasu Z, Abubakar IS, Babashani M, Galadanci HS. Domestic violence among women living with HIV/AIDS in Kano, Northern Nigeria. African Journal of Reproductive Health. 2011;15:41-49
51. National Population Commission (NPC) [Nigeria] and ICF Macro. Nigeria demographic and health survey 2008. Abuja, Nigeria: National Population Commission and ICF Macro; 2009.
52. Silverman JG, Decker MR, Saggurti N, Balaiah D, Raj A. Intimate partner violence and HIV infection among married Indian women. Journal of the American Medical Association. 2008;300:703-710.
53. McCree DH, Koenig LJ, Basile KC, Fowler D, Green Y. Addressing the intersection of HIV and intimate partner violence among women with or at risk for HIV in the United States. J Womens Health (Larchmt). 2015;24(5):331–335. [PubMed: 25973798]
DOI:10.1089/jwh.2015.5301.
54. Pyra M, Weber K, Wilson TE, et al. Sexual minority status and violence among HIV infected and at-risk women. J Gen Intern Med. 2014;29(8):1131–1138.
DOI:10.1007/s11606-014-2832-y. [PubMed: 24700180]

55. Zierler S, Cunningham WE, Andersen R et al. Violence victimization after HIV infection in a U.S. probability sample of adult patients in primary care [published correction appears in Am J Public Health. 2000;90(3):447]. Am J Public Health. 2000;90(2):208–215. [PubMed: 10667181]. DOI:10.2105/ajph.90.2.208.
56. Obionu IM, Echefu CO, Chinweokwu VU, Chineme WO, Gabriel AU, Aguwa EN. Intimate partner violence among HIV-positive service users of a tertiary care and treatment facility in Enugu State, Nigeria. HIV AIDS Rev 2021;20(4):275-280. DOI:https://doi.org/10.5114/hivar.2021.111896
57. Nybergh L, Taft C, Enander V, Krantz G. Self-reported exposure to intimate partner violence among women and men in Sweden: Results from a population-based survey. BMC Public Health. 2013;13(1):845. Available:https://doi.org/10.1186/1471-2458-13-845.
58. Elmquist J, Hamel J, Shorey RC, Labrecque L, Ninnemann A, Stuart GL. Motivations for intimate partner violence in men and women arrested for domestic violence and court referred to batterer intervention programs. Partner Abuse. 2014;5(4):359–374. Available:https://doi.org/10.1891/1946-6560.5.4.359].
59. Ansley B, Lemons-Lyn, Amy R. Baugher, Sharoda Dasgupta, Jennifer L. Fagan, Sharon G. Smith, R. Luke Shouse. Intimate Partner Violence Experienced by Adults with Diagnosed HIV in the U.S. Am J Prev Med. 2021 June;60(6):747–756. DOI:10.1016/j.amepre.2020.12.019).
60. Siemieniuk RA, Krentz HB, Gish JA, Gill MJ. Domestic violence screening: prevalence and outcomes in a Canadian HIV population. AIDS Patient Care STDS 2010;24:763–70;
61. Bryan N, Davidov DM, Dick T, Bassler J, Fisher M. Intimate partner violence experiences among men living with HIV in rural Appalachia. AIDS and Behav 2019; 23: 3002–14
62. Howe TR. Marriages & Families in the 21st Century: A Bioecological Approach. Wiley-Blackwell; 2012;44. Available:https://books.google.com.ng/books?hl=en&lr=&id=tcRJ4VpjE2MC&oi=fnd&pg=PR15&dq=Howe+TR.+Families+in+crisis:+violence,+abuse,+and+neglect:+intimat
- e+partner+violence:+marital+rape.+In+Howe+TR.+Marriages+and+families+in+the+21st+century,+a+bioecological+appro (Accessed: 20.12.2023)
63. Abiodun O, Sodeinde K, Bamidele F, Ojinni Y, Adekeye J, Ohiaogu F, Mbonu F. Intimate Partner Violence Among Women Accessing HIV-Care Services at Tertiary Hospitals in Ogun State, Nigeria: Implications for Policy and Practice. J Interpers Violence. 2022 Jan;37(1-2):58-78. Epub 2020 Feb 28. PMID: 32108537.
64. Gyuse AMI, Ushie AP. Pattern of domestic violence among pregnant women in Jos, Nigeria. South African Family Practice 2009; 51(4):56-59
65. Ameh N, Abdul MA. Prevalence of domestic violence among pregnant women in Zaria, Nigeria. Annals of African Medicine 2004;2(1):4-6.
66. National Population Commission (NpopC) [Nigeria] and ORC Macro. Nigeria Demographic and Health Survey 2008 Calverton, Maryland: National Population Commission and ORC Macro.2004:169-172
67. Ezechi OC, Gab-Okafor C, Onwujekwe DI, et al. Intimate partner violence and correlates in pregnant HIV positive Nigerians. Arch Gynecol Obstet 2009; 280:745–52
68. Brown LL, Van Zyl MAR. Mitigating intimate partner violence among South African women testing HIV positive during mobile counseling and testing. AIDS Care 2018;30:65–71
69. Gichane MW, Moracco KE, Thirumurthy H, Okitolonda W, Behets F, Yotebieng M. Intimate partner violence and prevention of mother to child transmission of HIV: evidence from Kinshasa, Democratic Repub Congo. 2018;1–11
70. Closson K, McLinden T, Parry R, Lee M, Gibbs A, Kibel M, Wang L, Trigg J, Braitstein P, Pick N, Parashar S, Montaner JSG, Kaida A, Hogg RS. Severe intimate partner violence is associated with all-cause mortality among women living with HIV. AIDS. 2020 Aug 1;34(10):1549-1558. DOI:10.1097/QAD.0000000000002581. PMID: 32675565
71. World Health Organization. WHO multi-country study on women's health and domestic violence against women: Initial results on prevalence, health outcomes and women's responses: World Health Organization; 2005.

72. Wu Y, Wang Y, Long H, Liu L, Dai L, Cao W, Liu J, Liu M. Prevalence of Intimate Partner Violence and Associated Factors Among People With HIV: A Large-Sample Cross-Sectional Study in China. *Journal of Infectious Diseases* 2023;228:1592-1599.
73. Li Y, Marshall CM, Rees HC, Nunez A, Ezeanolue EE, Ehiri JE. Intimate partner violence and HIV infection among women: a systematic review and meta-analysis. *J Int AIDS Soc* 2014;17:18-45.
74. Wolff B, Blanc AK, Gage AJ. Who decides? Women's status and negotiation of sex in Uganda. *Cult Health Sex.* 2000;2:303–22.
75. Moher D, Liberati A, Tetzlaff J, et al. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med.* 2009;6: e1000097.
76. Shamu S, Zarowsky C, Shefer T, et al. Intimate partner violence after disclosure of HIV test results among pregnant women in Harare, Zimbabwe. *PLoS One.* 2014; 9: e109447.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/114191>