



Efficacy of Azoxystrobin 7.5% and Propiconazole 12.5% SE Against Sheath Blight in Rice

S. B. Gowdar ^{a*} and Narappa G ^b

^a Department of Plant Pathology, College of Agriculture, Gangavathi, Koppal - 583 227 (Karnataka), India.

^b ICAR-Krishi Vigyan Kendra, Gangavathi, Koppal - 583 227 (Karnataka), India.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JEAI/2024/v46i52375

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/114542>

Original Research Article

Received: 11/01/2024

Accepted: 15/03/2024

Published: 19/03/2024

ABSTRACT

Among the fungal diseases causing significant yield loss in rice, sheath blight caused by *Rhizoctonia solani* Kuhn is the most important ranked the second most important disease worldwide after blast and a serious threat in rice growing areas of the world and causes more economic yield losses. The field experiment was conducted with seven treatments and replicated three times in RBD design at ARS, Gangavathi to know the effect of Azoxystrobin 7.5% + Propiconazole 12.5% SE against sheath blight in rice. The variety BPT-5204 was sown in plot size of 5 X 5 m² with all regular agronomic practices followed as per the standard package of practice of University of Agricultural Sciences, Raichur. It has been found that the fungicide Azoxystrobin 7.5% + Propiconazole 12.5% SE at different doses evaluated was effective in reducing the severity of rice sheath blight and thereby increased the rice grain yield. The treatment Azoxystrobin 7.5% + Propiconazole 12.5% SE @ 625 ml/ha recorded minimum sheath blight of 13.70 and 16.11 per

++ Associate Professor;

*Corresponding author: E-mail: sbgowdar@gmail.com;

cent PDI with yield of 64.60 q/ha and 63.20 q/ha compared to control treatment 48.70 and 57.96 per cent PDI with grain yield of 54.10 q/ha and 52.40 q/ha during the first and second season, respectively. In the current study, along with reducing sheath blight severity, fungicide treatments minimized grain yield losses in 2 years. Field trial conducted clearly indicated that Azoxystrobin 7.5% + Propiconazole 12.5% SE @ 500 and 625 ml/ha dose can effectively control sheath blight of rice. Azoxystrobin 7.5% + Propiconazole 12.5% w/v SE @ 500 ml/ha was at par with higher dose and resulted better yield than other treatments.

Keywords: Azoxystrobin 7.5% + propiconazole 12.5% SE; fungicide; rice; sheath blight.

1. INTRODUCTION

“There are about 50 different biotic factors that can cause potential yield loss in rice including fungi, bacteria, viruses, nematodes and insects. Of the disease-causing organisms, fungal pathogens impose a greater challenge in sustaining rice production” [1]. “Among the fungal diseases causing significant yield loss in rice, sheath blight is the most important ranked the second most important rice disease worldwide after blast” [2,3,4]. “Sheath blight of rice caused by *Rhizoctonia solani* Kuhn is an important disease of rice and a serious threat in rice growing areas of the world and causes more economic yield losses” [5,6,7], ranging from 20 to 50 per cent depending on the severity of infection [8,9] and also 5-10 per cent yield loss in subtropical low land paddy cultivars of Asia [5]. “The yield losses ranging from 4 to 50 per cent have been reported depending on the crop stage at the time of infection, severity of the disease and environmental conditions” [10,11,12]. The potential losses due to sheath blight disease are reported to be 50 to 54.3 per cent alone in India [13,14] and this disease is particularly most prevalent in intensive rice cultivation system due to excess use of nitrogenous fertilizers.

“In the absence of effective host plant resistance against sheath blight pathogen in rice, the management of sheath blight disease is mainly carried out through the use of chemicals” [15]. “Fungicide based management of sheath blight disease is successful at field level in majority of the cases” [16,17,18,19,20]. “Foliar spray and seed treatment are the most popular method of fungicidal application against *R. solani*. Even though both systemic and non-systemic fungicides are used for chemical management, systemic fungicides offer better management of this disease” [15]. “Timely application of selective fungicides between panicle differentiation and heading stage offers effective protection against this disease. Periodical monitoring of the rice field and application of fungicides at the initial

stages of infection especially at booting stage is recommended for managing sheath blight in susceptible varieties” [21,22]. “Most of the fungicides like benomyl, carbendazim, chloroneb, captafol, mancozeb, zineb, edifenphos, lprobenphos, thiophanate, carboxin etc. have been found effective for the control of the disease under field conditions” [23,24,25]. “Out of these benomyl, carbendazim, edifenphos and lprobenphos were the most effective chemicals” [13]. “For many years, strobilurin fungicides have been the backbone for management of rice sheath blight. It has been opined that *R. solani*, would not develop fungicide resistance or would be slow to develop resistance” [26]. “Presently, the Strobilurin group of systemic fungicides are the most preferred chemical group to manage sheath blight disease in rice” [27]. “Azoxystrobin from this group is very effective for not only controlling the disease but also found to enhance yield as well” [8]. The combinatory chemical formulation such as Azoxystrobin 18.2% + Difenoconazole 11.4% [19,28], Trifloxystrobin 25% + Tebuconazole 50% [29,30], etc., are recommended to manage the disease. Recently, many combination fungicides such as kresoxim methyl 40% + hexaconazole 8%, azoxystrobin 18.2% + difenoconazole 11.4% SC, trifloxystrobin 25% + tebuconazole 50% 75 WG, have been shown to control the sheath blight disease under field condition [31,32]. It is vital to look for a novel molecule with a distinct mode of action because the constant use of fungicides belonging to the same group and having the same mode of action would cause the same fungi to evolve resistant strains. In order to determine the field effectiveness of a combination fungicide, Azoxystrobin 7.5% + Propiconazole 12.5% SE, against sheath blight in rice, the current study was conducted.

2. MATERIALS AND METHODS

The field experiment was conducted with seven treatments and replicated three times in RBD design at ARS, Gangavati to know the effect of Azoxystrobin 7.5% + Propiconazole 12.5% SE

against sheath blight in Rice. The variety BPT-5204 was sown in plot size of 5 X 5 m² with all regular agronomic practices followed as per the standard package of practice of University of Agricultural Sciences, Raichur. The evaluation of the fungicide was done along with standard checks and untreated control against the incidences of sheath blight disease in rice. Treatments details were as follows - T1 - Azoxystrobin 7.5% + Propiconazole 12.5% SE @ 375 ml / ha; T2 - Azoxystrobin 7.5% + Propiconazole 12.5% SE @ 500ml / ha; T3 - Azoxystrobin 7.5% + Propiconazole 12.5% SE @ 625 ml / ha; T4 - Azoxystrobin 23 % SC @ 500 ml / ha; T5 - Propiconazole 25% EC @ 500 ml / ha; T6 – Validamycin 3% L @ 2000 ml / ha; T7 - Untreated control.

“The fungicides were applied as foliar spray treatment in the replicated plots just after the appearance of sheath blight disease in the main field at 45DAT. The plots were inspected regularly to see the disease development and further two more spray were applied at an interval of 15 days on 60 and 75 Days after Transplanting (DAT). Observations were recorded on disease severity in each treatment after three sprays as per the standard method. The incidence of disease were recorded on leaves as Per cent Disease Index (PDI) on the basis of scoring of the diseases as per the disease rating scale of Standard Evaluation System (SES) for rice” [33]. In the present study, observations for disease incidence were recorded from the randomly selected twenty clumps / hills per plot for recording the disease severity in each replicated plots of the treatments. The observations were recorded on intensity of diseases were observed in each replicated plot for each treatment on 10th day after each spray. After 10 days of last spray, the final scoring of the disease incidence was recorded. Further, the scored data were converted into Per cent Disease Index (PDI) of plants using formula given by Wheeler [34].

$$PDI = \frac{\text{Sum of numerical values}}{\text{Number of leaves observed}} \times \frac{100}{\text{Maximum disease rating value}}$$

In order to record the yield, crop was harvested plot-wise from the individual replicated plots and average paddy yield was recorded and converted into q/ha.

The original PDI values were suitably transformed into arcsine transformed values and subjected to statistical analysis for drawing conclusions

3. RESULTS AND DISCUSSION

3.1 Sheath Blight Infection

It has been found that Azoxystrobin 7.5% + Propiconazole 12.5% SE @ 500- 625 ml/ha reduced the sheath blight infection more than rest of the treatments and improved the rice yield. The Azoxystrobin 7.5% + Propiconazole 12.5% SE was found to be effective in reducing the severity of the sheath blight disease. In the 1st season, the treatment Azoxystrobin 7.5% + Propiconazole 12.5% SE @ 625 ml/ha recorded least PDI of sheath blight disease incidence (13.70%) and was significantly superior over control treatment (48.70%). The same treatment was significantly on par with Azoxystrobin 7.5% + Propiconazole 12.5% SE @ 500 ml/ha treatment with PDI of 14.07 per cent. All other treatments were inferior to these two treatments though significantly superior to the untreated check. Among the various treatments, Azoxystrobin 23% SC @ 500ml/ha was found least effective with higher PDI of sheath blight (26.11%) (Table-1). During the 2nd season, the Azoxystrobin 7.5% + Propiconazole 12.5% SE was found to be effective in reducing the severity of the sheath blight disease. The treatment Azoxystrobin 7.5% + Propiconazole 12.5% SE @ 625 ml/ha recorded least PDI of sheath blight disease incidence (16.11%) and was significantly superior over control treatment (57.96%). The same treatment was significantly on par with Azoxystrobin 7.5% + Propiconazole 12.5% SE @ 500 ml/ha treatment with PDI of 16.48 per cent. All other treatments were inferior to these two treatments though significantly superior to the untreated check (57.96%). Among the various treatments, Azoxystrobin 23% SC @ 500ml/ha was found least effective with higher sheath blight (30.56%) (Table 2). These data were in accordance with Phelp and Soto [35] and Jones et al. [36]. These finding are in full agreement with the previous reports of fungicide combinations such as azoxystrobin + difenconazole [19], kresoxim methyl + hexaconazole, fluxapyroxad + epixiconazole [31], trifloxystrobin + tebuconazole [37,32,38], flutolanil + azoxystrobin, thiophonate methyl + azoxystrobin, tebuconazole + azoxystrobin, propiconazole + azoxystrobin [39] reported the better efficacy against sheath blight. Various reviews confirmed that strobilurin compounds found to be effective in controlling many diseases like sheath blight [40,32].

Table 1. Effect of Azoxystrobin 7.5% +Propiconazole 12.5% SE on sheath blight of rice during Kharif - 2011 (1st season)

Sl. No.	Treatment	Dose a.i. (g/ha)	Dose Formulation (ml/ha)	Sheath blight intensity on rice				Grain Yield (q/ha)
				Initial score	10 days after I spray	10 days after II spray	Terminal score (10 days after III spray)	
1	Azoxystrobin 7.5% + Propiconazole 12.5% SE	28.1+46.9	375	4.63 (12.43)*	12.78 (20.94)	17.22 (24.52)	25.19 (30.12)	59.40
2	Azoxystrobin 7.5% + Propiconazole 12.5% SE	37.5+62.5	500	4.07 (11.64)	6.67 (14.96)	12.41 (20.62)	14.07 (22.03)	63.50
3	Azoxystrobin 7.5% + Propiconazole 12.5% SE	46.9+78.1	625	4.44 (12.17)	6.11 (14.31)	11.85 (20.14)	13.70 (21.73)	64.60
4	Azoxystrobin 23% SC	125	500	4.26 (11.91)	13.15 (21.26)	20.37 (26.83)	26.11 (30.73)	59.10
5	Propiconazole 25% EC	125	500	4.07 (11.64)	8.96 (17.42)	14.15 (22.10)	18.37 (25.38)	61.00
6	Validamycin 3% L	60	2000	4.26 (11.91)	10.56 (18.96)	16.30 (23.81)	20.74 (27.09)	60.70
7	Control	-	-	4.81 (12.68)	26.30 (30.85)	33.52 (35.38)	48.70 (44.26)	54.10
CD at 5% level				NS	2.40	1.38	3.30	2.48

* Figures in the parentheses represent arcsine transformed values

Table 2. Effect of Azoxystrobin 7.5% +Propiconazole 12.5% SE on sheath blight of rice during Kharif - 2012 (2nd season)

Sl. No.	Treatment	Dose a.i. (g/ha)	Dose Formulation (ml/ha)	Sheath blight intensity on rice				Grain Yield (q/ha)
				Initial score	10 days after I spray	10 days after II spray	Terminal score (10 days after III spray)	
1	Azoxystrobin 7.5% + Propiconazole 12.5% SE	28.1+46.9	375	5.19 (13.16)*	15.37(23.08)	19.81 (26.43)	29.07 (32.63)	57.50
2	Azoxystrobin 7.5% + Propiconazole 12.5% SE	37.5+62.5	500	5.37 (13.40)	9.63(18.08)	15.93(23.52)	16.48 (23.66)	62.20
3	Azoxystrobin 7.5% + Propiconazole 12.5% SE	46.9+78.1	625	5.00(12.92)	8.70(17.16)	15.19(22.93)	16.11 (23.66)	63.20
4	Azoxystrobin 23% SC	125	500	5.56(13.63)	18.52(25.49)	25.37(30.24)	30.56 (33.56)	57.60
5	Propiconazole 25% EC	125	500	5.37(13.40)	11.93(20.21)	18.85(25.73)	22.37 (28.23)	59.00
6	Validamycin 3% L	60	2000	5.74(13.86)	14.81(22.64)	20.00(26.57)	24.81 (29.88)	58.70
7	Control	-	-	5.74(13.86)	29.81(33.10)	34.44(35.94)	57.96 (49.58)	52.40
CD at 5% level				NS	2.12	2.18	3.24	3.00

* Figures in the parentheses represent arcsine transformed values

3.2 Yield

Due to varying degree of reduction in sheath blight disease incidences the difference in the yield level between treated and untreated plots was very much significant. During the 1st season, the maximum grain yield of 64.60 q/ha was recorded with Azoxystrobin 7.5% + Propiconazole 12.5% SE @ 625 ml/ha followed by the same fungicide @ 500ml/ha which recorded 63.50 q/ha was statistically superior and comparatively effective than rest of the treatments. The treatments Propiconazole 25% EC, Validamycin 3% L, and Azoxystrobin 23% SC recorded the yield of 61.00, 60.70 and 59.40 q/ha, respectively. The lowest grain yield (54.10 q/ha) was recorded in untreated control (Table 1). During the 2nd season, the maximum grain yield of 63.20 q/ha was recorded with Azoxystrobin 7.5% + Propiconazole 12.5% w/v SE @ 625 ml/ha followed by the same fungicide @ 500ml/ha which recorded 62.20 q/ha was statistically superior and comparatively effective than rest of the treatments tested. The treatments Propiconazole 25% EC, Validamycin 3% L and Azoxystrobin 23% SC recorded the yield of 59.00, 58.70 and 57.60 q/ha, respectively. The lowest grain yield was recorded with 52.40 q/ha in untreated control (Table 2). In the current study, along with reducing sheath blight severity, fungicide treatments minimized grain yield losses in 2 years. The results are in agreement with previous studies that reported the efficacy of azoxystrobin [3], azoxystrobin and flutolanil [8] against sheath blight and in minimizing yield losses. Application of fungicides has been reported to enhance the crop yield due to reduction in disease load [41,40]. The present results are in conformity with those of previous reports Bhuvaneshwari and Raju [19], Bag et al. [42] reported that fungicides application increases the yield of rice, mainly due to reduced disease severity of sheath blight.

4. CONCLUSION

The fungicide Azoxystrobin 7.5% + Propiconazole 12.5% SE at different doses were evaluated at Agriculture Research Station, Gangavati and was found effective in reducing the severity of rice sheath blight and thereby increased the rice grain yield. Field trial conducted clearly indicated that Azoxystrobin 7.5% + Propiconazole 12.5% SE @ 500 and 625 ml/ha dose can effectively control sheath blight of rice. Azoxystrobin 7.5% +

Propiconazole 12.5% w/v SE @ 500 ml/ha was at par with higher dose and resulted better yield than other treatments.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Webster RK, Gunnell PS. Compendium of Rice Diseases. St Paul, MN: American Phytopathological Society, viii–62; 1992.
2. Pan X, Zou J, Chen Z, Lu J, Yu H, Li H. Tagging major quantitative trait loci for sheath blight resistance in a rice variety, Jasmine 85. Chin. Sci. Bull. 1999;44:1783–1789.
3. Groth DE. Azoxystrobin rate and timing effects on rice sheath blight incidence and severity and rice grain and milling yields. Plant Dis. 2005;89:1171–1174.
4. Zhou XG, Jo YK. Disease management. In: Way MO, McCauley GM, Zhou XG, Wilson LT, Brandy M, (Eds.), Texas Rice Production Guidelines. Texas Rice Research Foundation. 2014;44–57.
5. Ou SH. Rice Diseases, 2nd Edn. Commonwealth Mycological Institute, Surrey; 1985.
6. Savary S, Willocquet L, Elazegui FA, Castilla N, Teng PS. Rice pest constraints in tropical Asia: Quantification and yield loss due to rice pests in a range of production situations. Plant Dis. 2000;84:357-369.
7. Savary S, Teng PS, Willocquet L, Nutter FW Jr. Quantification and modeling of crop losses: A review of purposes. Annual Rev. Phytopathol. 2006;44:89-112.
8. Groth DE, Bond JA. Effect of cultivars and fungicides on rice sheath blight, yield, and quality. Plant Dis. 2007;91:1647–1650.
9. Margani R, Widadi S. Utilizing Bacillus to inhibit the growth and infection by sheath blight pathogen, Rhizoctonia solani in rice. IOP Conf. Ser. Earth Environ. Sci. 2018; 142:012070.
10. Singh SK, Shukla V, Singh H, Sinha AP. Current status and impact of sheath blight in rice (*Oryza sativa* L.) - A review. Agric. Rev. 2004;25(4):289-297.
11. Zheng A, Lin R, Zhang D, Qin P, Xu L, Ai P, Ding L, Wang Y, Chen Y, Liu Y. The evolution and pathogenic mechanisms of

- the rice sheath blight pathogen. Nat. Commun. 2013;4:1424.
12. Bhunkal N, Singh Ramm, Mehta N. Assessment of losses and identification of slow blighting genotypes against sheath blight of rice. J. Mycol. Pl. Pathol. 2015;45:285-292.
 13. Roy AK. Sheath blight of rice in India. Indian Phytopath. 1993;46:97-205.
 14. Rajan CPD. Estimation of yield losses due to sheath blight of rice. Indian Phytopathol. 1987;40:174-177.
 15. Naik RG, Jayalakshmi K, Naik TB. Efficacy of fungicides on the management of sheath blight of rice. Int. J. Curr. Microbiol. Appl. Sci. 2017;6:611-614.
 16. Kandhari J, Gupta RL, Kandari J. Efficacy of fungicides and resistance inducing chemicals against sheath blight of rice. J. Mycological Res. 2003;41
 17. Kandhari J, Gupta RL. Efficacy of fungicides and resistance inducing chemicals against sheath blight of rice. Journal of Mycological Research. 2003; 41:67-69
 18. Groth DE, Bond JA. Initiation of rice sheath blight epidemics and effect of application timing of azoxystrobin on disease incidence, severity, yield, and milling quality. Plant Disease. 2006;90:1073-1076.
 19. Bhuvaneswari V, Raju KS. Efficacy of new combination fungicide against rice sheath blight caused by *Rhizoctonia solani* (Kühn). J. Rice Res. 2012;5:57-61.
 20. Kumar PMK, Gowda SDK, Rishikant M, Kumar KN, Gowda PKT, Vishwanath K. Impact of fungicides on rice production in India. In: Fungicides – Showcases of integrated plant disease management from around the world. 2013;77-98.
 21. Singh R, Sunder S, Kumar P. Sheath blight of rice: Current status and perspectives. Indian Phytopathol. 2016;69:340-351.
 22. Uppala S, Zhou XG. Field efficacy of fungicides for management of sheath blight and narrow brown leaf spot of rice. Crop Prot. 2018;104:72-77.
 23. Dash SC, Panda S. Chemical control of rice sheath blight disease. Indian Phytopath. 1984;37: 79-82.
 24. Kannaiyan S, Prasad NN. Effect of foliar spray of certain fungicides on the control of sheath blight of rice. Madras Agric. J. 1984;71:111-114.
 25. Singh R, Sinha AP. Comparative efficacy of local bioagents, commercial bioformulation and fungicide for the management of sheath blight of rice, under glass house conditions. Indian Phytopath. 2004;57:494-496.
 26. Robinson E. Resistant sheath blight changing practices in Louisiana; 2013. Available:<http://deltafarmpress.com/rice/resistant-sheath-blight-changing-practices-louisiana>, Accessed date: 24 July 2015.
 27. Yellareddygar SKR, Reddy MS, Kloepper JW, Lawrence KS, Fadamiro H. Rice sheath blight: A review of disease and pathogen management approaches. J. Plant Pathol. Microbiol. 2014;5:1000241.
 28. Kumar P, Ahlawat S, Chauhan R, Kumar A, Singh R, Kumar A. *In vitro* and field efficacy of fungicides against sheath blight of rice and post-harvest fungicide residue in soil, husk, and brown rice using gas chromatography-tandem mass spectrometry. Environ. Monit. Assess. 2018;190:503.
 29. Shahid AA, Shahbaz M, I Ali M. A comparative study of the commercially available fungicides to control sheath blight of rice in Lahore. J. Plant Pathol. Microb. 2014;5:2157-7471.
 30. Rashid MM, Bhuiyan MR, Dilzahan HA, Hamid MA, Hasan N, Khan MAI. Biological control of rice sheath blight disease (*Rhizoctonia solani*) using bio-pesticides and bio-control agents. Bangladesh Rice J. 2020;24:47-58.
 31. Prasanna Kumar MK, Veerabhadraswamy AL. Appraise a combination of fungicides against blast and sheath blight diseases of paddy (*Oryzae sativa* L). J. Exp. Biol. Agri. Sci. 2014;2:49-57.
 32. Bag MK. Efficacy of a new fungicide 'Trifloxystrobin 25% + Tebuconazole 50%' 75WG against Sheath Blight (*Rhizoctonia solani* Kühn) of Rice. Journal of Crop and Weed. 2009;5(1):224-226.
 33. International Rice Research Institute. Standard Evaluation System for Rice, 5th edn., Nov. 2002, INGER, Genetic Resources Center, IRRI, P.O. Box 933, 1099, Manila, Philippines. 2002;56.
 34. Wheeler BEJ. An introduction to plant diseases. John Wiley and Sons Ltd; London, UK. 1969;254.
 35. Phelp RH, Soto A. Rice diseases at CARONI and approaches to the control In Proc. Monitoring tour and workshop on Integrated Pest Management in rice in Caribbean. held in Guyana and Trinidad and Tobago. 7-11 October 1991; 1993.
 36. Jones RE, Belmar SB, Jeger MJ. Evaluation of benomyl and propiconazole

- for controlling sheath blight of rice caused by *Rhizoctonia solani*. Plant Disease. 1987;71:222-229
37. Bag MK, Saha S. Fungitoxic effect of Nativo 75WG (trifloxystrobin 25% + tebuconazole 50%) on grain discolouration (GD) disease of rice in West Bengal. Pestology. 2009;33:47-49.
38. Visalakshmi V, Raju MRB, Rao AU, Kumar KM, Satyanarayana NH. Compatibility and efficacy of insecticide and fungicide combinations on major pests and sheath blight of paddy. Nat. Env. Poll. Tech. 2016;15:233-235.
39. Jin C, Wu R, Hu H, Jifeng W, Xiaoxia W. Control effect of 6 types of azoxystrobin mixture on rice sheath blight. Chin. Agric. Sci. Bull.; 2013. Available:http://en.cnki.com.cn/Article_en/CJFDTOTAL-ZNTB201318031.htm, Accessed date: 22 April 2016.
40. Bag MK. Performance of a new generation fungicide Metominostrobin 20SC against sheath blight disease of rice. J. Mycopathol. Res. 2011;49(1):167-169.
41. Biswas A, Bag MK. Strobilurins in management of sheath blight disease of rice: A review. Pestol. 2010;34:23-26.
42. Bag MK, Yadav M, Mukherjee AK. Bioefficacy of Strobilurin based fungicides against rice sheath blight disease. Transcriptomics. 2016;4:128.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/114542>