



Comparative Studies of Lepidopteran Pest Incidence on JL 24 and DH 256 Varieties of Groundnut

Kolli Bharghavi ^{a++*}, Rohini Sugandi ^{b#},
R. Channakeshava ^{c†}, Basavaraj. S. Yenagi ^{d‡}
and Burjikindi Madhuri ^{a‡}

^a Department of Entomology, Professor Jayashankar Telangana State Agricultural University, Hyd-500030 (Telangana), India.

^b Institute of Organic Farming, University of Agricultural Sciences, Dharwad – 580005 (Karnataka), India.

^c AICRP on Soybean, Main Agricultural Research Station, University of Agricultural Sciences, Dharwad – 580005 (Karnataka), India.

^d AICRP on Groundnut, Main Agricultural Research Station, University of Agricultural Sciences, Dharwad – 580005 (Karnataka), India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.56557/UPJOZ/2024/v45i94016

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://prh.mbimph.com/review-history/3400>

Original Research Article

Received: 01/02/2024

Accepted: 05/04/2024

Published: 11/04/2024

ABSTRACT

This experiment compares the incidence of lepidopteran pests under protected and unprotected groundnut cultivation at AICRP on Groundnut, MARS, Dharwad, Karnataka in the kharif season of 2021–2022. In order to protect the protected plot from pests, protection measures were

⁺⁺ M. Sc (Agril. Entomology);

[#] Scientist (Entomology);

[†] Scientist (Agronomy);

[‡] Ph.D. Scholar;

*Corresponding author: Email: bhargavireddykolli@gmail.com;

implemented at 50 DAS using flubendiamide 20 WG @ 0.5g/l for leaf-eating caterpillars. Based on the crop's phenology, observations were made. *Spodoptera litura*, *Thysanoplusia orichalcea*, *Helicoverpa armigera*, and *Maruca vitrata*, were deemed to be the main pests, with the remaining pests being of less significance based on their population density and type of damage. When comparing unprotected JL-24 and DH-256 crop plots to protected plots, the majority of major and minor pests were infested during the vegetative to maturity stage, with maximum infestation occurring during the pod formation and pod filling stages of the crop. In comparison to unprotected plots, protected plots had the highest pod and haulm yields.

Keywords: Groundnut; lepidopteran pests; protected plot; unprotected plot.

1. INTRODUCTION

Groundnut (*Arachis hypogaea* L.) is a valuable cash crop for millions of small-scale farmers in the semi-arid tropical regions of South America. One of the most significant oil seed crops farmed in India, it provides around 30% of the country's total oil supply. Farmers in India with limited resources primarily cultivate groundnuts as a crop under rainfed systems [1]. India grows groundnuts on an area of around 55.71 lakh hectares during the kharif, rabi, and summer seasons. In 2020–21, the crop produced 102 lakh tonnes and had a productivity of 1831 kg/ha [2]. Abiotic and biotic stresses experienced by the crop during growth are the cause of low groundnut productivity. Diseases and pests are the main biotic stresses on groundnut production. There are more than 100 insects known to feed on groundnuts in India, some of which can be highly harmful and significantly lower yields viz., 26 to 100% by *S. litura* and *H. armigera* [3], 25-100% by *A. albistriga* [4]. Among them, red hairy caterpillar *Amsacta albistriga* (Walker), tobacco caterpillar *Spodoptera litura* (Fabricius), gram caterpillar *Helicoverpa armigera* (Hübner), semilooper *Thysanoplusia orichalcea* (Fabricius), leaf miner *Aproaerema modicella* (Deventer) are considered as an important destructive insect pests [5]. Studies reveal that 15 - 20 percent of the total oilseed production is lost directly or indirectly by the attack of insect and mite pests every year [6]. The population of pests and their natural enemies on groundnuts may have shifted recently as a result of varying climatic conditions [7]. Therefore, a thorough understanding of the behaviour of insect pests and their natural enemies on groundnuts will aid in predicting any outbreaks and alerting those who need to be warned. The status and order of pest appearance during the crop period, crop losses and types of damages, and detailed information about a pest complex are all crucial for developing an economically viable, ecologically sound, and socially acceptable pest

management strategy [8]. Field research was done in this regard to determine the prevalence of pests on cultivars of groundnuts that are susceptible and moderately resistant varieties.

2. MATERIALS AND METHODS

The experiment was conducted in the field at AICRP on Groundnut, Main Agricultural Research Station, Dharwad, Karnataka during *kharif* season of 2021-22. Field studies were carried out to determine the comparative incidence of major insect pests on two different groundnut cultivars JL-24 (susceptible) and Dh-256 (moderately resistant). Both varieties were sown over a plot size of 10 m x 10 m at a spacing of 30 cm x 10 cm. Two separate protected and unprotected JL-24 and Dh-256 plots (10m x 10m) (specify the plot) were maintained throughout the study period. In unprotected plots recommended package of practices was followed to raise the crop except crop protection measures in order to avoid influence of chemicals on insect population. Whereas in protected plots, chemical protection (at 50 DAS with flubendiamide 20 WG @ 0.5g/l for leaf eating caterpillars) was given on a need basis. Observations on different species of insects were recorded from 10 randomly selected plants. Chronological appearance [seedling (10-25 days), vegetative (25-35 days), flower initiation (35-45 days), pegging and pod formation (45-75 days), pod filling (75-90 days), maturity (90-110 days) and harvesting (110-120 days) stage] of the insect pests, their nature and quantity of damage, feeding behaviour and yield reductions were documented. Observations were recorded visually on the standing crop at weekly intervals.

Metarhizium rileyi: *M. rileyi* is the producer of airborne conidia on conidiophores, which are infectious propagules that start the pathogenesis. Conidia stick to the surfaces of their hosts, germinate, and create a germ tube that pierces the host's cuticle, colonizes the insect's

hemocoel, and ultimately kills the insect. The fungus's mycelium was visible on the surface of the larval integument, and as sporulation began, the fungus's white growth changed to an olive-green hue [9].

Sl. NPV: The NPV infection caused the larvae to become inactive, enlarge and become slightly glossy, change the colour of their cuticles to a darker brown, and eventually die. The larvae's lysed cuticle, which becomes frail and secretes a clear liquid that can be either pink or brownish milky white depending on the infection's severity, were the other obvious symptoms. Larva ascending to a higher altitude prior to demise (Tree Top Disease) [10].

3. RESULTS AND DISCUSSION

3.1 Insect Pests and Natural Enemies Observed on Groundnut

Twelve species of insect pests belonging to 3 orders and 9 families of pests were found to infest the groundnut at AICRP on Groundnut, Main Agricultural Research Station, Dharwad during *kharif* season 2021-22. Among these, defoliators *viz.*, *Spodoptera litura*, *Thysanoplusia orichalcea*, *Helicoverpa armigera*, *Maruca vitrata*, *Aproaerema modicella* and *Spilarctia obliqua*; Natural enemies like entomopathogens were recorded in both protected and unprotected plot of both the varieties of groundnut.

3.2 Unprotected Plot

3.2.1 Leaf eating caterpillars

In both the varieties larval populations (*S. litura*, *T. orichalcea*, *H. armigera*, *Maruca vitrata* and *A. modicella*) peaked at the time of pegging and pod formation in unprotected plots of JL-24 and Dh-256 (42.25 and 30.00 /10 plants, 23.9 and 18.00/10 plants, 17.43 and 12.31 /10 plants, 19.91 and 17.60 /10 plants and 17.04 and 7.20 /10 plants) (Table 1). Highest larval population of *S. obliqua* was found in the JL-24 plot (between 5.00 and 35.75 /10 plants) as compared with Dh-256 (2.50 to 29.21 /10 plants).

3.2.2 Natural enemies

Metarhizium rileyi and *Spodoptera litura* Nuclear polyhydrosis virus (*Sl. NPV*) populations from flowering to pod filling of JL-24 and Dh-256 (5.40 to 42.50 and 2.00 to 24.50; 2.04 to 13.20 and 1.00 to 7.00 cadavers/10 plants) (Table 2).

3.3 Protected Plot

3.3.1 Leaf eating caterpillars

In JL-24 and Dh-256 varieties the larval populations of *S. litura* range from 7.95 to 24.68 and 2.15 to 18.50/10 plants from vegetative to maturity stage. Highest population of *T. orichalcea* and *H. armigera* recorded at flower initiation of JL-24 (19.00 and 12.05/10 plants) and Dh-256 (15.29 and 9.75/10 plants) (Table 1). Population of *M. vitrata* from vegetative to maturity stage (4.57 to 16.00 and 1.58 and 11.00/10 plants) of both varieties. *A. modicella* vegetative to pod filling stage of JL-24 (1.00 to 8.80/10 plants) and pegging and pod formation in Dh-256 (5.48 and 2.00/10 plants). Highest larval population of *S. obliqua* was recorded during harvesting stage of JL-24 plot (25.50/10 plants) Dh-256 (19.51 /10 plants).

3.3.2 Natural enemies

M. rileyi and *Sl. NPV* populations from flowering and pegging and pod formation of JL-24 (10.51 and 6.08; 8.91 and 1.89 cadavers/10 plant) and Dh-256 (8.24 and 5.00; 4.82 and 0.75 cadavers/10 plants) were observed in Table 2.

The above results shows that all the sucking pests and defoliators significantly differed from unprotected plot and protected plots of both the varieties and the highest populations were recorded in JL-24 than Dh-256. This supports the findings of Pal et al., [11] that the Dh- 256 is tolerant to defoliators *viz.*, *S. litura* and leaf miner. JL-24 is susceptible to defoliators and sucking pests with confirmation of research conducted on biochemical and biophysical characters like phenols, wax, trichome density was lower as well as higher sugars in JL-24, which the harbouring of pests of groundnut [12]. In protected plots recorded lowest population because of using insecticides for controlling leaf eating caterpillars *i.e.*, flubendiamide 20 WG @ 0.5g/l. In order to ensure crop yields and reduce post-harvest losses, the use of pesticides, such as insecticides, has evolved from ancient times to become a crucial and strictly necessary agricultural component. Plant protection products are highly toxic to the pests they are intended to control. Utilising them is intended to lessen or get rid of pests totally. Sridhar and Sharma [13] found that, Flubendiamide 20 WG, provided significantly better protection up to 15 days after spraying to soybean crop from the semilooper (1.75 larvae / m row length [mrl]) and *S. litura*

Table 1. Comparative assessment of groundnut varieties JL-24 and Dh-256 for leaf-eating caterpillar incidence

Sl. No	Stage of the crop	<i>Spodoptera litura</i> /10 plants				<i>Thysanoplusia orichalcea</i> /10 plants				<i>Helicoverpa armigera</i> /10 plants			
		JL-24		Dh-256		JL-24		Dh-256		JL-24		Dh-256	
		UP	P	UP	P	UP	P	UP	P	UP	P	UP	P
1	Seedling	-	-	-	-	-	-	-	-	-	-	-	-
2	Vegetative	9.56	9.69	5.96	6.01	10.58	11.61	5.00	5.01	5.00	4.82	2.05	3.00
3	Flower initiation	25.98	24.68	16.98	18.50	18.18	19.00	15.31	15.29	12.00	12.05	9.24	9.75
4	Pegging and pod formation	42.25	13.95	30.00	9.75	23.91	11.24	18.00	9.28	17.43	8.15	12.31	7.40
5	Pod filling	29.86	10.92	18.56	5.66	10.32	5.92	7.01	2.00	9.30	3.21	7.20	2.53
6	Maturity	11.01	7.95	6.01	2.15	3.01	-	1.00	-	2.00	-	1.00	-
7	Harvesting	-	-	-	-	-	-	-	-	-	-	-	-
T_(Cal)		2.78		2.62		2.71		2.54		2.52		2.45	
T_(Tab) @ 0.05		2.44		2.44		2.44		2.44		2.44		2.44	
Sl. No	Stage of the crop	<i>Maruca vitrata</i> /10 plants				<i>Aproaerema modicella</i> /10 plants				<i>Spilarctia obliqua</i> /10 plants			
		JL-24		Dh-256		JL-24		Dh-256		JL-24		Dh-256	
		UP	P	UP	P	UP	P	UP	P	UP	P	UP	P
1	Seedling	-	-	-	-	-	-	-	-	-	-	-	-
2	Vegetative	-	-	-	-	-	-	-	-	-	-	-	-
3	Flower initiation	15.60	16.00	12.00	11.00	9.00	8.80	5.29	5.48	5.00	5.00	2.50	3.00
4	Pegging and pod formation	19.91	10.50	17.60	7.56	17.04	4.00	7.20	2.00	15.75	9.80	12.50	7.50
5	Pod filling	18.68	8.50	15.67	5.60	16.10	1.00	3.00	-	24.20	12.70	14.65	9.65
6	Maturity	12.57	4.57	8.10	1.58	2.00	-	-	-	30.50	19.50	24.00	15.62
7	Harvesting	-	-	-	-	-	-	-	-	35.75	25.50	29.21	19.51
T_(Cal)		2.73		2.64		3.32		2.48		2.85		2.67	
T_(Tab) @ 0.05		2.44		2.44		2.44		2.44		2.44		2.44	

UP: Unprotected; P: Protected

Table 2. Comparative incidence of natural enemies of insect pests on groundnut var. JL-24 and Dh-256

Sl. No.	Stage of the crop	<i>Metarhizium rileyi</i> (Cadavers/10 plants)				<i>Spodoptera liture</i> Nuclear polyhydrosis virus (Sl. NPV) (Infected larvae /10 plants)			
		JL-24		Dh-256		JL-24		Dh-256	
		UP	P	UP	P	UP	P	UP	P
1	Seedling	-	-	-	-	-	-	-	-
2	Vegetative	-	-	-	-	-	-	-	-
3	Flower initiation	21.00	10.51	15.50	8.24	10.40	8.91	7.00	4.82
4	Pegging and pod formation	42.50	6.08	24.50	5.00	13.20	1.89	9.23	0.75
5	Pod filling	5.04	-	2.00	-	2.04	-	1.00	-
6	Maturity	-	-	-	-	-	-	-	-
7	Harvesting	-	-	-	-	-	-	-	-
T_(Cal)		1.58		2.01		1.71		1.63	
T_(Tab) @ 0.05		2.44		2.44		2.44		2.44	

UP: Unprotected; P: Protected

(0.58 larvae / mrl) than untreated check (39.17 and 4.58 larvae / mrl). Tatagar et al. [14] who reported that Flubendiamide 20 WG @ 60 g a.i./ha was most effective in chilli against *S. litura* and *H. armigera* resulting in highest yield with lowest fruit damage. Sapekar et al. [15], noticed flubendiamide 39.35% SC @ 3 ml superior insecticide which gives maximum protection against semilooper (1.22 larvae / mrl), *S. litura* (2.53 larvae/mrl), *H. armigera* and bihar hairy caterpillar (0.81 larvae / mrl) as compared with untreated check (4.73, 2.40, 4.42 and 4.36 larvae/mrl).

Unprotected plots documented highest number of natural enemies. The present findings were corroborated by Manu, [16], who observed *M. rileyi* in vegetative stage (1.91 cadaver/mrl) and highest was reported at 70 DAS (17.50 cadaver/mrl). The occurrence of the illness coincides with the peak pest activity of the crop; *M. rileyi* infected *S. litura* was seen from 35 DAS and peaked at 65-70 DAS with 33.70%; insecticides were found to be extremely damaging to *M. rileyi* growth [17,18,19].

4. CONCLUSION

The pest incidence varied over the course of the crop phenological stages. The abundance of pests varied between the varieties in addition to the protected and unprotected plots. B Dh-256 tolerance to the main groundnut pests and hence it helps in limiting the use of particular insecticides for the targeted pest. Which in turn results in improved management and lower protection costs for farmers, who can then ensure higher yields and higher profits.

ACKNOWLEDGEMENT

We are grateful to Department of Agricultural Entomology, AICRP on Groundnut (UASD) and University of Agricultural Sciences, Dharwad for their unwavering support throughout the research.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Daphale KD, Mutkule DS, Ambilwade PP. Bio efficacy of bio-pesticides and insecticides against defoliators of kharif

groundnut. Biological Forum – An International Journal. 2022; 14(4a): 462-466.

2. Anonymous; 2022. Available:www.agricoop.gov.in
3. Waykule PK, Mutkule DS, Jadhav AS, Dhormare AP and Ingale AS. Bio-efficacy of different insecticides against gram pod borer (*Helicoverpa armigera* H.) on groundnut. Journal of Entomology and Zoology Studies. 2020 8(6): 354-358.
4. Pandiarajan J, Suganya T, Krishnan M. Gut resident microbes in groundnut pest *Amsacta albistriga* (Red Hairy Caterpillar). Current Research in Microbiology and Biotechnology. 2014;2:340-346.
5. Amin PW, Mohammad. Groundnut pest research at ICRISAT. Proceedings of the international workshop on groundnut. ICRISAT Center, Patancheru, Andhra Pradesh, India. 1980;158-166.
6. Biswas GC, Das GP. Insect and mite pests diversity in the oilseed crops ecosystems in Bangladesh. Bangladesh Journal of Zoology. 2011;39(2):232-235.
7. Srinivasa Rao M, Ranga Rao GV, Venkateswarlu B. Impact of climate change on insect pests. National Symposium on Climate Change and Rainfed Agriculture. February, 2010; 43-54.
8. Bijur S, Verma S. Sequential appearance of insect pests and natural enemy complex of on Pea crop. Indian Journal of Entomology. 1995;57(4):373-377.
9. Visalakshi M, Varma PK, Sekhar VC, Bharathalaxmi M, Manisha BL, Upendhar S. Studies on mycosis of *Metarhizium (Nomuraea) rileyi* on *Spodoptera frugiperda* infesting maize in Andhra Pradesh, India. Egyptian Journal of Biological Pest Control. 2020;30:1-10.
10. Bhattarai UR Li F, Katuwal Bhattarai M, Masoudi A, Wang D. Phototransduction and circadian entrainment are the key pathways in the signaling mechanism for the baculovirus induced tree-top disease in the lepidopteran larvae. Scientific Reports. 2018;8(1):17528.
11. Pal KK, Singh R, Singh AL. Handbook of groundnut varieties of India released during 2001-2021. ICAR-Directorate of Groundnut Research, P. B. 5, Junagadh, Gujarat, India. 2021;124.
12. Mohammad Saleem A, Gopalakrishna Naidu K, Tippannavar PS, Nadaf HL.

- Biophysical and biochemical mechanism of resistance to *Spodoptera litura* in groundnut (*Arachis hypogaea* L.). Journal of Entomology and Zoology Studies. 2019; 4:86-96.
13. Sridhar Y, Sharma AN. Flubendiamide, a novel insecticide for management of lepidopteron defoliators in soybean. Legume Research-An International Journal. 2015;38(4):551-554.
 14. Tatagar MH, Mohankumar HD, Shiva Prasad M, Mesta RK. Bio-efficacy of flubendiamide 20 WG against chilli fruit borers, *Helicoverpa armigera* (Hub.) and *Spodoptera litura* (Fb.). Karnataka. Journal of Agricultural Sciences. 2009;22(3): 579-581.
 15. Sapekar AS, Sonkamble MM, Matre YB. Bio-efficacy of Different Insecticides against of Major Defoliators on Soybean. International Journal of Current Microbiology and Applied Sciences. 2020; 11:2561-2569.
 16. Manu N. Crop loss estimation and management of leaf eating caterpillars in soybean. M. Sc. (Agri.) Thesis University of Agricultural Sciences, Dharwad, Karnataka, India; 2012.
 17. Patil RK. Ecofriendly approaches for the management of *Spodoptera litura* (F.) in groundnut. Ph.D. Thesis University of Agricultural Sciences, Dharwad, Karnataka, India; 2000.
 18. Anitha. Studies on the groundnut pod borers. M. Sc. (Agri.) Thesis Acharya N.G Ranga Agricultural University, Guntur, Andhra Pradesh, India; 1992.
 19. Raju CEP, Tayde AR. Field evaluation of selected insecticides and botanical against mustard aphid, *Lipaphis erysimi* (Kalt.) on mustard, *Brassica juncea* L. International Journal of Plant & Soil Science. 2022; 34(22):1188-1193.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here: