



Productivity and Economics of Hydroponic Fodders as Influenced by Harvesting Time

**Ayesha Ashique^{a++*}, A.B. Chorey^{b#}, M.M. Ganvir^{bt}
and V. Chaitanya^{a++}**

^a Department of Agronomy, Dr. Panjabrao Deshmukh Krishi Vidyapeeth, Akola (Maharashtra)-444104, India.

^b All India Coordinated Research Project on Dryland Agriculture, Dr. Panjabrao Deshmukh Krishi Vidyapeeth, Akola (Maharashtra)-444104, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JSRR/2024/v30i62040

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/115682>

Original Research Article

Received: 22/02/2024

Accepted: 26/04/2024

Published: 01/05/2024

ABSTRACT

An investigation was conducted at Maharashtra Animal and Fishery Science University, Panjabrao Deshmukh Krishi Vidyapeeth, Akola to study the Productivity and economics of hydroponic fodders as influenced by harvesting time at Department of Agronomy.

The investigation was carried out in a factorial randomized block design (FRBD) with three replicas. The study consisted of four different times of harvest (9th, 11th, 13th and 15th days of harvesting) and two fodder crops (maize and cowpea). The results revealed that the higher green fodder yield (26.64 kgm⁻²) and dry matter yield (3.06 kgm⁻²) were recorded on the 15th day of harvesting in

⁺⁺ PG Student;

[#] Chief Scientist;

[†] Agronomis;

*Corresponding author: E-mail: ayeshaashique18@gmail.com;

fodder crop Cowpea, whereas lower green fodder yield (18.58 kgm⁻²) and dry matter yield (2.28 kgm⁻²) were recorded on the 9th day of harvesting in Maize. In terms of economics, the 15th day of harvesting has recorded a higher gross monetary return, net monetary return, and benefit-cost ratio in Cowpea.

Keywords: Hydroponics; fodder; FRBD; B:C ratio; cowpea; harvesting time; antioxidants.

1. INTRODUCTION

In India Livestock contributes about 25% of the total agricultural GDP. A good fodder supply is required for better livestock production. The fodder, which is produced hydroponically, has a short growth period of 7–10 days, and only water is used to produce a fodder of shoot and root combinations [1].

As per the 20th Livestock Census 2019, the livestock population of the country is 535.78 million, including 192.49 million (35.94%) cattle, 109.85 million (20.45%) buffaloes, 74.26 million (13.87%) sheep, 148.88 million (27.80%) goats and 9.06 million pigs (1.69%). The increasing trend in the livestock population, along with the intensive rearing system, has resulted in increased demands for feed and fodder in the country. The land allocation for the cultivation of green fodder is limited to only 5% of the gross cropped area [2]. Presently, India is facing a net shortfall of 35.6% green fodder (Deo et al., 2021). The scarcity of adequate land, fertilizer, irrigation, and labour along with the unavailability of green fodder throughout the year, are major constraints for fodder production and sustainable dairy farming [3]. Further, fodder production is adversely affected by climate change through the erratic distribution of rainfall [4].

These problems can be overcome by hydroponics technology to produce green fodder for farm animals (Sneath and Mc Intosh, 2003; [5]. Hydroponic fodders are a rich source of antioxidants in form of β -carotene, Vitamins A, E and C [6] and rich in limiting amino acid, lysine. Hydroponic fodder is produced from cereal seeds like maize, barley, oats, millets and legumes like alfalfa, clover, horse gram and cowpea, which have high germinating capacity and fast-growing power. It is reported that hydroponic fodder increases digestibility of nutrients and milk yield. Hydroponics fodder is a good source of chlorophyll, which is very important with a poor-quality roughage-based diet during drought conditions. Hydroponics requires 1 to 2 litres of water to produce 1 kilogram of fodder, compared

to conventional fodder production, which requires 80–90 litres of water to grow 1 kilogram of green fodder. It saves about 95–97% of water and land required for fodder cultivation in a conventional system. [7]. Hence, present investigation was undertaken to study the hydroponic fodder production with the following objective:-

1. To evaluate the productivity and quality of fodder with different crops in hydroponic condition.
2. To study the optimum time of harvest for higher green fodder yield and quality of fodder.
3. To assess the economics of hydroponics fodder production in different treatments.

2. MATERIALS AND METHODS

The location of the experiment was at hydroponics facility of Maharashtra Animal and Fishery Science University, PDKV, Akola. The experiment was conducted using factorial RBD with two factors involving two crops and four times of harvesting, each repeated three times with a combination of a total 8 treatments. Each tray was made with vinyl fibre of dimensions 1.5 × 1 × 0.15 ft³, with three 1.5 mm holes designed at both sides. The seeds were dried under direct sunlight in open air one day before seed washing then it is washed in the bucket and the seeds were allowed to settle for 5 minutes where the light-weight floating seeds were taken out. After draining, the seeds and trays were washed with 0.1% sodium hypochlorite solution to control the formation of mould. The washed seeds are soaked in fresh water in a 1:2 proportion (1 part seeds and 2 parts water) in a container. Maize seeds were soaked for 24 hours, and cowpeas for 6-7 hours. After soaking, water is drained out and placed in a clean, wet gunny or cotton cloth airtight for sprouting. After 24 hours seeds were transferred into the tray. The automated fogging was adopted in the study, and the automation was set at one-minute for every two hours for maize and 30 seconds for every three hours for cowpea.

3. RESULTS AND DISCUSSION

The current study's findings, as well as relevant discussion, have been summarized under the following headings.

3.1 Effect of Crops

The data in Table 1 shows that the green fodder yield and dry matter yield were significantly influenced by the different fodder crops. The maximum green fodder yield of 23.48 kgm⁻² was recorded with Cowpea, which was significantly higher than Maize (22.17 kgm⁻²). The higher green biomass yield in Cowpea might be due to the higher yield parameters like fresh weight of shoot and roots. Similarly, the results revealed that the maximum dry matter yield was 2.92 kgm⁻² with Cowpea, which was significantly higher than Maize (2.50 kgm⁻²). The higher dry weight of Cowpea is directly related to total dry matter accumulation. These results are in conformity with the findings of Vennila [8]; Naik et al. [9]; Gnanaraj and Sundaram [10].

3.2 Effect of Harvesting Time

The data in Table 1 revealed that green fodder yield and dry matter yield were significantly influenced by the harvesting time. The maximum green fodder yield of 26.64 kgm⁻² was recorded with 15th day of harvesting, which was significantly superior from all the harvesting times. Biomass yield of the fodder crop depends on both external and internal factors [11]. The increase in dry matter accumulation in seedlings has increased fresh biomass yield of the hydroponically grown maize and cowpea with respect to time. The maximum dry matter yield 3.06 kgm⁻² was recorded in treatment 15th day of harvesting) and lowest on 9th day of harvesting

(2.28 kgm⁻²). Higher dry matter accumulation during different growth stages of Cowpea was mainly ascribed to increased photosynthetic ability and their partitioning to different parts with the advancement of age. These results are in conformity with the findings of Naik et al. [9].

3.3 Effect of Crops

Data presented in Table 2 revealed that the hydroponics fodder Cowpea has recorded a higher gross monetary return of Rs. 1570.26 m⁻² and was found to be significantly superior to the fodder Maize Rs. 1186 m⁻². This might be due to the fact that the market value of cowpea fodder is high as it is a protein crop. So, it has resulted in the highest gross return [12,13]. Cowpea has recorded higher net monetary return as Rs 477.76 m⁻² and found significantly superior over the Maize Rs. 281.05 m⁻². B:C ratio of fodder Cowpea (1.43) was found significantly superior over the fodder Maize (1.30). This might be due to higher market value of hydroponic cowpea fodder.

3.4 Effect of Harvest Time

The data presented in Table 2 further revealed that the harvesting time at H₄ (15th day of harvesting) recorded higher gross monetary return, net monetary return and benefit:cost ratio [14]. 15th day of harvesting Rs.1610m⁻² and was found significantly superior over the others. As the age advances, the total green fodder yield increases, which in turn increases the gross monetary return with harvesting on the 15th day after sowing. Higher net monetary return was observed on 15th day of harvesting Rs.579.79m⁻². This might be due to the fact that as the age advances the total green fodder increased and resulted in increased gross returns, but the

Table 1. Green fodder yield and Dry matter yield (kg m⁻²) of hydroponically grown fodder crops as influenced by the time of harvest

Treatments	Greenfodder yield (kgm ⁻²)	Dry matter yield (kgm ⁻²)
C1- Maize	22.17	0.23
C2-Cowpea	23.48	0.27
SE (m) ±	0.09	0.01
CD at 1%	0.27	0.03
H1- Harvesting at 9 th day	18.58	0.22
H2-Harvesting at 11 th day	20.93	0.23
H3-Harvesting at 13 th day	25.12	0.26
H4-Harvesting at 15 th day	26.64	0.28
SE(m) ±	0.46	0.01
CD at 1%	1.41	0.04

Table 2. Economics of hydroponically grown fodder crops as influenced by the time of harvest

Treatments	GMR Rs m ⁻²	NMR Rs m ⁻²	Cost of Cultivation Rs m ⁻²	B:C Ratio
Factor A- Crop				
C1- Maize	1186.28	281.05	905.21	1.30
C2-Cowpea	1570.26	477.76	1092.50	1.43
SE (m) ±	19.08	19.08	-	-
CD at 1%	57.86	57.86	-	-
Factor B – Harvesting time				
H1- Harvesting at 9 th day	1117.97	151.14	966.80	1.15
H2-Harvesting at 11 th day	1265.66	277.48	988.17	1.27
H3-Harvesting at 13 th day	1518.73	509.19	1009.54	1.49
H4-Harvesting at 15 th day	1610.70	579.80	1030.90	1.55
SE(m) ±	26.98	26.98	-	-
CD at 1%	81.83	81.83	-	-
Interaction				
SE (m) ±	38.15	38.15	-	-
CD at 1%	115.73	115.72	-	-
GM	1378.27	379.40	998.85	1.37

additional savings in seed and labour cost resulted marginal increase in higher net returns. Higher benefit cost ratio was observed on 15th day of harvesting (1.55) and lowest on 9th day of harvesting (1.15).

4. CONCLUSIONS

From the above study, it was observed that higher green fodder yield and dry matter yield was recorded in cowpea as compared to maize, whereas in harvesting time higher green fodder yield and dry matter yield were recorded at 15th day of harvesting and was superior over other day of harvesting.

In terms of economics, it can be concluded that cowpea has a higher gross return, net return and B:C ratio. Harvesting the crop at 15th day has resulted in higher return but 13th day of harvesting time was found more economical due to higher increase in the rate of gross and net monetary return.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Al-Juthery HW, Al-Fadhly JT, Ali EA. HM, Al-Tae RAHG. Role of some nanofertilizers and atonikin maximizing for production of hydroponically-grown barley fodder. *Int. J. Agric. Stat. Sci.* 2019;15:565-570.
2. Shit N. Hydroponic fodder production: an alternative technology for sustainable livestock production in India. *Exploratory Animal & Medical Research.* 2019;9(2):108-119.
3. Hazra CR. Feed and forage resources for sustainable livestock development. *Range Management and Agroforestry.* 2014;35(1):1-14.
4. Gunasekaran S, Bandeswaran C, Valli C. Effect of different nutrient solutions on the biomass yield and nutritive value of low-cost hydroponic fodder maize. *Indian Journal of Animal Sciences.* 2017;87(11):1418-1419.
5. Naik PK, Dhawaskar BD, Fatarpekar DD, Chakurkar EB, Swain BKSingh NP. Nutrient changes with the growth of hydroponics cowpea (*Vigna unguiculata*) sprouts. *Indian Journal of Animal Nutrition.* 2016;33(3):357-359.
6. Gaikwad DJ, Kalasare RS, Jaswanth DS, Sravani PS, Pramanic S. Morpho-Physiological Responses Associated with Hydroponically Grown Maize Fodder. *Indian Journal of Natural Sciences.* 2022;13(72):43228-43232.
7. Al-karaki GN, Al-hashimi M. Green fodder production and water use efficiency of some forage crops under hydroponic condition. *Intern. Schol. Res. Network.* 2012:1-5.
8. Vennila C. Productivity, nutritive value, growth rate, biomass yield and economics of different hydroponic green fodders

- for livestock. International Journal of Livestock Research. 2018;8(9): 261-270.
9. Naik PK, Swain BK, Singh NP. Production and utilisation of hydroponics fodder. Indian Journal of Animal Nutrition. 2015;32(1):1-9.
 10. Gnanaraj PT, Sundaram SM. Productivity and water use efficiency and nutritional composition of yellow maize fodder under hydroponic condition. Journal of Pharmacognosy and Phytochemistry. 2020;9(3):243-246.
 11. Bekuma A. Nutritional Benefit and Economic Value of Hydroponics Fodder Production Technology in Sustainable Livestock Production Against Climate Change-A Mini-Review. Advances in Applied Sciences. 2019; 4(1):23.
 12. Ganguly S, Lalitha BS, Raddy G, Khan S. Influence of Combinations of Different Fodder Crops on Growth and Biomass Yield under Hydroponic Condition. Int. J. Curr. Microbiol. App. Sci. 2020;9(9): 498-509.
 13. Jolad R, Sivakumar SD, Babu C, Sriharan N. Performance of Different Crops under Hydroponics Fodder Production System. Madras Agricultural Journal. 2018; 105(1-3):50-55.
 14. Kumari S, Pradhan P, Yadav R, Kumar S. Hydroponic techniques: A soilless cultivation in agriculture. Journal of pharmacognosy and phytochemistry. 2018;7(1):1886-1891.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/115682>